



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 25, 2023 – 01:40 PM EDT

PDB ID : 5VB0  
Title : Crystal structure of fosfomycin resistance protein FosA3  
Authors : Klontz, E.; Guenther, S.; Silverstein, Z.; Sundberg, E.  
Deposited on : 2017-03-28  
Resolution : 2.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	144	 79% 16% ..
1	B	144	 81% 10% 8%
1	C	144	 8% 76% 23% .
1	D	144	 3% 84% 8% 8%
1	E	144	 85% 8% 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	144	 89% 7%
1	G	144	 82% 10% 8%
1	H	144	 83% 13%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fosfomycin resistance protein FosA3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	139	1101	701	193	201	6	0	0	0
1	B	133	1047	670	177	194	6	0	0	0
1	C	143	1119	710	198	205	6	0	0	0
1	D	132	1039	666	176	191	6	0	0	0
1	E	133	1037	664	173	194	6	0	0	0
1	F	139	1101	702	194	199	6	0	0	0
1	G	132	1033	663	173	191	6	0	0	0
1	H	140	1109	706	195	202	6	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	HIS	-	expression tag	UNP D7UQM0
A	140	HIS	-	expression tag	UNP D7UQM0
A	141	HIS	-	expression tag	UNP D7UQM0
A	142	HIS	-	expression tag	UNP D7UQM0
A	143	HIS	-	expression tag	UNP D7UQM0
A	144	HIS	-	expression tag	UNP D7UQM0
B	139	HIS	-	expression tag	UNP D7UQM0
B	140	HIS	-	expression tag	UNP D7UQM0
B	141	HIS	-	expression tag	UNP D7UQM0
B	142	HIS	-	expression tag	UNP D7UQM0
B	143	HIS	-	expression tag	UNP D7UQM0
B	144	HIS	-	expression tag	UNP D7UQM0
C	139	HIS	-	expression tag	UNP D7UQM0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	140	HIS	-	expression tag	UNP D7UQM0
C	141	HIS	-	expression tag	UNP D7UQM0
C	142	HIS	-	expression tag	UNP D7UQM0
C	143	HIS	-	expression tag	UNP D7UQM0
C	144	HIS	-	expression tag	UNP D7UQM0
D	139	HIS	-	expression tag	UNP D7UQM0
D	140	HIS	-	expression tag	UNP D7UQM0
D	141	HIS	-	expression tag	UNP D7UQM0
D	142	HIS	-	expression tag	UNP D7UQM0
D	143	HIS	-	expression tag	UNP D7UQM0
D	144	HIS	-	expression tag	UNP D7UQM0
E	139	HIS	-	expression tag	UNP D7UQM0
E	140	HIS	-	expression tag	UNP D7UQM0
E	141	HIS	-	expression tag	UNP D7UQM0
E	142	HIS	-	expression tag	UNP D7UQM0
E	143	HIS	-	expression tag	UNP D7UQM0
E	144	HIS	-	expression tag	UNP D7UQM0
F	139	HIS	-	expression tag	UNP D7UQM0
F	140	HIS	-	expression tag	UNP D7UQM0
F	141	HIS	-	expression tag	UNP D7UQM0
F	142	HIS	-	expression tag	UNP D7UQM0
F	143	HIS	-	expression tag	UNP D7UQM0
F	144	HIS	-	expression tag	UNP D7UQM0
G	139	HIS	-	expression tag	UNP D7UQM0
G	140	HIS	-	expression tag	UNP D7UQM0
G	141	HIS	-	expression tag	UNP D7UQM0
G	142	HIS	-	expression tag	UNP D7UQM0
G	143	HIS	-	expression tag	UNP D7UQM0
G	144	HIS	-	expression tag	UNP D7UQM0
H	139	HIS	-	expression tag	UNP D7UQM0
H	140	HIS	-	expression tag	UNP D7UQM0
H	141	HIS	-	expression tag	UNP D7UQM0
H	142	HIS	-	expression tag	UNP D7UQM0
H	143	HIS	-	expression tag	UNP D7UQM0
H	144	HIS	-	expression tag	UNP D7UQM0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mn 1 1	0	0
2	B	1	Total Mn 1 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0
2	E	1	Total Mn 1 1	0	0
2	F	1	Total Mn 1 1	0	0
2	G	1	Total Mn 1 1	0	0
2	H	1	Total Mn 1 1	0	0


- Molecule 3 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ni 2 2	0	0
3	C	2	Total Ni 2 2	0	0
3	F	1	Total Ni 1 1	0	0
3	H	1	Total Ni 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Fosfomycin resistance protein FosA3




- Molecule 1: Fosfomycin resistance protein FosA3




- Molecule 1: Fosfomycin resistance protein FosA3




- Molecule 1: Fosfomycin resistance protein FosA3




- Molecule 1: Fosfomycin resistance protein FosA3








- Molecule 1: Fosfomycin resistance protein FosA3



- Molecule 1: Fosfomycin resistance protein FosA3



- Molecule 1: Fosfomycin resistance protein FosA3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.61Å 87.61Å 357.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 2.69 29.75 – 2.69	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.75-2.69) 100.0 (29.75-2.69)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 2.68Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, $R_{free}$	0.205 , 0.249 0.212 , 0.258	Depositor DCC
$R_{free}$ test set	1920 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	57.5	Xtrriage
Anisotropy	0.715	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8600	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	4/1133 (0.4%)	0.59	1/1538 (0.1%)
1	B	0.34	0/1074	0.60	0/1456
1	C	0.39	0/1152	0.69	1/1566 (0.1%)
1	D	0.28	0/1066	0.50	0/1445
1	E	0.44	2/1064 (0.2%)	0.68	3/1445 (0.2%)
1	F	0.33	0/1133	0.60	1/1537 (0.1%)
1	G	0.32	0/1060	0.57	0/1438
1	H	0.42	1/1141 (0.1%)	0.60	1/1547 (0.1%)
All	All	0.39	7/8823 (0.1%)	0.61	7/11972 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	ARG	CZ-NH1	-7.95	1.22	1.33
1	A	56	ARG	NE-CZ	-7.92	1.22	1.33
1	A	56	ARG	CZ-NH2	-6.28	1.24	1.33
1	E	132	LYS	CD-CE	-6.01	1.36	1.51
1	H	75	GLU	CD-OE1	-5.99	1.19	1.25

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	132	LYS	CA-CB-CG	10.19	135.82	113.40
1	E	132	LYS	CB-CG-CD	7.69	131.58	111.60
1	H	19	LEU	CA-CB-CG	-7.62	97.78	115.30
1	F	129	ARG	CG-CD-NE	6.66	125.80	111.80
1	E	132	LYS	CG-CD-CE	-5.57	95.18	111.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1101	0	1034	14	0
1	B	1047	0	1004	12	0
1	C	1119	0	1041	28	0
1	D	1039	0	1000	10	0
1	E	1037	0	982	5	0
1	F	1101	0	1041	10	0
1	G	1033	0	989	9	0
1	H	1109	0	1048	16	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	2	0	0	0	0
3	C	2	0	0	0	0
3	F	1	0	0	0	0
3	H	1	0	0	0	0
All	All	8600	0	8139	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:TRP:HB3	1:B:105:LEU:HD11	1.55	0.88
1:C:15:LEU:HG	1:C:19:LEU:HD11	1.57	0.87
1:B:53:GLU:O	1:B:56:ARG:NH2	2.09	0.85
1:C:98:GLU:OE2	1:C:101:SER:N	2.15	0.80
1:F:129:ARG:O	1:F:129:ARG:HD2	1.88	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/144 (94%)	132 (98%)	3 (2%)	0	100	100
1	B	129/144 (90%)	126 (98%)	3 (2%)	0	100	100
1	C	141/144 (98%)	134 (95%)	7 (5%)	0	100	100
1	D	128/144 (89%)	125 (98%)	3 (2%)	0	100	100
1	E	129/144 (90%)	127 (98%)	1 (1%)	1 (1%)	19	40
1	F	135/144 (94%)	134 (99%)	1 (1%)	0	100	100
1	G	128/144 (89%)	125 (98%)	3 (2%)	0	100	100
1	H	136/144 (94%)	133 (98%)	3 (2%)	0	100	100
All	All	1061/1152 (92%)	1036 (98%)	24 (2%)	1 (0%)	51	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	57	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/117 (96%)	108 (96%)	4 (4%)	35	61
1	B	107/117 (92%)	104 (97%)	3 (3%)	43	70
1	C	112/117 (96%)	110 (98%)	2 (2%)	59	81
1	D	106/117 (91%)	105 (99%)	1 (1%)	78	91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	105/117 (90%)	103 (98%)	2 (2%)	57	80
1	F	112/117 (96%)	110 (98%)	2 (2%)	59	81
1	G	105/117 (90%)	103 (98%)	2 (2%)	57	80
1	H	113/117 (97%)	111 (98%)	2 (2%)	59	81
All	All	872/936 (93%)	854 (98%)	18 (2%)	53	78

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	106	ASP
1	H	106	ASP
1	H	29	ARG
1	C	106	ASP
1	F	129	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	121	GLN
1	H	24	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	139/144 (96%)	-0.15	1 (0%) 87 88	44, 57, 80, 98	0
1	B	133/144 (92%)	-0.05	2 (1%) 73 74	41, 56, 86, 120	0
1	C	143/144 (99%)	0.38	11 (7%) 13 11	48, 75, 104, 113	0
1	D	132/144 (91%)	0.13	4 (3%) 50 49	46, 63, 93, 107	0
1	E	133/144 (92%)	-0.03	2 (1%) 73 74	45, 59, 82, 100	0
1	F	139/144 (96%)	-0.11	1 (0%) 87 88	47, 61, 88, 100	0
1	G	132/144 (91%)	0.08	3 (2%) 60 60	48, 63, 84, 102	0
1	H	140/144 (97%)	0.23	4 (2%) 51 51	46, 70, 94, 109	0
All	All	1091/1152 (94%)	0.06	28 (2%) 56 55	41, 62, 92, 120	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	99	GLY	6.0
1	B	94	ASP	5.3
1	D	94	ASP	4.7
1	D	1	MET	4.5
1	G	54	GLN	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NI	C	202	1/1	0.87	0.21	87,87,87,87	0
2	MN	D	201	1/1	0.91	0.20	54,54,54,54	0
3	NI	H	202	1/1	0.92	0.08	89,89,89,89	0
2	MN	F	201	1/1	0.93	0.22	56,56,56,56	0
3	NI	A	202	1/1	0.93	0.21	81,81,81,81	0
3	NI	F	202	1/1	0.94	0.05	85,85,85,85	0
3	NI	C	203	1/1	0.94	0.04	87,87,87,87	0
2	MN	B	201	1/1	0.96	0.24	52,52,52,52	0
3	NI	A	203	1/1	0.96	0.03	99,99,99,99	0
2	MN	E	201	1/1	0.97	0.30	59,59,59,59	0
2	MN	A	201	1/1	0.97	0.21	50,50,50,50	0
2	MN	C	201	1/1	0.98	0.26	57,57,57,57	0
2	MN	G	201	1/1	0.98	0.27	60,60,60,60	0
2	MN	H	201	1/1	0.98	0.32	62,62,62,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.