

Full wwPDB X-ray Structure Validation Report (i)

May 26, 2020 - 03:30 am BST

PDB ID	:	2VH5
Title	:	CRYSTAL STRUCTURE OF HRAS(G12V) - ANTI-RAS FV (disulfide free
		mutant) COMPLEX
Authors	:	Tanaka, T.; Williams, R.L.; Rabbitts, T.H.
Deposited on		
$\operatorname{Resolution}$:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


• :	4.02b-467
:	1.8.5 (274361), CSD as541be (2020)
:	NOT EXECUTED
:	NOT EXECUTED
:	1.1.7(2018)
; ;	20191225.v01 (using entries in the PDB archive December 25 th 2019)
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.11
	l :) : 5 : 5 :) :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069(2.70-2.70)
Sidechain outliers	138945	3069(2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	Н	114	83%	16%	•
2	L	104	67% 27%		6%
3	R	166	81%	16%	•••



2VH5

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 3081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ANTI-RAS FV HEAVY CHAIN.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Н	114	Total	C EGQ	N 154	0	S o	0	0	0
			893	568	154	169	2			

• Molecule 2 is a protein called ANTI-RAS FV LIGHT CHAIN.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	L	104	Total 785	C 494	N 128	O 160	${ m S} { m 3}$	0	0	0

• Molecule 3 is a protein called GTPASE HRAS.

Mol	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace
3	R	166	Total 1325	C 827	N 228	O 263	S 7	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
R	12	VAL	GLY	engineered mutation	UNP P01112

• Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Н	2	Total Zn 2 2	0	0
4	R	1	Total Zn 1 1	0	0

• Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf
F	D	1	Total	С	Ν	Ο	Р	0	0
5	п	T	32	10	5	14	3	0	0

• Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	R	1	Total Mg 1 1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	Н	10	Total O 10 10	0	0
7	L	9	Total O 9 9	0	0
7	R	23	TotalO2323	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ANTI-RAS FV HEAVY CHAIN





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	75.86Å 85.40 Å 63.08 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.18 - 2.70	Depositor
% Data completeness	99.1 (42.18-2.70)	Depositor
(in resolution range)	55.1 (42.10 2.10)	Depositor
R_{merge}	0.13	Depositor
R _{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.211 , 0.291	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3081	wwPDB-VP
Average B, all atoms $(Å^2)$	40.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	Н	0.52	0/913	0.64	0/1235
2	L	0.50	0/801	0.69	1/1088~(0.1%)
3	R	0.60	1/1344~(0.1%)	0.69	1/1815~(0.1%)
Al	l All	0.55	1/3058~(0.0%)	0.68	2/4138~(0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	R	31	GLU	CB-CG	5.97	1.63	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	R	108	ASP	CB-CG-OD1	7.01	124.61	118.30
2	L	13	LEU	CA-CB-CG	5.93	128.93	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Н	893	0	874	10	1
2	L	785	0	771	20	0

Continued on next page...



Mol	•	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	R	1325	0	1300	20	3
4	Η	2	0	0	0	0
4	R	1	0	0	0	0
5	R	32	0	12	2	0
6	R	1	0	0	0	0
7	Н	10	0	0	1	0
7	L	9	0	0	0	0
7	R	23	0	0	1	0
All	All	3081	0	2957	49	3

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A. (Interatomic	Clash
Atom-1	Atom-2	${ m distance}~({ m \AA})$	overlap (Å)
2:L:10:PRO:O	2:L:104:THR:HG22	1.58	1.01
3:R:112:VAL:HG22	3:R:140:PRO:HG2	1.69	0.73
3:R:108:ASP:OD1	3:R:166:HIS:NE2	2.28	0.67
2:L:81:GLN:HB3	2:L:82:PRO:HD2	1.76	0.66
3:R:30:ASP:OD1	3:R:31:GLU:OE2	2.15	0.64
3:R:20:THR:O	3:R:24:ILE:HG12	1.98	0.64
2:L:21:VAL:HA	2:L:22:THR:HB	1.80	0.62
2:L:63:ARG:CZ	2:L:81:GLN:HG3	2.30	0.60
2:L:41:LYS:NZ	2:L:83:GLU:HB2	2.17	0.59
3:R:111:MET:HB3	3:R:139:ILE:HG21	1.86	0.58
1:H:83:MET:HE1	1:H:111:VAL:HG21	1.87	0.57
1:H:22:ALA:HB2	1:H:96:VAL:HG21	1.88	0.56
1:H:99:GLY:HA3	7:H:2006:HOH:O	2.06	0.56
3:R:32:TYR:CD1	5:R:1167:GTP:H5"	2.42	0.55
3:R:1:MET:O	3:R:1:MET:SD	2.66	0.54
1:H:34:MET:HB3	1:H:79:LEU:HD22	1.89	0.53
2:L:32:SER:HA	2:L:70:GLY:H	1.75	0.50
2:L:22:THR:O	2:L:22:THR:HG22	2.13	0.49
2:L:5:GLN:HB3	2:L:28:SER:HB3	1.94	0.49
2:L:22:THR:H	2:L:76:THR:HA	1.77	0.49
2:L:31:ILE:HG21	2:L:92:GLN:HG3	1.92	0.49
2:L:41:LYS:HZ3	2:L:83:GLU:HB2	1.76	0.49
2:L:92:GLN:NE2	2:L:95:MET:H	2.11	0.49
2:L:93:SER:HA	2:L:98:MET:HG2	1.95	0.48
3:R:98:GLU:OE1	3:R:102:ARG:NH2	2.46	0.47

Continued on next page...



		Interatomic	Clash
Atom-1	Atom-2	$distance (m \AA)$	overlap (Å)
1:H:22:ALA:CB	1:H:96:VAL:HG21	2.44	0.47
1:H:39:GLN:HE22	2:L:40:GLN:HE22	1.62	0.47
1:H:43:LYS:HA	1:H:43:LYS:HD2	1.78	0.47
1:H:102:PHE:N	1:H:102:PHE:CD1	2.83	0.46
3:R:68:ARG:HG2	3:R:71:TYR:OH	2.15	0.46
2:L:33:SER:O	2:L:52:SER:HA	2.15	0.46
1:H:6:GLU:HA	1:H:21:SER:O	2.17	0.45
3:R:154:ASP:HB2	7:R:2019:HOH:O	2.17	0.44
1:H:41:PRO:HA	1:H:42:GLY:HA2	1.74	0.44
3:R:72:MET:O	3:R:104:LYS:HD3	2.18	0.44
2:L:80:LEU:HD23	2:L:84:ASP:HB2	1.98	0.44
3:R:84:ILE:HD12	3:R:123:ARG:HB3	2.00	0.44
3:R:41:ARG:HA	3:R:53:LEU:O	2.17	0.44
3:R:108:ASP:CG	3:R:166:HIS:CD2	2.91	0.44
3:R:29:VAL:O	3:R:29:VAL:HG23	2.18	0.44
3:R:111:MET:O	3:R:139:ILE:HB	2.19	0.42
2:L:5:GLN:HA	2:L:5:GLN:HE21	1.84	0.42
2:L:41:LYS:HE2	2:L:83:GLU:O	2.20	0.42
3:R:117:LYS:HG2	5:R:1167:GTP:C6	2.55	0.41
3:R:87:THR:HG23	3:R:124:THR:OG1	2.20	0.41
2:L:60:VAL:HA	2:L:61:PRO:HD3	1.81	0.41
2:L:4:ILE:HG22	2:L:29:GLN:HE21	1.86	0.40
3:R:84:ILE:HD11	3:R:118:CYS:HA	2.03	0.40
3:R:108:ASP:OD1	3:R:166:HIS:CD2	2.74	0.40

Continued from previous page...

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:1:GLU:OE2	3:R:166:HIS:NE2[4_556]	1.97	0.23
3:R:27:HIS:CE1	3:R:98:GLU:OE2[4_556]	2.05	0.15
3:R:31:GLU:OE2	3:R:107:ASP:OD2[4_556]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	Η	112/114~(98%)	108~(96%)	4 (4%)	0	100	100
2	L	102/104~(98%)	96 (94%)	4 (4%)	2(2%)	7	19
3	R	164/166~(99%)	156~(95%)	8 (5%)	0	100	100
All	All	378/384 (98%)	360~(95%)	16 (4%)	2(0%)	29	54

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	22	THR
2	L	32	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Н	94/94~(100%)	89~(95%)	5 (5%)	22 48
2	L	90/90~(100%)	$81 \ (90\%)$	9 (10%)	7 18
3	R	145/145~(100%)	138~(95%)	7 (5%)	25 53
All	All	329/329~(100%)	308 (94%)	21 (6%)	17 39

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	5	LEU
1	Н	12	VAL
1	Н	20	LEU
1	Н	89	GLU
1	Н	102	PHE
2	L	5	GLN
2	L	11	SER
2	L	12	SER

Continued on next page...



Mol	Chain	Res	Type
2	L	13	LEU
2	L	29	GLN
2	L	35	LEU
2	L	44	GLU
2	L	76	THR
2	L	105	LYS
3	R	1	MET
3	R	29	VAL
3	R	31	GLU
3	R	52	LEU
3	R	70	GLN
3	R	102	ARG
3	R	107	ASP

Continued from previous page...

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	Н	77	ASN
2	L	5	GLN
2	L	29	GLN
2	L	40	GLN
2	L	92	GLN
3	R	70	GLN
3	R	95	GLN
3	R	129	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
5	GTP	R	1167	6	$26,\!34,\!34$	1.01	2 (7%)	$33,\!54,\!54$	1.70	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	R	1167	6	-	1/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
5	R	1167	GTP	C6-N1	2.87	1.38	1.33
5	R	1167	GTP	C2-N1	2.13	1.39	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
5	R	1167	GTP	N3-C2-N1	-5.37	120.06	127.22
5	R	1167	GTP	C2-N3-C4	3.82	119.72	115.36
5	R	1167	GTP	PB-O3B-PG	-3.33	121.39	132.83
5	R	1167	GTP	C5-C6-N1	-2.85	119.54	123.43
5	R	1167	GTP	C6-N1-C2	2.74	120.28	115.93
5	R	1167	GTP	PA-O3A-PB	-2.07	125.71	132.83
5	R	1167	GTP	N2-C2-N1	2.04	120.43	117.25

There are no chirality outliers.

All (1) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
5	R	1167	GTP	PA-O3A-PB-O1B

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	R	$11\overline{67}$	GTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

