

# wwPDB X-ray Structure Validation Summary Report (i)

### May 15, 2020 – 03:47 am BST

PDB ID : 6VH5

Title: Crystal structure of prephenate dehydratase from brucella melitensis biovar

abortus 2308 in complex with phenylalanine

Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2020-01-09

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove) roteins) : Engh & Huber (2001)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries},  ext{resolution range}( ext{Å})) \end{aligned}$
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	308	83%	8%	9%
1	В	308	85%	6%	9%
1	С	308	86%	5%	9%
1	D	308	82%	9%	8%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Prephenate dehydratase: Amino acid-binding ACT.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	281	Total	С	N	О	S	0	0	0
1	A	201	2146	1358	373	404	11	0	U	0
1	В	280	Total	С	N	О	S	0	0	0
1	Б		2094	1328	361	395	10	U	U	
1	С	279	Total	С	N	О	S	0	0	0
1		219	2084	1322	359	393	10	0	U	
1	D	282	Total	С	N	О	S	0	1	0
1		282	2119	1346	365	398	10	0	1	

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q2YPM5
A	-19	ALA	-	expression tag	UNP Q2YPM5
A	-18	HIS	-	expression tag	UNP Q2YPM5
A	-17	HIS	_	expression tag	UNP Q2YPM5
A	-16	HIS	-	expression tag	UNP Q2YPM5
A	-15	HIS	_	expression tag	UNP Q2YPM5
A	-14	HIS	_	expression tag	UNP Q2YPM5
A	-13	HIS	-	expression tag	UNP Q2YPM5
A	-12	MET	_	expression tag	UNP Q2YPM5
A	-11	GLY	-	expression tag	UNP Q2YPM5
A	-10	THR	-	expression tag	UNP Q2YPM5
A	-9	LEU	-	expression tag	UNP Q2YPM5
A	-8	GLU	-	expression tag	UNP Q2YPM5
A	-7	ALA	_	expression tag	UNP Q2YPM5
A	-6	GLN	_	expression tag	UNP Q2YPM5
A	-5	THR	_	expression tag	UNP Q2YPM5
A	-4	GLN	=	expression tag	UNP Q2YPM5
A	-3	GLY	-	expression tag	UNP Q2YPM5
A	-2	PRO		expression tag	UNP Q2YPM5
A	-1	GLY	-	expression tag	UNP Q2YPM5
A	0	SER	_	expression tag	UNP Q2YPM5



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-20	MET	_	expression tag	UNP Q2YPM5
В	-19	ALA	_	expression tag	UNP Q2YPM5
В	-18	HIS	-	expression tag	UNP Q2YPM5
В	-17	HIS	_	expression tag	UNP Q2YPM5
В	-16	HIS	-	expression tag	UNP Q2YPM5
В	-15	HIS	_	expression tag	UNP Q2YPM5
В	-14	HIS	_	expression tag	UNP Q2YPM5
В	-13	HIS	-	expression tag	UNP Q2YPM5
В	-12	MET	-	expression tag	UNP Q2YPM5
В	-11	GLY	-	expression tag	UNP Q2YPM5
В	-10	THR	-	expression tag	UNP Q2YPM5
В	-9	LEU	-	expression tag	UNP Q2YPM5
В	-8	GLU	-	expression tag	UNP Q2YPM5
В	-7	ALA	-	expression tag	UNP Q2YPM5
В	-6	GLN	-	expression tag	UNP Q2YPM5
В	-5	THR	-	expression tag	UNP Q2YPM5
В	-4	GLN	-	expression tag	UNP Q2YPM5
В	-3	GLY	-	expression tag	UNP Q2YPM5
В	-2	PRO	-	expression tag	UNP Q2YPM5
В	-1	GLY	-	expression tag	UNP Q2YPM5
В	0	SER	-	expression tag	UNP Q2YPM5
С	-20	MET	_	expression tag	UNP Q2YPM5
С	-19	ALA	_	expression tag	UNP Q2YPM5
С	-18	HIS	_	expression tag	UNP Q2YPM5
С	-17	HIS	_	expression tag	UNP Q2YPM5
С	-16	HIS	_	expression tag	UNP Q2YPM5
С	-15	HIS	_	expression tag	UNP Q2YPM5
С	-14	HIS	_	expression tag	UNP Q2YPM5
С	-13	HIS	_	expression tag	UNP Q2YPM5
С	-12	MET	_	expression tag	UNP Q2YPM5
С	-11	GLY	_	expression tag	UNP Q2YPM5
С	-10	THR	_	expression tag	UNP Q2YPM5
С	-9	LEU	-	expression tag	UNP Q2YPM5
С	-8	GLU	_	expression tag	UNP Q2YPM5
С	-7	ALA	-	expression tag	UNP Q2YPM5
С	-6	GLN	-	expression tag	UNP Q2YPM5
С	-5	THR	-	expression tag	UNP Q2YPM5
С	-4	GLN	-	expression tag	UNP Q2YPM5
С	-3	GLY	-	expression tag	UNP Q2YPM5
С	-2	PRO	-	expression tag	UNP Q2YPM5
С	-1	GLY	-	expression tag	UNP Q2YPM5
С	0	SER	-	expression tag	UNP Q2YPM5




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MET	-	expression tag	UNP Q2YPM5
D	-19	ALA	-	expression tag	UNP Q2YPM5
D	-18	HIS	_	expression tag	UNP Q2YPM5
D	-17	HIS	-	expression tag	UNP Q2YPM5
D	-16	HIS	_	expression tag	UNP Q2YPM5
D	-15	HIS	_	expression tag	UNP Q2YPM5
D	-14	HIS	_	expression tag	UNP Q2YPM5
D	-13	HIS	_	expression tag	UNP Q2YPM5
D	-12	MET	_	expression tag	UNP Q2YPM5
D	-11	GLY	_	expression tag	UNP Q2YPM5
D	-10	THR	_	expression tag	UNP Q2YPM5
D	-9	LEU	_	expression tag	UNP Q2YPM5
D	-8	GLU	_	expression tag	UNP Q2YPM5
D	-7	ALA	-	expression tag	UNP Q2YPM5
D	-6	GLN	-	expression tag	UNP Q2YPM5
D	-5	THR	-	expression tag	UNP Q2YPM5
D	-4	GLN	-	expression tag	UNP Q2YPM5
D	-3	GLY	=	expression tag	UNP Q2YPM5
D	-2	PRO	-	expression tag	UNP Q2YPM5
D	-1	GLY	=	expression tag	UNP Q2YPM5
D	0	SER	-	expression tag	UNP Q2YPM5

 $\bullet$  Molecule 2 is PHENYLALANINE (three-letter code: PHE) (formula:  $\mathrm{C_9H_{11}NO_2}).$ 



$\mathbf{Mol}$	Chain	Residues	Atoms				ZeroOcc	$\mathbf{AltConf}$
2	A	1	Total 12	C 9	N 1	O 2	0	0






 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C N O 12 9 1 2	0	0
2	С	1	Total C N O 12 9 1 2	0	0
2	D	1	Total C N O 12 9 1 2	0	0

 $\bullet$  Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\mathrm{C_2H_6O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	В	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	52	Total O 52 52	0	0
4	В	50	Total O 50 50	0	0
4	С	29	Total O 29 29	0	0
4	D	37	Total O 37 37	0	0




Chain D:


# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prephenate dehydratase: Amino acid-binding ACT









# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	226.95Å 61.25Å 120.05Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 112.95° 90.00°	Depositor
Resolution (Å)	41.41 - 2.40	Depositor
Resolution (A)	41.41 - 2.40	EDS
% Data completeness	98.2 (41.41-2.40)	Depositor
(in resolution range)	98.5 (41.41-2.40)	EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.79 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.17.1	Depositor
D D.	0.207 , 0.251	Depositor
$R, R_{free}$	0.207 , $0.251$	DCC
$R_{free}$ test set	1878 reflections $(3.18\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.4	Xtriage
Anisotropy	0.919	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.30 , 45.8	EDS
L-test for twinning <sup>2</sup>	$ < L > = 0.51, < L^2> = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 19.78% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.35	0/2189	0.54	0/2975	
1	В	0.34	0/2136	0.53	0/2911	
1	С	0.32	0/2127	0.53	0/2899	
1	D	0.36	0/2166	0.54	0/2949	
All	All	0.34	0/8618	0.53	0/11734	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

# 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2146	0	2082	14	0
1	В	2094	0	2003	13	0
1	С	2084	0	1976	7	0
1	D	2119	0	2031	17	0
2	A	12	0	8	1	0
2	В	12	0	8	1	0
2	С	12	0	8	0	0
2	D	12	0	8	1	0
3	A	4	0	6	1	0



$\alpha \cdots$	· ·	•	
Continued	trom	nromanne	naae
-	110111	picolous	payc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	8	0	12	3	0
3	С	4	0	6	1	0
3	D	8	0	12	0	0
4	A	52	0	0	1	0
4	В	50	0	0	0	0
4	С	29	0	0	0	0
4	D	37	0	0	0	0
All	All	8683	0	8160	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll}  ext{Interatomic} \  ext{distance} \ ( ext{\AA}) \end{array}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
1:B:106:ILE:HD12	1:B:125:ILE:HD11	1.72	0.70
1:B:96:ARG:NH2	1:B:162:GLU:OE1	2.21	0.69
1:B:188:PRO:HD3	1:B:194:ILE:HG13	1.75	0.67
1:D:188:PRO:HD3	1:D:194:ILE:HG13	1.77	0.66
1:B:66:HIS:ND1	1:B:241:TYR:HB2	2.16	0.60

There are no symmetry-related clashes.

# 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	279/308 (91%)	275 (99%)	4 (1%)	0	100	100
1	В	278/308 (90%)	275 (99%)	3 (1%)	0	100	100
1	С	277/308 (90%)	274 (99%)	3 (1%)	0	100	100
1	D	281/308 (91%)	275 (98%)	6 (2%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
All	All	1115/1232 (90%)	1099 (99%)	16 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	219/249 (88%)	217 (99%)	2 (1%)	78	90	
1	В	$208/249 \ (84\%)$	208 (100%)	0	100	100	
1	С	$205/249 \ (82\%)$	203 (99%)	2 (1%)	76	88	
1	D	211/249 (85%)	209 (99%)	2 (1%)	78	90	
All	All	843/996 (85%)	837 (99%)	6 (1%)	84	92	

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	143	SER
1	D	265	LYS
1	С	168	SER
1	A	268	ARG
1	D	181	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	HIS

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trens	Chain	D. a. T.:1.		В	ond leng	$\operatorname{gths}$	Е	ond ang	gles
MIOI	Type	Chain	m Res	Link	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	D	303	-	3,3,3	0.46	0	2,2,2	0.43	0
3	EDO	В	303	_	3,3,3	0.50	0	2,2,2	0.28	0
3	EDO	D	302	_	3,3,3	0.60	0	2,2,2	0.23	0
3	EDO	С	302	-	3,3,3	0.58	0	2,2,2	0.08	0
3	EDO	A	302	_	3,3,3	0.56	0	2,2,2	0.14	0
3	EDO	В	302	_	3,3,3	0.59	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	303	_	-	0/1/1/1	-
3	EDO	В	303	-	-	0/1/1/1	=
3	EDO	D	302	-	-	0/1/1/1	-
3	EDO	С	302	_	-	1/1/1/1	-
3	EDO	A	302	-	-	1/1/1/1	=
3	EDO	В	302	-	-	1/1/1/1	=

There are no bond length outliers.

There are no bond angle outliers.



There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	EDO	O1-C1-C2-O2
3	В	302	EDO	O1-C1-C2-O2
3	С	302	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	303	EDO	1	0
3	С	302	EDO	1	0
3	A	302	EDO	1	0
3	В	302	EDO	2	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	281/308 (91%)	-0.14	5 (1%) 68	66	41, 66, 94, 135	0
1	В	280/308 (90%)	-0.05	6 (2%) 63	61	43, 67, 103, 133	0
1	С	279/308 (90%)	-0.03	7 (2%) 57	55	41, 75, 109, 152	0
1	D	282/308 (91%)	-0.08	4 (1%) 75	73	37, 68, 102, 134	0
All	All	1122/1232 (91%)	-0.08	22 (1%) 65	63	37, 68, 104, 152	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	106	ILE	6.4
1	A	233	GLY	6.0
1	С	112	CYS	5.9
1	В	233	GLY	3.9
1	В	66	HIS	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	EDO	В	303	4/4	0.65	0.39	93,96,97,99	0
3	EDO	В	302	4/4	0.69	0.33	66,68,69,73	0
3	EDO	A	302	4/4	0.80	0.22	71,72,72,75	0
3	EDO	D	303	4/4	0.80	0.22	69,81,91,100	0
3	EDO	D	302	4/4	0.89	0.20	48,64,64,66	0
3	EDO	С	302	4/4	0.89	0.14	49,58,65,65	0
2	PHE	В	301	12/12	0.92	0.20	43,52,59,60	0
2	PHE	С	301	12/12	0.94	0.18	42,52,59,60	0
2	PHE	A	301	12/12	0.95	0.17	39,48,61,63	0
2	PHE	D	301	12/12	0.95	0.17	42,53,56,61	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

