



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 5, 2024 – 06:14 AM EST

PDB ID : 1VLZ
Title : UNCOUPLED PHOSPHORYLATION AND ACTIVATION IN BACTERIAL CHEMOTAXIS: THE 2.1 ANGSTROM STRUCTURE OF A THREONINE TO ISOLEUCINE MUTANT AT POSITION 87 OF CHEY
Authors : Ganguli, S.; Wang, H.; Matsumura, P.; Volz, K.
Deposited on : 1995-04-21
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	128	 75% 19% 6%
1	B	128	 70% 21% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHEY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	128	954	609	156	183	6	0	0	0
1	B	127	944	604	158	176	6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	ILE	THR	conflict	UNP P06143
B	87	ILE	THR	conflict	UNP P06143

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	92	Total	O	0	0
			92	92		
2	B	80	Total	O	0	0
			80	80		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CHEY

Chain A:  75% 19% 6%



- Molecule 1: CHEY

Chain B:  70% 21% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.04Å 72.47Å 36.15Å 90.00° 109.06° 90.00°	Depositor
Resolution (Å)	10.00 – 2.05 30.91 – 2.05	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.05) 87.8 (30.91-2.05)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.37 (at 2.05Å)	Xtrriage
Refinement program	PROFFT	Depositor
R, R_{free}	0.156 , (Not available) 0.162 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtrriage
Anisotropy	0.073	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 95.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.048 for -h-1,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2070	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/966	2.13	28/1304 (2.1%)
1	B	0.89	0/956	2.04	29/1290 (2.2%)
All	All	0.90	0/1922	2.09	57/2594 (2.2%)

There are no bond length outliers.

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ARG	NE-CZ-NH1	26.54	133.57	120.30
1	A	18	ARG	NE-CZ-NH2	-16.39	112.11	120.30
1	B	75	ASP	CB-CG-OD1	15.12	131.90	118.30
1	B	22	ARG	NE-CZ-NH1	-14.89	112.85	120.30
1	B	73	ARG	NE-CZ-NH2	-14.41	113.10	120.30
1	A	41	ASP	CB-CG-OD2	14.33	131.20	118.30
1	B	19	ARG	NE-CZ-NH1	13.27	126.93	120.30
1	A	12	ASP	CB-CG-OD2	12.98	129.99	118.30
1	B	19	ARG	NE-CZ-NH2	-12.00	114.30	120.30
1	A	64	ASP	CB-CG-OD2	-10.17	109.15	118.30
1	B	22	ARG	NE-CZ-NH2	9.81	125.21	120.30
1	A	64	ASP	CB-CG-OD1	9.67	127.00	118.30
1	B	51	TYR	CB-CG-CD2	-9.13	115.52	121.00
1	B	19	ARG	CD-NE-CZ	8.81	135.94	123.60
1	A	22	ARG	NE-CZ-NH1	8.35	124.47	120.30
1	B	126	LYS	CA-CB-CG	7.88	130.74	113.40
1	B	32	ASN	N-CA-CB	-7.79	96.58	110.60
1	A	78	MET	CA-CB-CG	7.52	126.08	113.30
1	A	75	ASP	CB-CG-OD1	7.44	125.00	118.30
1	A	118	GLU	CG-CD-OE2	-7.34	103.62	118.30
1	B	56	SER	N-CA-CB	-7.22	99.66	110.50
1	B	64	ASP	CB-CG-OD1	7.07	124.66	118.30
1	A	127	LEU	CA-CB-CG	6.85	131.06	115.30
1	B	51	TYR	CB-CG-CD1	6.84	125.11	121.00
1	B	100	GLN	CA-CB-CG	6.72	128.18	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	32	ASN	CB-CA-C	6.57	123.54	110.40
1	A	34	GLU	OE1-CD-OE2	6.55	131.16	123.30
1	A	12	ASP	CB-CG-OD1	-6.53	112.42	118.30
1	B	81	LEU	N-CA-CB	-6.53	97.35	110.40
1	B	117	GLU	CG-CD-OE2	-6.08	106.13	118.30
1	A	41	ASP	CB-CG-OD1	-6.03	112.87	118.30
1	B	45	LYS	CA-CB-CG	-6.02	100.16	113.40
1	A	93	GLU	OE1-CD-OE2	-5.81	116.33	123.30
1	B	49	GLY	N-CA-C	5.74	127.44	113.10
1	A	38	ASP	CB-CG-OD2	5.73	123.46	118.30
1	A	27	GLU	CG-CD-OE2	-5.72	106.86	118.30
1	A	118	GLU	CG-CD-OE1	5.71	129.72	118.30
1	A	83	VAL	CA-CB-CG2	5.69	119.43	110.90
1	B	41	ASP	CB-CA-C	5.55	121.50	110.40
1	A	91	LYS	CB-CA-C	-5.51	99.37	110.40
1	B	37	GLU	OE1-CD-OE2	5.48	129.88	123.30
1	A	106	TYR	CA-CB-CG	-5.42	103.10	113.40
1	A	102	GLY	CA-C-O	-5.42	110.85	120.60
1	A	66	LEU	CB-CA-C	5.41	120.47	110.20
1	B	75	ASP	CB-CG-OD2	-5.33	113.50	118.30
1	A	22	ARG	CD-NE-CZ	-5.27	116.22	123.60
1	B	5	GLU	CG-CD-OE1	-5.25	107.81	118.30
1	A	106	TYR	CB-CG-CD1	-5.23	117.86	121.00
1	A	27	GLU	OE1-CD-OE2	5.23	129.58	123.30
1	B	34	GLU	N-CA-CB	5.22	120.00	110.60
1	A	119	LYS	N-CA-CB	5.22	120.00	110.60
1	A	34	GLU	CG-CD-OE2	-5.16	107.99	118.30
1	B	73	ARG	NH1-CZ-NH2	5.12	125.03	119.40
1	B	18	ARG	NE-CZ-NH2	-5.07	117.77	120.30
1	B	26	LYS	N-CA-CB	5.07	119.72	110.60
1	B	57	ASP	CB-CG-OD1	5.01	122.81	118.30
1	B	117	GLU	OE1-CD-OE2	5.01	129.31	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	954	0	951	17	0
1	B	944	0	947	13	0
2	A	92	0	0	3	0
2	B	80	0	0	0	0
All	All	2070	0	1898	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:LEU:HD23	1:B:129:MET:SD	2.12	0.89
1:B:75:ASP:O	1:B:79:SER:HB3	1.95	0.66
1:A:58:TRP:HE3	1:A:85:MET:HE2	1.62	0.65
1:A:34:GLU:OE1	2:A:151:HOH:O	2.14	0.65
1:A:58:TRP:CE3	1:A:85:MET:HE2	2.35	0.62
1:A:43:LEU:O	1:A:47:GLN:HG2	2.04	0.56
1:B:43:LEU:HD11	1:B:71:THR:HG21	1.88	0.56
1:B:118:GLU:O	1:B:122:LYS:HG3	2.06	0.56
1:A:94:ASN:ND2	2:A:196:HOH:O	2.16	0.55
1:B:34:GLU:HG3	1:B:51:TYR:OH	2.08	0.53
1:B:127:LEU:HB3	1:B:129:MET:HG3	1.91	0.53
1:B:3:ASP:C	1:B:5:GLU:H	2.15	0.49
1:B:15:SER:HB2	1:B:19:ARG:HH11	1.76	0.49
1:A:38:ASP:HB2	1:A:61:PRO:O	2.15	0.46
1:A:118:GLU:HG2	2:A:212:HOH:O	2.16	0.45
1:B:124:PHE:HD1	1:B:129:MET:SD	2.40	0.45
1:A:26:LYS:HB2	1:A:26:LYS:HE3	0.97	0.43
1:A:34:GLU:HG3	1:A:51:TYR:OH	2.19	0.43
1:B:81:LEU:HA	1:B:82:PRO:HD3	1.88	0.43
1:A:24:LEU:O	1:A:28:LEU:HG	2.19	0.43
1:A:106:TYR:CD1	1:A:106:TYR:C	2.92	0.43
1:B:41:ASP:O	1:B:45:LYS:HG3	2.18	0.42
1:A:85:MET:O	1:A:106:TYR:HA	2.18	0.42
1:A:58:TRP:CE3	1:A:85:MET:CE	3.02	0.42
1:A:81:LEU:HA	1:A:82:PRO:HD3	1.95	0.42
1:A:106:TYR:O	1:A:119:LYS:HE2	2.19	0.42
1:B:9:LEU:HD11	1:B:36:ALA:HB2	2.03	0.41
1:B:7:LYS:HE2	1:B:32:ASN:OD1	2.20	0.41
1:A:25:LEU:HA	1:A:25:LEU:HD23	1.81	0.40
1:A:47:GLN:HE21	1:A:47:GLN:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/128 (98%)	124 (98%)	2 (2%)	0	100	100
1	B	125/128 (98%)	121 (97%)	3 (2%)	1 (1%)	19	10
All	All	251/256 (98%)	245 (98%)	5 (2%)	1 (0%)	34	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	90	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/102 (94%)	93 (97%)	3 (3%)	40	33
1	B	94/102 (92%)	92 (98%)	2 (2%)	53	48
All	All	190/204 (93%)	185 (97%)	5 (3%)	46	39

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	GLN
1	A	61	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	127	LEU
1	B	53	PHE
1	B	79	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/128 (100%)	-0.61	0 100 100	7, 12, 25, 34	0
1	B	127/128 (99%)	-0.57	1 (0%) 86 88	7, 17, 31, 50	0
All	All	255/256 (99%)	-0.59	1 (0%) 92 93	7, 14, 30, 50	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	129	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.