

wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2023 – 03:37 AM EDT

PDB ID : 6VLC

Title : Crystal structure of UDP-GlcNAc 2-epimerase from Neisseria meningitidis

bound to UDP-GlcNAc

Authors: Fisher, A.J.; Hurlburt, N.K.

Deposited on : 2020-01-23

Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 12438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called UDP-N-acetylglucosamine 2-epimerase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	371	Total	С	N	О	S	0	0	0
1	A	3/1	2926	1874	485	557	10	0	U	
1	В	366	Total	С	N	О	S	0	0	0
1	Ъ	300	2877	1847	471	549	10	0	U	
1	С	371	Total	С	N	О	S	0	0	0
1		3/1	2926	1874	485	557	10	U	U	
1	D	365	Total	С	N	О	S	0	0	0
1	D	309	2873	1845	470	548	10	U	U	U

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	373	LEU	-	expression tag	UNP A0A0U1RGY0
A	374	GLU	-	expression tag	UNP A0A0U1RGY0
A	375	HIS	-	expression tag	UNP A0A0U1RGY0
A	376	HIS	-	expression tag	UNP A0A0U1RGY0
A	377	HIS	-	expression tag	UNP A0A0U1RGY0
A	378	HIS	-	expression tag	UNP A0A0U1RGY0
A	379	HIS	-	expression tag	UNP A0A0U1RGY0
A	380	HIS	-	expression tag	UNP A0A0U1RGY0
В	373	LEU	-	expression tag	UNP A0A0U1RGY0
В	374	GLU	-	expression tag	UNP A0A0U1RGY0
В	375	HIS	-	expression tag	UNP A0A0U1RGY0
В	376	HIS	-	expression tag	UNP A0A0U1RGY0
В	377	HIS	-	expression tag	UNP A0A0U1RGY0
В	378	HIS	-	expression tag	UNP A0A0U1RGY0
В	379	HIS	-	expression tag	UNP A0A0U1RGY0
В	380	HIS	-	expression tag	UNP A0A0U1RGY0
С	373	LEU	-	expression tag	UNP A0A0U1RGY0
С	374	GLU	-	expression tag	UNP A0A0U1RGY0
С	375	HIS	-	expression tag	UNP A0A0U1RGY0
С	376	HIS		expression tag	UNP A0A0U1RGY0
С	377	HIS	-	expression tag	UNP A0A0U1RGY0


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	378	HIS	-	expression tag	UNP A0A0U1RGY0
С	379	HIS	-	expression tag	UNP A0A0U1RGY0
С	380	HIS	-	expression tag	UNP A0A0U1RGY0
D	373	LEU	-	expression tag	UNP A0A0U1RGY0
D	374	GLU	-	expression tag	UNP A0A0U1RGY0
D	375	HIS	ı	expression tag	UNP A0A0U1RGY0
D	376	HIS	-	expression tag	UNP A0A0U1RGY0
D	377	HIS	-	expression tag	UNP A0A0U1RGY0
D	378	HIS	-	expression tag	UNP A0A0U1RGY0
D	379	HIS	-	expression tag	UNP A0A0U1RGY0
D	380	HIS	-	expression tag	UNP A0A0U1RGY0

• Molecule 2 is URIDINE-DIPHOSPHATE-N-ACETYLGLUCOSAMINE (three-letter code: UD1) (formula: $C_{17}H_{27}N_3O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	Λ	1	Total	С	N	О	Р	0	0
	A	1	39	17	3	17	2	U	U
9	C	1	Total	С	N	О	Р	0	0
		1	39	17	3	17	2	U	0

• Molecule 3 is water.

\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	179	Total O 179 179	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	207	Total O 207 207	0	0
3	С	169	Total O 169 169	0	0
3	D	203	Total O 203 203	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	124.88Å 129.74Å 213.39Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.97 - 2.15	Depositor
% Data completeness	98.9 (38.97-2.15)	Depositor
(in resolution range)	,	-
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.29 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.168 , 0.212	Depositor
Wilson B-factor (\mathring{A}^2)	33.8	Xtriage
Anisotropy	0.034	Xtriage
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.036 for -k,-h,-l	Xtriage
Total number of atoms	12438	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Res	Link	Во	ond leng	$ ag{ths}$	Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UD1	A	400	-	38,41,41	0.45	0	57,62,62	2.12	9 (15%)
2	UD1	С	400	-	38,41,41	0.47	0	57,62,62	1.95	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UD1	A	400	-	-	6/26/63/63	0/3/3/3
2	UD1	С	400	-	-	6/26/63/63	0/3/3/3

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	A	400	UD1	C1'-C2'-N2'	7.89	124.60	111.00
2	A	400	UD1	O5'-C1'-O1'	-6.94	102.29	111.36
2	С	400	UD1	O5'-C1'-O1'	-6.94	102.30	111.36
2	С	400	UD1	C1'-C2'-N2'	6.25	121.77	111.00
2	A	400	UD1	PB-O1'-C1'	5.25	140.03	119.74

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400	UD1	O5'-C1'-O1'-PB
2	A	400	UD1	C3'-C2'-N2'-C7'
2	A	400	UD1	C8'-C7'-N2'-C2'
2	A	400	UD1	O7'-C7'-N2'-C2'
2	С	400	UD1	O5'-C1'-O1'-PB


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

