

# Full wwPDB EM Validation Report (i)

Jan 6, 2025 – 10:56 AM EST

PDB ID : 8VLR

EMDB ID : EMD-43342

Title : Cryo-EM structure of native H2AK119bu nucleosome at 2.6

Authors: Wang, Y.; Zhang, K.

Deposited on : 2024-01-12

Resolution : 2.60 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113

MolProbity : 4.02b-467

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : 1.9.13

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	98	88%	11% •
1	Е	98	94%	6%
2	В	84	86%	14%
2	F	84	95%	5%
3	С	108	84%	15% •
3	G	108	91%	9%
4	D	93	91%	9%
4	Н	93	92%	8%

Continued on next page...



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Length	Quality of chain						
5	K	136	54%	46%					
6	L	136	74%	26%					



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 21181 atoms, of which 9524 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Histone H3.1.

	$\mathbf{Mol}$	Chain	Residues		${f Atoms}$					AltConf	Trace	
	1	Λ	98	Total	С	Н	N	О	S	0	0	
	1	Α	90	1652	508	846	156	138	4	0		
İ	1	E	98	Total	С	Н	N	О	S	0	0	
	1	E	90	1652	508	846	156	138	4	0	U	

• Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	${f Atoms}$					AltConf	Trace	
2	D	84	Total	С	Н	N	О	S	0	0
2	Ъ	04	1398	424	725	133	115	1	U	U
2	Г	84	Total	С	Н	N	О	S	0	0
<u> </u>	Г	04	1398	424	725	133	115	1	U	U

• Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	С	108	Total	С	Н	N	О	0	0
3		100	1735	526	900	165	144	0	
9	C	108	Total	С	Н	N	О	0	0
3	G	108	1735	526	900	165	144	0	U

• Molecule 4 is a protein called Histone H2B type 1-A.

Mol	Chain	Residues		Atoms						Trace
4	D	93	Total	С	Н	N	О	S	0	0
4	ע	90	1482	458	754	128	140	2	U	. 0
4	П	93	Total	С	Н	N	О	S	0	0
4	Н	90	1482	458	754	128	140	2	0	U

• Molecule 5 is a DNA chain called DNA (136-MER).



Mol	Chain	Residues			Aton	ns			AltConf	Trace
5	I/	136	Total	С	Н	N	О	Р	0	0
5	IX.	190	4320	1330	1539	494	821	136	U	U

 $\bullet$  Molecule 6 is a DNA chain called DNA (136-MER).

Mol	Chain	Residues			Aton	ns			AltConf	Trace
6	Т	136	Total	С	Н	N	О	Р	0	0
0	L	130	4327	1333	1535	515	809	135	U	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





 $\bullet$  Molecule 3: Histone H2A type 1-B/E

Chain G:

91%

Molecule 4: Histone H2B type 1-A

Chain D: 91% 9%



• Molecule 4: Histone H2B type 1-A

Chain H: 92% 8%



• Molecule 5: DNA (136-MER)

Chain K: 54% 46%





• Molecule 6: DNA (136-MER)

Chain L: 74% 26%





# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1288371	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.845	Depositor
Minimum map value	-3.876	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.110	Depositor
Recommended contour level	0.321	Depositor
Map size (Å)	341.75998, 341.75998, 341.75998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.068, 1.068, 1.068	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Во	ond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.39	0/818	0.60	0/1097
1	Е	0.35	0/818	0.58	0/1097
2	В	0.38	0/680	0.61	0/908
2	F	0.36	0/680	0.60	0/908
3	С	0.33	0/845	0.57	0/1139
3	G	0.34	0/845	0.56	0/1139
4	D	0.34	0/739	0.51	0/994
4	Н	0.34	0/739	0.50	0/994
5	K	0.75	0/3116	0.99	1/4806 (0.0%)
6	L	0.77	0/3135	0.94	0/4838
All	All	0.59	0/12415	0.81	1/17920 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
5	K	80	DT	O4'-C1'-N1	5.12	111.59	108.00

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	806	846	844	9	0
1	Е	806	846	844	4	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	673	725	722	7	0
2	F	673	725	722	2	0
3	С	835	900	897	12	0
3	G	835	900	897	7	0
4	D	728	754	751	6	0
4	Н	728	754	751	5	0
5	K	2781	1539	1539	52	0
6	L	2792	1535	1535	27	0
All	All	11657	9524	9502	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
3:C:11:ARG:NH1	3:C:12:ALA:O	2.19	0.74
5:K:89:DC:H2'	5:K:90:DT:H72	1.70	0.71
3:C:36:LYS:HE3	3:C:36:LYS:HA	1.79	0.65
5:K:13:DC:H2'	5:K:14:DT:H72	1.82	0.61
5:K:23:DT:H2'	5:K:24:DT:H72	1.83	0.60
3:G:40:SER:OG	3:G:42:ARG:O	2.23	0.57
3:C:32:ARG:NH1	5:K:29:DA:OP1	2.38	0.56
4:D:42:ILE:HD13	4:D:63:MET:HB3	1.88	0.56
5:K:37:DT:H2'	5:K:38:DT:H71	1.89	0.55
5:K:69:DC:H2'	5:K:70:DT:H72	1.89	0.55
3:G:69:ALA:O	3:G:73:ASN:ND2	2.39	0.54
5:K:13:DC:C2'	5:K:14:DT:H72	2.38	0.54
5:K:48:DT:H2"	5:K:49:DC:C6	2.44	0.53
5:K:127:DA:H2"	5:K:128:DT:H71	1.92	0.52
1:A:75:ALA:O	2:B:20:LYS:NZ	2.43	0.52
6:L:204:DG:H2"	6:L:205:DG:C8	2.47	0.50
6:L:166:DT:H2"	6:L:167:DT:C6	2.47	0.50
6:L:219:DA:H2"	6:L:220:DT:O5'	2.12	0.50
5:K:43:DA:H2"	5:K:44:DC:O5'	2.12	0.49
5:K:54:DA:H4'	5:K:55:DA:OP1	2.13	0.49
3:C:77:ARG:HH11	3:C:77:ARG:HG3	1.78	0.49
5:K:69:DC:C2'	5:K:70:DT:H72	2.44	0.48
5:K:44:DC:H2'	5:K:45:DT:H72	1.96	0.48
5:K:46:DG:N2	6:L:248:DA:C2	2.82	0.48
5:K:141:DT:H2"	5:K:142:DA:C8	2.49	0.47



 $Continued\ from\ previous\ page...$ 

Continuea from pret		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\rm \mathring{A})$	overlap (Å)
3:C:51:LEU:HD13	4:D:74:ILE:HG21	1.95	0.47
5:K:90:DT:H4'	5:K:91:DT:OP1	2.15	0.47
6:L:180:DT:H2'	6:L:181:DA:C8	2.50	0.47
6:L:209:DG:H4'	6:L:210:DT:OP1	2.15	0.47
5:K:45:DT:H2"	5:K:46:DG:C8	2.50	0.47
3:C:54:VAL:HG21	4:D:99:VAL:HG21	1.98	0.46
5:K:22:DC:H2'	5:K:23:DT:H72	1.98	0.46
1:A:73:GLU:HG3	2:B:22:LEU:HA	1.98	0.46
1:A:79:LYS:CG	1:A:80:THR:HG23	2.46	0.46
5:K:86:DT:H2"	5:K:87:DG:C8	2.51	0.46
5:K:128:DT:H2"	5:K:129:DC:C6	2.51	0.46
6:L:163:DA:H2"	6:L:164:DG:C8	2.51	0.46
2:B:19:ARG:NH2	5:K:51:DA:OP2	2.42	0.45
6:L:165:DA:H2"	6:L:166:DT:C6	2.51	0.45
5:K:105:DT:H2"	5:K:106:DA:C8	2.52	0.45
5:K:37:DT:H2"	5:K:38:DT:C6	2.51	0.45
1:E:103:LEU:CD2	2:F:57:VAL:HG11	2.46	0.45
5:K:58:DG:H4'	5:K:59:DG:OP1	2.16	0.45
3:C:102:ILE:HG23	4:D:62:ILE:HG12	1.99	0.45
5:K:109:DA:H4'	5:K:110:DA:OP1	2.17	0.45
3:C:26:PRO:O	3:C:30:VAL:HG23	2.17	0.44
5:K:27:DA:C2	6:L:267:DG:N2	2.85	0.44
1:A:74:ILE:HD11	2:B:62:LEU:HB3	2.00	0.44
6:L:242:DT:H2"	6:L:243:DG:O4'	2.17	0.44
1:A:126:LEU:HD22	1:E:113:HIS:CG	2.52	0.44
3:G:32:ARG:NH2	6:L:176:DA:OP1	2.49	0.44
5:K:14:DT:H2"	5:K:15:DG:C8	2.53	0.44
6:L:263:DT:H2'	6:L:264:DT:H72	1.99	0.44
4:D:106:GLU:OE2	4:D:109:LYS:NZ	2.49	0.44
3:G:71:ARG:HH11	3:G:71:ARG:HG2	1.82	0.44
5:K:111:DA:C8	5:K:112:DT:H72	2.53	0.44
6:L:258:DT:H2"	6:L:259:DA:C8	2.53	0.43
2:F:35:ARG:O	2:F:39:ARG:HG2	2.19	0.43
4:H:54:GLY:HA3	6:L:167:DT:OP1	2.18	0.43
5:K:48:DT:C2'	5:K:49:DC:C6	3.02	0.43
3:G:54:VAL:HG21	4:H:99:VAL:HG21	2.01	0.43
5:K:54:DA:H2"	5:K:55:DA:O5'	2.19	0.43
5:K:106:DA:C2	6:L:187:DT:C7	3.01	0.43
5:K:43:DA:H4'	5:K:44:DC:OP1	2.18	0.42
5:K:89:DC:H4'	5:K:90:DT:OP1	2.19	0.42
5:K:119:DT:H2"	5:K:120:DT:C5	2.53	0.42



 $Continued\ from\ previous\ page...$ 

Continuea from pret		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap(Å)
1:A:69:ARG:NH2	6:L:237:DT:OP1	2.46	0.42
5:K:103:DG:C8	5:K:104:DT:H72	2.55	0.42
3:G:73:ASN:H	3:G:73:ASN:HD22	1.67	0.42
3:C:77:ARG:HD3	5:K:19:DG:H5"	2.01	0.42
1:A:108:ASN:O	1:A:112:ILE:HG12	2.19	0.42
2:B:35:ARG:O	2:B:39:ARG:HG2	2.19	0.42
3:C:13:LYS:O	3:C:14:ALA:HB3	2.19	0.42
5:K:45:DT:C2'	5:K:46:DG:C8	3.03	0.42
5:K:81:DG:H2"	5:K:82:DA:C8	2.55	0.42
5:K:131:DG:N2	6:L:163:DA:C2	2.88	0.42
5:K:120:DT:H2"	5:K:121:DG:C8	2.54	0.42
6:L:266:DT:H2"	6:L:267:DG:C8	2.55	0.42
4:H:42:ILE:HD11	4:H:67:VAL:HG21	2.01	0.42
6:L:219:DA:H4'	6:L:220:DT:OP1	2.20	0.42
1:A:78:PHE:HB2	2:B:70:VAL:HG11	2.00	0.42
4:H:38:TYR:O	4:H:42:ILE:HG12	2.19	0.42
6:L:226:DT:H2"	6:L:227:DG:C8	2.55	0.42
5:K:33:DG:H1'	5:K:34:DT:H5"	2.00	0.42
5:K:90:DT:H2"	5:K:91:DT:O5'	2.20	0.42
6:L:227:DG:H2"	6:L:228:DA:C8	2.55	0.42
6:L:216:DT:H2"	6:L:217:DG:C8	2.55	0.41
3:C:76:THR:O	3:C:76:THR:CG2	2.67	0.41
5:K:25:DC:H2"	5:K:26:DC:C5	2.55	0.41
5:K:38:DT:H2"	5:K:39:DA:C8	2.55	0.41
6:L:251:DT:H2'	6:L:252:DT:H71	2.02	0.41
3:C:62:ILE:HD12	4:D:66:PHE:CE2	2.55	0.41
1:E:106:ASP:OD2	1:E:131:ARG:NE	2.44	0.41
6:L:251:DT:C2'	6:L:252:DT:H71	2.51	0.41
3:G:26:PRO:O	3:G:30:VAL:HG23	2.21	0.41
5:K:37:DT:C2'	5:K:38:DT:H71	2.50	0.41
6:L:177:DG:C8	6:L:178:DT:H72	2.56	0.41
1:A:103:LEU:CD2	2:B:57:VAL:HG11	2.51	0.41
5:K:70:DT:H2"	5:K:71:DG:C8	2.56	0.41
5:K:96:DT:H4'	5:K:97:DG:OP1	2.21	0.41
6:L:193:DC:H2"	6:L:194:DT:C6	2.56	0.41
1:E:45:THR:O	1:E:49:ARG:HG3	2.21	0.40
4:H:53:THR:HG22	4:H:54:GLY:O	2.20	0.40
5:K:16:DC:H2"	5:K:17:DC:C6	2.56	0.40
5:K:86:DT:C2'	5:K:87:DG:C8	3.05	0.40
5:K:89:DC:C2'	5:K:90:DT:H72	2.45	0.40
5:K:129:DC:H2"	5:K:130:DT:C7	2.50	0.40



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
5:K:138:DC:C2'	5:K:139:DA:C8	3.04	0.40
6:L:166:DT:H2'	6:L:167:DT:H72	2.03	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	96/98~(98%)	91 (95%)	5 (5%)	0	100	100
1	$\mathbf{E}$	96/98 (98%)	96 (100%)	0	0	100	100
2	В	82/84 (98%)	81 (99%)	1 (1%)	0	100	100
2	F	82/84 (98%)	81 (99%)	1 (1%)	0	100	100
3	C	106/108 (98%)	104 (98%)	2 (2%)	0	100	100
3	G	106/108 (98%)	104 (98%)	2 (2%)	0	100	100
4	D	91/93 (98%)	88 (97%)	3 (3%)	0	100	100
4	Н	91/93 (98%)	90 (99%)	1 (1%)	0	100	100
All	All	750/766 (98%)	735 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	85/85 (100%)	83 (98%)	2 (2%)	44	70
1	$\mathbf{E}$	85/85 (100%)	85 (100%)	0	100	100
2	В	69/69 (100%)	65 (94%)	4 (6%)	17	36
2	F	69/69 (100%)	68 (99%)	1 (1%)	62	82
3	$\mathbf{C}$	85/85 (100%)	81 (95%)	4 (5%)	22	45
3	G	85/85 (100%)	84 (99%)	1 (1%)	67	85
4	D	82/82 (100%)	82 (100%)	0	100	100
4	Н	82/82 (100%)	81 (99%)	1 (1%)	67	85
All	All	642/642 (100%)	629 (98%)	13 (2%)	50	74

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ASP
1	A	78	PHE
2	В	24	ASP
2	В	25	ASN
2	В	79	LYS
2	В	93	GLN
3	С	11	ARG
3	С	74	LYS
3	С	91	GLU
3	С	101	THR
2	F	23	ARG
3	G	76	THR
4	Н	57	SER

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
3	G	110	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry (i)

There are no ligands in this entry.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-43342. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

#### 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



#### 6.1.2 Raw map





The images above show the map projected in three orthogonal directions.




### 6.2 Central slices (i)

#### 6.2.1 Primary map








X Index: 160


Y Index: 160

Z Index: 160

#### 6.2.2 Raw map







X Index: 160

Y Index: 160


Z Index: 160


The images above show central slices of the map in three orthogonal directions.




### 6.3 Largest variance slices (i)

#### 6.3.1 Primary map








X Index: 152


Y Index: 156

Z Index: 144

#### 6.3.2 Raw map







X Index: 174

Y Index: 148


Z Index: 144

The images above show the largest variance slices of the map in three orthogonal directions.




### 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



#### 6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



### 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.321. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

### 6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $155~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $140~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 7.3 Rotationally averaged power spectrum (i)




\*Reported resolution corresponds to spatial frequency of 0.385  $\rm \mathring{A}^{-1}$ 



### 8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

#### 8.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.385  $\rm \mathring{A}^{-1}$ 



### 8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	2.60	-	-	
Author-provided FSC curve	2.60	2.90	2.67	
Unmasked-calculated*	3.24	3.75	3.28	


<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.24 differs from the reported value 2.6 by more than 10 %



## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-43342 and PDB model 8VLR. Per-residue inclusion information can be found in section 3 on page 6.


### 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.321 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.321).



### 9.4 Atom inclusion (i)



At the recommended contour level, 92% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.



### 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.321) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9250	0.6020
A	0.9650	0.6410
В	0.9540	0.6390
С	0.9580	0.6380
D	0.9690	0.6390
Е	0.9790	0.6510
F	0.9470	0.6360
G	0.9740	0.6510
Н	0.9780	0.6460
K	0.8920	0.5560
L	0.8870	0.5590



