

Full wwPDB X-ray Structure Validation Report (i)

Nov 5, 2023 – 11:09 AM EST

:	6VNJ
:	JAK2 JH1 in complex with PN4-014
:	Davis, R.R.; Schonbrunn, E.
	2020-01-29
:	1.90 Å(reported)
	:

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	:::::::::::::::::::::::::::::::::::::::	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	308	<mark>6%</mark> 84%	11%	•••
1	В	308	87%	7%	6%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	В	290		C 1525			Р 1	S 14	0	1	0
1	А	299	Total 2473	C 1571	N 429	0 458	Р 1	S 14	0	1	0

• Molecule 1 is a protein called Tyrosine-protein kinase JAK2.


Chain	Residue	Modelled	Actual	Comment	Reference
В	825	HIS	-	expression tag	UNP O60674
В	826	HIS	-	expression tag	UNP O60674
В	827	HIS	-	expression tag	UNP O60674
В	828	HIS	-	expression tag	UNP O60674
В	829	HIS	-	expression tag	UNP O60674
В	830	HIS	-	expression tag	UNP O60674
В	831	HIS	-	expression tag	UNP O60674
В	832	HIS	-	expression tag	UNP O60674
В	833	GLU	-	expression tag	UNP O60674
В	834	ASN	-	expression tag	UNP O60674
В	835	LEU	-	expression tag	UNP O60674
В	836	TYR	-	expression tag	UNP O60674
В	837	PHE	-	expression tag	UNP O60674
В	838	GLN	-	expression tag	UNP O60674
В	839	GLY	-	expression tag	UNP O60674
А	825	HIS	-	expression tag	UNP O60674
А	826	HIS	-	expression tag	UNP O60674
А	827	HIS	-	expression tag	UNP O60674
А	828	HIS	-	expression tag	UNP O60674
А	829	HIS	-	expression tag	UNP O60674
А	830	HIS	-	expression tag	UNP O60674
А	831	HIS	-	expression tag	UNP O60674
А	832	HIS	-	expression tag	UNP O60674
А	833	GLU	-	expression tag	UNP O60674
А	834	ASN	_	expression tag	UNP O60674

There are 30 discrepancies between the modelled and reference sequences:



Contentia	Continuacia from precious page						
Chain	Residue	Modelled	Actual	Comment	Reference		
А	835	LEU	-	expression tag	UNP O60674		
А	836	TYR	-	expression tag	UNP O60674		
А	837	PHE	-	expression tag	UNP O60674		
А	838	GLN	-	expression tag	UNP O60674		
А	839	GLY	-	expression tag	UNP O60674		

• Molecule 2 is 3-[4-(2-{[4-(piperidin-4-yl)phenyl]amino}-6,7-dihydro-5H-pyrrolo[2,3-d]pyr imidin-4-yl)-1H-pyrazol-1-yl]propanenitrile (three-letter code: R5S) (formula: C₂₃H₂₆N₈) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total C N 31 23 8	0	0
2	А	1	Total C N 31 23 8	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	122	Total O 122 122	0	0
3	А	142	Total O 142 142	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41	Depositor
Cell constants	112.13Å 112.13Å 70.58Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	-
Resolution (Å)	43.90 - 1.90	Depositor
	43.90 - 1.90	EDS
% Data completeness	$96.6\ (43.90\text{-}1.90)$	Depositor
(in resolution range)	99.6 (43.90 - 1.90)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.76 (at 1.89 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.14-3260_3260	Depositor
D D	0.185 , 0.211	Depositor
R, R_{free}	0.183 , 0.206	DCC
R_{free} test set	1494 reflections (2.17%)	wwPDB-VP
Wilson B-factor $(Å^2)$	38.3	Xtriage
Anisotropy	0.338	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.37, 51.3	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.034 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5201	wwPDB-VP
Average B, all atoms $(Å^2)$	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: R5S, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.57	0/2510	0.70	1/3378~(0.0%)
1	В	0.57	0/2436	0.67	0/3277
All	All	0.57	0/4946	0.68	1/6655~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1
1	В	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	975	ARG	NE-CZ-NH2	-5.10	117.75	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	840	ASP	Peptide
1	В	859	ASN	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2473	0	2442	30	0
1	В	2402	0	2382	12	1
2	А	31	0	0	0	0
2	В	31	0	0	0	0
3	А	142	0	0	5	1
3	В	122	0	0	5	1
All	All	5201	0	4824	42	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	A00111-2	distance (Å)	overlap (Å)
1:B:1021:TYR:OH	3:B:1301:HOH:O	1.92	0.87
1:A:908:ASP:OD2	3:A:1301:HOH:O	1.97	0.81
1:A:840:ASP:O	3:A:1302:HOH:O	2.00	0.80
1:B:850:LYS:NZ	3:B:1303:HOH:O	2.18	0.76
1:A:942:GLN:HG2	1:A:1051:ILE:HB	1.68	0.74
1:B:877:GLU:OE2	3:B:1302:HOH:O	2.07	0.71
1:A:840:ASP:HB2	1:A:843:GLN:HB2	1.71	0.71
1:A:1052:GLU:O	3:A:1303:HOH:O	2.09	0.69
1:A:894:ASP:OD1	1:A:897:ARG:NH2	2.28	0.65
1:B:952:LYS:NZ	1:B:988:ASN:OD1	2.36	0.58
1:B:868:TYR:O	1:B:876:GLY:HA3	2.04	0.57
1:A:1072:GLN:O	1:A:1075:VAL:HG12	2.06	0.56
1:A:841:PRO:O	1:A:842:THR:OG1	2.16	0.56
1:A:989:ARG:HD3	3:A:1407:HOH:O	2.07	0.54
1:A:938:ARG:O	1:A:942:GLN:HG3	2.08	0.53
1:B:887:SER:O	1:B:887:SER:OG	2.21	0.52
1:A:885:GLN:HE21	1:A:885:GLN:HA	1.75	0.52
1:A:868:TYR:O	1:A:876:GLY:HA3	2.10	0.52
1:A:1104:GLU:HG2	1:A:1114:PRO:HD3	1.91	0.52
1:A:840:ASP:CB	1:A:843:GLN:HB2	2.41	0.51
1:A:1053:LYS:NZ	1:A:1054:SER:OG	2.26	0.51



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1104:GLU:OE2	3:B:1304:HOH:O	2.19	0.50
1:A:1053:LYS:CE	1:A:1054:SER:H	2.26	0.49
1:A:975:ARG:HD3	1:A:997:LEU:O	2.13	0.48
1:B:877:GLU:OE2	1:B:914:LYS:HE2	2.15	0.47
1:A:952:LYS:NZ	1:A:988:ASN:OD1	2.47	0.47
1:B:842:THR:HG22	1:B:913:TYR:OH	2.16	0.46
1:A:1053:LYS:N	1:A:1053:LYS:HE3	2.30	0.45
1:A:885:GLN:HG3	3:A:1354:HOH:O	2.17	0.45
1:A:1052:GLU:HG2	1:A:1053:LYS:CE	2.48	0.44
1:A:966:TYR:OH	1:A:970:LYS:NZ	2.48	0.43
1:A:920:ALA:HA	1:A:921:GLY:HA2	1.62	0.43
1:A:1068:ASP:OD1	1:A:1068:ASP:N	2.49	0.43
1:A:976:ASP:HB2	1:A:997:LEU:HD12	1.99	0.43
1:B:848:HIS:CG	1:B:870:PRO:HA	2.55	0.42
1:B:943:LYS:NZ	3:B:1312:HOH:O	2.53	0.42
1:A:857:LYS:HE2	1:A:885:GLN:OE1	2.21	0.41
1:B:1068:ASP:OD1	1:B:1068:ASP:N	2.50	0.41
1:A:1053:LYS:HE3	1:A:1054:SER:H	1.85	0.41
1:A:861:GLY:HA2	1:A:885:GLN:HG2	2.03	0.40
1:A:1053:LYS:CD	1:A:1054:SER:H	2.34	0.40
1:A:902:LEU:HG	1:A:913:TYR:HB2	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:883:LYS:NZ	1:B:1007:PTR:O3P[2_654]	1.30	0.90
3:B:1375:HOH:O	3:A:1369:HOH:O[3_554]	1.97	0.23

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	296/308~(96%)	288~(97%)	6(2%)	2(1%)	22	12
1	В	287/308~(93%)	281 (98%)	6(2%)	0	100	100
All	All	583/616~(95%)	569~(98%)	12 (2%)	2~(0%)	41	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	841	PRO
1	А	835	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric C		Percentiles
1	А	271/280~(97%)	265~(98%)	6(2%)	52 47
1	В	264/280~(94%)	261 (99%)	3 (1%)	73 73
All	All	535/560~(96%)	526~(98%)	9~(2%)	60 57

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	918	TYR
1	В	945	LYS
1	В	1080	GLU
1	А	840	ASP
1	А	885	GLN
1	А	906	GLN
1	А	918	TYR
1	А	922	ARG
1	А	1053	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mo	True	Chain Res		Link	Bo	ond leng	$_{\rm ths}$	В	ond ang	les
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	В	1007	1	$15,\!16,\!17$	1.31	1 (6%)	19,22,24	0.84	1 (5%)
1	PTR	А	1007	1	15,16,17	1.30	2 (13%)	19,22,24	0.96	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	В	1007	1	-	1/10/11/13	0/1/1/1
1	PTR	А	1007	1	-	2/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	1007	PTR	OH-CZ	-4.31	1.30	1.40
1	А	1007	PTR	OH-CZ	-3.82	1.32	1.40
1	А	1007	PTR	P-OH	2.05	1.62	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	1007	PTR	O2P-P-OH	3.22	115.31	105.24
1	В	1007	PTR	O3P-P-O2P	2.35	116.62	107.64



6VNJ

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	1007	PTR	CZ-OH-P-O1P
1	В	1007	PTR	CZ-OH-P-O1P
1	А	1007	PTR	CZ-OH-P-O3P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	В	1007	PTR	0	1

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Res	Link	Bond lengths			Bond angles		
Mol	Mol Type Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	R5S	А	1201	-	31,35,35	2.70	14 (45%)	34,48,48	1.57	9 (26%)
2	R5S	В	1201	-	31,35,35	2.68	12 (38%)	34,48,48	1.72	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R5S	А	1201	-	-	0/12/30/30	0/5/5/5



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R5S	В	1201	-	-	0/12/30/30	0/5/5/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
2	В	1201	R5S	C26-N30	8.12	1.49	1.36
2	А	1201	R5S	C26-N30	7.08	1.48	1.36
2	В	1201	R5S	C11-N12	6.54	1.49	1.36
2	А	1201	R5S	C31-N05	-6.05	1.29	1.35
2	А	1201	R5S	C11-N12	5.89	1.48	1.36
2	А	1201	R5S	C09-N10	-3.99	1.28	1.34
2	В	1201	R5S	C31-N05	-3.91	1.31	1.35
2	А	1201	R5S	C23-C24	-3.60	1.32	1.38
2	В	1201	R5S	C09-N10	-3.51	1.29	1.34
2	В	1201	R5S	C13-N12	3.48	1.48	1.40
2	А	1201	R5S	C13-N12	3.41	1.48	1.40
2	В	1201	R5S	C24-C13	3.25	1.44	1.39
2	В	1201	R5S	C14-C15	-3.14	1.33	1.38
2	А	1201	R5S	C14-C15	-3.00	1.33	1.38
2	А	1201	R5S	C03-C02	2.83	1.58	1.46
2	В	1201	R5S	C03-C02	2.81	1.58	1.46
2	В	1201	R5S	C23-C24	-2.51	1.34	1.38
2	В	1201	R5S	C26-N25	-2.43	1.30	1.34
2	А	1201	R5S	C24-C13	2.39	1.43	1.39
2	В	1201	R5S	C08-C09	2.34	1.51	1.49
2	А	1201	R5S	N06-N05	2.33	1.38	1.35
2	А	1201	R5S	C26-N25	-2.23	1.30	1.34
2	В	1201	R5S	C18-C17	-2.22	1.47	1.53
2	А	1201	R5S	C18-C17	-2.20	1.47	1.53
2	А	1201	R5S	C08-C09	2.17	1.51	1.49
2	А	1201	R5S	C15-C16	2.13	1.42	1.39

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	1201	R5S	C07-N06-N05	3.68	108.06	104.23
2	В	1201	R5S	N30-C26-N25	3.08	129.87	125.05
2	В	1201	R5S	C22-C17-C16	-3.07	105.60	112.79
2	В	1201	R5S	C29-C28-C27	3.05	106.99	103.19
2	А	1201	R5S	C14-C15-C16	-2.80	118.38	121.20
2	А	1201	R5S	C23-C16-C15	2.78	121.76	118.29
2	А	1201	R5S	C27-C09-N10	-2.76	118.23	122.78



6VNJ

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	1201	R5S	N30-C26-N25	2.56	129.06	125.05
2	В	1201	R5S	C23-C24-C13	-2.56	117.34	120.30
2	В	1201	R5S	C27-C09-N10	-2.53	118.61	122.78
2	В	1201	R5S	C23-C16-C15	2.38	121.26	118.29
2	А	1201	R5S	C19-N20-C21	2.36	117.11	110.34
2	А	1201	R5S	C07-N06-N05	2.31	106.63	104.23
2	В	1201	R5S	C14-C15-C16	-2.29	118.89	121.20
2	В	1201	R5S	C14-C13-N12	-2.19	113.30	120.64
2	А	1201	R5S	C23-C16-C17	-2.04	115.81	121.11
2	А	1201	R5S	C18-C17-C22	2.02	113.78	109.56
2	А	1201	R5S	C08-C09-N10	2.01	117.76	115.09

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sup Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2		$OWAB(Å^2)$	Q < 0.9	
1	А	298/308~(96%)	0.38	20 (6%)	17	20	31, 46, 81, 114	0
1	В	289/308~(93%)	0.30	22 (7%)	13	15	30, 45, 91, 112	0
All	All	587/616~(95%)	0.34	42 (7%)	15	17	30, 45, 85, 114	0

All (42) RSRZ outliers are listed below:

		Res Type		RSRZ
1	А	922	ARG	7.1
1	В	920	ALA	6.8
1	В	858 GLY		5.5
1	В	842	THR	5.5
1	В	922	ARG	5.5
1	В	918	TYR	4.7
1	В	1068	ASP	4.6
1	А	920	ALA	4.6
1	В	1067	ASN	4.5
1	А	921	GLY	4.4
1	В	859	ASN	4.2
1	А	1068	ASP	4.1
1	В	921	GLY	4.0
1	А	1053	LYS	3.9
1	В	1072	GLN	3.8
1	А	1132	GLY	3.5
1	В	843	GLN	3.4
1	В	919	SER	3.2
1	А	1070	GLN	3.1
1	А	918	TYR	3.1
1	В	1053	LYS	3.1
1	А	840	ASP	3.0
1	А	1067	ASN	3.0
1	В	1065	ILE	2.8



Mol	Chain	Res	Type	RSRZ	
1	А	923	ARG	2.8	
1	А	860	PHE	2.8	
1	А	834	ASN	2.8	
1	А	1011	LYS	2.7	
1	В	1131	ALA	2.7	
1	А	1072	GLN	2.5	
1	В	1070	GLN	2.5	
1	В	845	GLU	2.5	
1	В	1069	LYS	2.4	
1	В	1076	PHE	2.2	
1	А	859	ASN	2.2	
1	А	1069	LYS	2.2	
1	А	1003	GLN	2.1	
1	В	860	PHE	2.1	
1	В	923	ARG	2.0	
1	В	947	ARG	2.0	
1	А	984	VAL	2.0	
1	А	919	SER	2.0	

Continued from previous page...

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	$Q{<}0.9$
1	PTR	А	1007	16/17	0.92	0.12	48,59,92,97	0
1	PTR	В	1007	16/17	0.93	0.10	40,52,90,94	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q < 0.9
2	R5S	В	1201	31/31	0.97	0.11	$30,\!35,\!52,\!56$	0
2	R5S	А	1201	31/31	0.97	0.16	30,37,63,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

