



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2023 – 02:45 AM EDT

PDB ID : 6VQY
Title : HLA-B*27:05 presenting an HIV-1 7mer peptide
Authors : Pymm, P.; Tenzer, S.; Wee, E.; Weimershaus, M.; Burgevin, A.; Kollnberger, S.; Gerstoft, J.; Josephs, T.M.; Ladell, K.; McLaren, J.E.; Appay, V.; Price, D.A.; Fugger, L.; Bell, J.I.; Hansjorg, S.; Van Endert, P.; Harkiolaki, M.; Iversen, A.K.N.
Deposited on : 2020-02-06
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 275 | Total 2250 | C 1400 | N 408 | O 435 | S 7 | 0 | 1 | 0 |
| 1 | C | 275 | Total 2244 | C 1396 | N 407 | O 434 | S 7 | 0 | 0 | 0 |


- Molecule 2 is a protein called Beta-2-microglobulin.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|--------------|----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 2 | B | 99 | Total 829 | C 528 | N 140 | O 158 | S 3 | 0 | 0 | 0 |
| 2 | D | 99 | Total 829 | C 528 | N 140 | O 158 | S 3 | 0 | 0 | 0 |

- Molecule 3 is a protein called 7-mer peptide.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------------|---------|---------|--------|---------|---------|-------|
| | | | Total | C | N | O | | | |
| 3 | G | 7 | Total 62 | C 43 | N 12 | O 7 | 0 | 0 | 0 |
| 3 | F | 7 | Total 62 | C 43 | N 12 | O 7 | 0 | 0 | 0 |

- Molecule 4 is ARGININE (three-letter code: ARG) (formula: C₆H₁₅N₄O₂).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---|---------|---------|
| | | | Total | C | N | O | | |
| 4 | A | 1 | 11 | 6 | 4 | 1 | 0 | 0 |
| 4 | C | 1 | 11 | 6 | 4 | 1 | 0 | 0 |
| 4 | C | 1 | 11 | 6 | 4 | 1 | 0 | 0 |

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| | | | Total | C | O | | |
| 5 | A | 1 | 6 | 3 | 3 | 0 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---------|---------|
| 5 | C | 1 | Total | C | O | 0 | 0 |
| | | | 6 | 3 | 3 | | |

- Molecule 6 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 6 | A | 40 | Total | O | 0 | 0 |
| | | | 40 | 40 | | |
| 6 | B | 20 | Total | O | 0 | 0 |
| | | | 20 | 20 | | |
| 6 | C | 33 | Total | O | 0 | 0 |
| | | | 33 | 33 | | |
| 6 | D | 19 | Total | O | 0 | 0 |
| | | | 19 | 19 | | |
| 6 | G | 2 | Total | O | 0 | 0 |
| | | | 2 | 2 | | |
| 6 | F | 3 | Total | O | 0 | 0 |
| | | | 3 | 3 | | |

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics i

EDS failed to run properly - this section is therefore incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | P 1 21 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 45.35Å 129.86Å 90.26Å 90.00° 104.38° 90.00° | Depositor |
| Resolution (Å) | 87.44 – 2.57 | Depositor |
| % Data completeness (in resolution range) | 99.2 (87.44-2.57) | Depositor |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | 0.23 | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 336.89 (at 2.58Å) | Xtrriage |
| Refinement program | PHENIX 1.16_3549 | Depositor |
| R, R_{free} | 0.211 , 0.238 | Depositor |
| Wilson B-factor (Å ²) | 26.7 | Xtrriage |
| Anisotropy | 0.607 | Xtrriage |
| L-test for twinning ² | $\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$ | Xtrriage |
| Estimated twinning fraction | 0.327 for h,-k,-h-l | Xtrriage |
| Reported twinning fraction | 0.380 for h,-k,-h-l | Depositor |
| Outliers | 0 of 32006 reflections | Xtrriage |
| Total number of atoms | 6438 | wwPDB-VP |
| Average B, all atoms (Å ²) | 28.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 5 | GOL | C | 303 | - | 5,5,5 | 0.90 | 0 | 5,5,5 | 1.04 | 0 |
| 4 | ARG | A | 301 | - | 9,10,11 | 0.41 | 0 | 5,11,13 | 0.28 | 0 |
| 4 | ARG | C | 301 | - | 9,10,11 | 0.41 | 0 | 5,11,13 | 0.26 | 0 |
| 5 | GOL | A | 302 | - | 5,5,5 | 0.90 | 0 | 5,5,5 | 1.02 | 0 |
| 4 | ARG | C | 302 | - | 9,10,11 | 0.41 | 0 | 5,11,13 | 0.26 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|-------|
| 5 | GOL | C | 303 | - | - | 2/4/4/4 | - |
| 4 | ARG | A | 301 | - | - | 1/8/9/11 | - |
| 4 | ARG | C | 301 | - | - | 1/8/9/11 | - |
| 5 | GOL | A | 302 | - | - | 2/4/4/4 | - |
| 4 | ARG | C | 302 | - | - | 1/8/9/11 | - |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 4 | C | 302 | ARG | O-C-CA-CB |
| 5 | A | 302 | GOL | C1-C2-C3-O3 |
| 5 | C | 303 | GOL | C1-C2-C3-O3 |
| 5 | A | 302 | GOL | O2-C2-C3-O3 |
| 5 | C | 303 | GOL | O2-C2-C3-O3 |
| 4 | C | 301 | ARG | CA-CB-CG-CD |
| 4 | A | 301 | ARG | N-CA-CB-CG |

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.