

Full wwPDB X-ray Structure Validation Report (i)

Nov 5, 2023 – 03:06 PM EST

PDB ID	:	6VS3
Title	:	JAK2 JH1 in complex with BL2-057
Authors	:	Davis, R.R.; Schonbrunn, E.
Deposited on	:	2020-02-10
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	308	85%	8%	6%
1	В	308	3% 80%	15%	••••



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	А	289	Total 2395	C 1521	1	0 442	Р 1	S 14	0	0	0
1	В	298	Total 2465	C 1567	N 427	0 456	Р 1	S 14	0	0	0

• Molecule 1 is a protein called Tyrosine-protein kinase JAK2.


Chain	Residue	Modelled	Actual	Comment	Reference
А	825	HIS	-	expression tag	UNP O60674
А	826	HIS	-	expression tag	UNP O60674
А	827	HIS	-	expression tag	UNP O60674
А	828	HIS	-	expression tag	UNP O60674
А	829	HIS	-	expression tag	UNP O60674
A	830	HIS	-	expression tag	UNP O60674
А	831	HIS	-	expression tag	UNP O60674
А	832	HIS	-	expression tag	UNP O60674
А	833	GLU	-	expression tag	UNP O60674
А	834	ASN	-	expression tag	UNP O60674
А	835	LEU	-	expression tag	UNP O60674
А	836	TYR	-	expression tag	UNP O60674
А	837	PHE	-	expression tag	UNP O60674
А	838	GLN	-	expression tag	UNP O60674
А	839	GLY	-	expression tag	UNP O60674
В	825	HIS	-	expression tag	UNP O60674
В	826	HIS	-	expression tag	UNP O60674
В	827	HIS	-	expression tag	UNP O60674
В	828	HIS	-	expression tag	UNP O60674
В	829	HIS	-	expression tag	UNP O60674
В	830	HIS	-	expression tag	UNP O60674
В	831	HIS	-	expression tag	UNP O60674
В	832	HIS	-	expression tag	UNP O60674
В	833	GLU	-	expression tag	UNP O60674
В	834	ASN	-	expression tag	UNP O60674

There are 30 discrepancies between the modelled and reference sequences:



Continued from previous page						
Chain	Residue	Modelled	Actual	Comment	Reference	
В	835	LEU	-	expression tag	UNP O60674	
В	836	TYR	-	expression tag	UNP O60674	
В	837	PHE	-	expression tag	UNP O60674	
В	838	GLN	-	expression tag	UNP O60674	
В	839	GLY	-	expression tag	UNP O60674	

• Molecule 2 is (3R)-3-cyclopentyl-3-[4-(2-{[4-(piperidin-4-yl)phenyl]amino}-7H-pyrrolo [2,3-d]pyrimidin-4-yl)-1H-pyrazol-1-yl]propanenitrile (three-letter code: R6V) (formula: C₂₈H₃₂N₈) (labeled as "Ligand of Interest" by depositor).



\mathbf{N}	lol	Chain	Residues	Atoms	ZeroOcc	AltConf
	2	А	1	Total C N 36 28 8	0	0
	2	В	1	Total C N 36 28 8	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	174	Total O 174 174	0	0
3	В	199	Total O 199 199	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41	Depositor
Cell constants	112.10Å 112.10Å 70.18Å	Denesiten
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.80 - 2.00	Depositor
Resolution (A)	79.27 - 2.00	EDS
% Data completeness	99.9 (43.80 - 2.00)	Depositor
(in resolution range)	$100.0 \ (79.27-2.00)$	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.58 (at 2.00 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.14-3260_3260	Depositor
R, R_{free}	0.184 , 0.218	Depositor
n, nfree	0.184 , 0.218	DCC
R_{free} test set	2946 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	32.7	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 52.6	EDS
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5305	wwPDB-VP
Average B, all atoms $(Å^2)$	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: R6V, $\rm PTR$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain Bo		nd lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.45	0/2429	0.60	0/3267
1	В	0.50	1/2502~(0.0%)	0.64	3/3367~(0.1%)
All	All	0.47	1/4931~(0.0%)	0.62	3/6634~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	В	841	PRO	N-CD	5.65	1.55	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	1012	GLU	C-N-CD	6.37	141.78	128.40
1	В	840	ASP	C-N-CD	5.51	139.98	128.40
1	В	842	THR	N-CA-CB	-5.31	100.22	110.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	1012	GLU	Peptide



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2395	0	2377	13	2
1	В	2465	0	2438	38	0
2	А	36	0	0	0	0
2	В	36	0	0	0	0
3	А	174	0	0	1	1
3	В	199	0	0	6	1
All	All	5305	0	4815	51	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1052:GLU:OE1	3:B:1301:HOH:O	1.74	1.02
1:B:1012:GLU:HG2	1:B:1013:PRO:C	1.89	0.91
1:B:1010:VAL:CG1	1:B:1012:GLU:HB3	2.02	0.89
1:B:1012:GLU:HG2	1:B:1013:PRO:O	1.86	0.73
1:B:1010:VAL:HG13	1:B:1012:GLU:HB3	1.68	0.73
1:B:1112:GLN:O	3:B:1302:HOH:O	2.08	0.72
1:A:1062:MET:HB3	1:A:1067:ASN:OD1	1.91	0.71
1:A:846:GLU:OE2	1:A:919:SER:OG	2.12	0.67
1:B:1104:GLU:OE1	3:B:1303:HOH:O	2.12	0.67
1:B:1080:GLU:OE1	3:B:1304:HOH:O	2.12	0.65
1:B:1015:GLU:OE2	1:B:1015:GLU:HA	1.98	0.64
1:A:868:TYR:O	1:A:876:GLY:HA3	2.00	0.62
1:B:943:LYS:HD3	1:B:944:HIS:NE2	2.15	0.62
1:B:1012:GLU:CG	1:B:1014:GLY:HA3	2.31	0.61
1:A:952:LYS:NZ	1:A:988:ASN:OD1	2.34	0.61
1:A:859:ASN:O	1:A:860:PHE:CG	2.58	0.57
1:B:919:SER:HB2	1:B:923:ARG:HB2	1.87	0.56
1:B:855:LEU:HD21	1:B:865:MET:CE	2.38	0.54
1:B:845:GLU:OE2	1:B:847:ARG:NH2	2.41	0.53
1:B:868:TYR:O	1:B:876:GLY:HA3	2.09	0.53
1:B:999:LYS:NZ	3:B:1305:HOH:O	2.25	0.53



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:A:914:LYS:NZ	3:A:1303:HOH:O	2.42	0.52
1:A:943:LYS:HD3	1:A:944:HIS:NE2	2.27	0.49
1:B:891:HIS:HD2	3:B:1400:HOH:O	1.95	0.49
1:B:967:LEU:HD22	1:B:972:TYR:HB2	1.93	0.49
1:B:841:PRO:HG2	1:B:843:GLN:OE1	2.13	0.48
1:B:1081:LEU:CD2	1:B:1086:GLY:HA3	2.43	0.48
1:A:1012:GLU:HG2	1:A:1013:PRO:HD2	1.96	0.47
1:B:1081:LEU:HD23	1:B:1086:GLY:HA3	1.96	0.47
1:B:1012:GLU:HG2	1:B:1014:GLY:N	2.28	0.47
1:B:855:LEU:HD21	1:B:865:MET:HE2	1.95	0.47
1:B:847:ARG:HE	1:B:847:ARG:HB3	1.60	0.46
1:A:1070:GLN:O	1:A:1073:MET:HB2	2.15	0.46
1:A:848:HIS:CG	1:A:870:PRO:HA	2.51	0.46
1:B:920:ALA:HA	1:B:921:GLY:HA2	1.59	0.46
1:B:976:ASP:HB2	1:B:997:LEU:HD12	1.97	0.45
1:B:1065:ILE:HG22	1:B:1081:LEU:HD12	1.99	0.45
1:B:918:TYR:CD1	1:B:922:ARG:HA	2.51	0.45
1:B:855:LEU:HD11	1:B:865:MET:HB2	1.98	0.45
1:B:838:GLN:CG	1:B:839:GLY:HA3	2.47	0.44
1:B:965:GLU:O	1:B:969:THR:HG23	2.18	0.44
1:B:902:LEU:HG	1:B:913:TYR:HB2	1.99	0.43
1:A:903:LYS:HE2	1:A:913:TYR:CZ	2.54	0.43
1:B:850:LYS:N	1:B:850:LYS:HD2	2.34	0.42
1:B:1083:LYS:HB2	1:B:1083:LYS:HE3	1.80	0.42
1:B:942:GLN:HG3	1:B:945:LYS:NZ	2.34	0.42
1:B:1104:GLU:HG2	1:B:1114:PRO:HD3	2.02	0.42
1:A:1066:GLY:O	1:A:1069:LYS:HG2	2.19	0.42
1:A:967:LEU:HD23	1:A:967:LEU:HA	1.87	0.41
1:B:1057:PRO:HB2	1:B:1058:PRO:HD3	2.02	0.41
1:B:838:GLN:HG3	1:B:839:GLY:HA3	2.03	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:883:LYS:NZ	1:A:1007:PTR:O3P[2_655]	1.30	0.90
1:A:883:LYS:CE	1:A:1007:PTR:O3P[2_655]	2.08	0.12
3:A:1333:HOH:O	3:B:1464:HOH:O[1_556]	2.12	0.08



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	286/308~(93%)	279~(98%)	7 (2%)	0	100 100
1	В	295/308~(96%)	287~(97%)	6~(2%)	2(1%)	22 16
All	All	581/616~(94%)	566 (97%)	13~(2%)	2 (0%)	41 37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	841	PRO
1	В	839	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	263/280~(94%)	262 (100%)	1 (0%)	91 93
1	В	270/280~(96%)	266~(98%)	4 (2%)	65 69
All	All	533/560~(95%)	528 (99%)	5 (1%)	78 83

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	918	TYR
1	В	843	GLN
1	В	850	LYS
1	В	1011	LYS



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type
1	В	1012	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	1070	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Dec	Tinle	Link Bond lengths			В	ond ang	les
10101	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	PTR	А	1007	1	$15,\!16,\!17$	1.18	1 (6%)	$19,\!22,\!24$	0.64	1 (5%)
1	PTR	В	1007	1	$15,\!16,\!17$	1.22	1 (6%)	19,22,24	0.64	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	А	1007	1	-	1/10/11/13	0/1/1/1
1	PTR	В	1007	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(\text{\AA})$	Ideal(Å)
1	В	1007	PTR	OH-CZ	-4.22	1.31	1.40
1	А	1007	PTR	OH-CZ	-4.12	1.31	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	1007	PTR	O2P-P-OH	2.42	112.82	105.24
1	А	1007	PTR	O2P-P-OH	2.16	111.98	105.24

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	1007	PTR	CZ-OH-P-O2P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	А	1007	PTR	0	2

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trune	Chain	Dec	Tinle	B	Bond lengths			Bond angles		
WIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
2	R6V	В	1201	-	37,41,41	2.17	12 (32%)	38,57,57	1.98	11 (28%)	
2	R6V	А	1201	-	37,41,41	2.21	13 (35%)	38,57,57	1.93	9 (23%)	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R6V	В	1201	-	-	1/14/38/38	0/6/6/6
2	R6V	А	1201	-	-	0/14/38/38	0/6/6/6

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	А	1201	R6V	C06-C05	-5.60	1.39	1.53
2	В	1201	R6V	C06-C05	-5.39	1.39	1.53
2	А	1201	R6V	C09-C05	5.37	1.67	1.53
2	В	1201	R6V	C09-C05	5.12	1.67	1.53
2	А	1201	R6V	C17-N18	4.58	1.45	1.36
2	В	1201	R6V	C17-N18	4.52	1.45	1.36
2	В	1201	R6V	C03-C02	3.63	1.54	1.46
2	А	1201	R6V	C08-C09	-3.22	1.38	1.51
2	А	1201	R6V	C29-C22	3.15	1.44	1.39
2	А	1201	R6V	C13-N10	-3.14	1.32	1.35
2	В	1201	R6V	C14-C15	3.07	1.53	1.49
2	В	1201	R6V	C13-N10	-3.07	1.32	1.35
2	А	1201	R6V	C03-C02	3.04	1.52	1.46
2	В	1201	R6V	C08-C09	-3.01	1.39	1.51
2	А	1201	R6V	C14-C15	2.94	1.53	1.49
2	А	1201	R6V	C19-N18	2.67	1.46	1.40
2	В	1201	R6V	C29-C22	2.66	1.43	1.39
2	А	1201	R6V	C30-C19	2.53	1.43	1.39
2	В	1201	R6V	C19-N18	2.44	1.46	1.40
2	А	1201	R6V	C33-C32	-2.42	1.36	1.43
2	А	1201	R6V	C05-C04	2.40	1.58	1.52
2	В	1201	R6V	C32-N36	2.36	1.39	1.34
2	В	1201	R6V	C07-C06	2.23	1.61	1.51
2	А	1201	R6V	C07-C06	2.19	1.60	1.51
2	В	1201	R6V	C33-C32	-2.07	1.37	1.43

All (25) bond length outliers are listed below:

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	1201	R6V	C24-C23-C22	-6.52	97.52	112.79
2	А	1201	R6V	C24-C23-C22	-6.48	97.60	112.79
2	В	1201	R6V	C17-N31-C32	4.37	120.24	115.28



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	А	1201	R6V	C13-N10-C04	-3.72	122.19	125.48
2	А	1201	R6V	C17-N31-C32	3.70	119.48	115.28
2	А	1201	R6V	C09-C05-C04	-3.33	106.19	112.73
2	В	1201	R6V	C09-C05-C04	-3.26	106.32	112.73
2	В	1201	R6V	C28-C23-C24	3.16	116.14	109.56
2	А	1201	R6V	C33-C15-N16	-3.14	119.08	121.95
2	В	1201	R6V	C28-C23-C22	3.11	120.07	112.79
2	В	1201	R6V	C33-C15-N16	-2.98	119.23	121.95
2	А	1201	R6V	C28-C23-C24	2.88	115.57	109.56
2	В	1201	R6V	C09-C05-C06	2.36	108.94	102.91
2	А	1201	R6V	C20-C19-N18	-2.17	113.37	120.64
2	А	1201	R6V	N31-C17-N16	-2.17	122.80	126.23
2	В	1201	R6V	C04-C03-C02	-2.09	108.94	112.12
2	В	1201	R6V	C29-C22-C23	-2.09	115.69	121.11
2	В	1201	R6V	C13-N10-C04	-2.03	123.68	125.48
2	А	1201	R6V	C09-C05-C06	2.02	108.08	102.91
2	В	1201	R6V	N31-C17-N16	-2.01	123.05	126.23

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	В	1201	R6V	C21-C22-C23-C24

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	288/308~(93%)	0.27	16 (5%) 24 23	23, 36, 79, 102	0
1	В	297/308~(96%)	0.16	9 (3%) 50 49	26, 39, 73, 102	0
All	All	585/616~(94%)	0.22	25 (4%) 35 34	23, 38, 78, 102	0

All (25) RSRZ outliers are listed below:

1 A 9 1 A 8 1 A 8 1 A 10 1 A 10 1 B 8 1 B 9 1 B 9 1 B 8 1 A 10 1 A 10 1 A 8	920 ALA 5.6 921 GLY 4.8 860 PHE 4.6 886 HIS 4.3 070 GLN 4.1 840 ASP 3.8 922 ARG 3.4 839 GLY 3.3 067 ASN 3.2
1 A 8 1 A 8 1 A 10 1 B 8 1 B 9 1 B 8 1 A 10 1 A 10 1 A 10 1 A 10 1 A 8 1 A 8	860 PHE 4.6 886 HIS 4.3 070 GLN 4.1 840 ASP 3.8 922 ARG 3.4 839 GLY 3.3
1 A 80 1 A 10 1 B 80 1 B 90 1 B 80 1 B 80 1 A 10 1 A 10 1 A 10 1 A 80	886 HIS 4.3 070 GLN 4.1 840 ASP 3.8 922 ARG 3.4 839 GLY 3.3
1 A 10 1 B 8 1 B 9 1 B 8 1 A 10 1 A 10 1 A 8 1 A 8	070 GLN 4.1 840 ASP 3.8 922 ARG 3.4 839 GLY 3.3
1 B 8 1 B 9 1 B 8 1 A 10 1 A 8	840 ASP 3.8 922 ARG 3.4 839 GLY 3.3
1 B 9 1 B 8 1 A 10 1 A 8	922 ARG 3.4 839 GLY 3.3
1 B 8 1 A 10 1 A 8	839 GLY 3.3
1 A 10 1 A 8	
1 A 8	067 ASN 3.2
1 A 10	890 GLU 3.2
	068 ASP 3.2
1 B 11	132 GLY 3.1
1 A 99	922 ARG 2.9
1 B 8	838 GLN 2.9
1 A 8	889 GLU 2.7
1 A 8	859 ASN 2.7
1 A 10	076 PHE 2.7
1 B 9	918 TYR 2.6
1 B 10	012 GLU 2.6
1 A 9	918 TYR 2.6
1 A 9	919 SER 2.4
1 B 84	841 PRO 2.4
1 A 10	065 ILE 2.4
1 B 10	



Mol	Chain	\mathbf{Res}	Type	RSRZ
1	А	893	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
1	PTR	В	1007	16/17	0.91	0.13	42,59,85,89	0
1	PTR	А	1007	16/17	0.93	0.12	30,43,90,96	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	R6V	В	1201	36/36	0.97	0.15	20,32,66,71	0
2	R6V	А	1201	36/36	0.98	0.13	21,28,43,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

