

Full wwPDB EM Validation Report (i)

Jul 10, 2024 – 02:09 PM EDT

PDB ID : 8VSU

EMDB ID : EMD-43506

Title : Cryo-EM structure of LKB1-STRADalpha-MO25alpha heterocomplex

Authors: Chan, L.M.; Courteau, B.J.; Verba, K.A.

Deposited on : 2024-01-24

Resolution : 2.86 Å(reported)

Based on initial models : 2WTK, .

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at $\frac{\text{https://www.wwpdb.org/validation/2017/EMValidationReportHelp}}{\text{with specific help available everywhere you see the } (i) symbol.$

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.13$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$		
Clashscore	158937	4297		
Ramachandran outliers	154571	4023		
Sidechain outliers	154315	3826		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	362	88%	• 9%				
2	С	449	67%	32%				
3	В	429	9% 69%	29%				



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 15240 atoms, of which 7646 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Calcium-binding protein 39.

Mol	Chain	Residues	Atoms						AltConf	Trace	
1	A	328	Total 5458	C 1736	H 2751	N 455	O 505	S 11	0	0	

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	342	GLY	-	expression tag	UNP Q9Y376
A	343	SER	-	expression tag	UNP Q9Y376
A	344	GLY	-	expression tag	UNP Q9Y376
A	345	ALA	-	expression tag	UNP Q9Y376
A	346	THR	-	expression tag	UNP Q9Y376
A	347	ASN	-	expression tag	UNP Q9Y376
A	348	PHE	-	expression tag	UNP Q9Y376
A	349	SER	-	expression tag	UNP Q9Y376
A	350	LEU	-	expression tag	UNP Q9Y376
A	351	LEU	-	expression tag	UNP Q9Y376
A	352	LYS	-	expression tag	UNP Q9Y376
A	353	GLN	-	expression tag	UNP Q9Y376
A	354	ALA	-	expression tag	UNP Q9Y376
A	355	GLY	-	expression tag	UNP Q9Y376
A	356	ASP	-	expression tag	UNP Q9Y376
A	357	VAL	-	expression tag	UNP Q9Y376
A	358	GLU	-	expression tag	UNP Q9Y376
A	359	GLU		expression tag	UNP Q9Y376
A	360	ASN	-	expression tag	UNP Q9Y376
A	361	PRO	-	expression tag	UNP Q9Y376
A	362	GLY	-	expression tag	UNP Q9Y376

• Molecule 2 is a protein called Serine/threonine-protein kinase STK11.

Mol	Chain	Residues		Atoms					AltConf	Trace
2	С	307	Total	С	Н	N	О	S	0	0
		301	4900	1555	2470	421	439	15		



There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	-15	PRO	-	expression tag	UNP Q15831
С	-14	ASP	-	expression tag	UNP Q15831
С	-13	TYR	-	expression tag	UNP Q15831
С	-12	LYS	-	expression tag	UNP Q15831
С	-11	ASP	-	expression tag	UNP Q15831
С	-10	ASP	-	expression tag	UNP Q15831
С	-9	ASP	-	expression tag	UNP Q15831
С	-8	ASP	-	expression tag	UNP Q15831
С	-7	LYS	-	expression tag	UNP Q15831
С	-6	GLU	-	expression tag	UNP Q15831
С	-5	ASN	-	expression tag	UNP Q15831
С	-4	LEU	-	expression tag	UNP Q15831
С	-3	TYR	-	expression tag	UNP Q15831
С	-2	PHE	-	expression tag	UNP Q15831
С	-1	GLN	-	expression tag	UNP Q15831
С	0	GLY	-	expression tag	UNP Q15831

• Molecule 3 is a protein called Isoform 3 of STE20-related kinase adapter protein alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace	
3	В	306	Total 4855	C 1558	H 2425	N 420	O 437	S 15	0	0

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	24	PRO	-	expression tag	UNP Q7RTN6
В	25	HIS	-	expression tag	UNP Q7RTN6
В	26	HIS	-	expression tag	UNP Q7RTN6
В	27	HIS	-	expression tag	UNP Q7RTN6
В	28	HIS	-	expression tag	UNP Q7RTN6
В	29	HIS	-	expression tag	UNP Q7RTN6
В	30	HIS	-	expression tag	UNP Q7RTN6
В	31	GLU	-	expression tag	UNP Q7RTN6
В	32	ASN	-	expression tag	UNP Q7RTN6
В	33	LEU	-	expression tag	UNP Q7RTN6
В	34	TYR	-	expression tag	UNP Q7RTN6
В	35	PHE	-	expression tag	UNP Q7RTN6
В	36	GLN	-	expression tag	UNP Q7RTN6
В	37	GLY	-	expression tag	UNP Q7RTN6
В	432	GLY	-	expression tag	UNP Q7RTN6


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
В	433	SER	-	expression tag	UNP Q7RTN6
В	434	GLY	-	expression tag	UNP Q7RTN6
В	435	ALA	-	expression tag	UNP Q7RTN6
В	436	THR	-	expression tag	UNP Q7RTN6
В	437	ASN	-	expression tag	UNP Q7RTN6
В	438	PHE	-	expression tag	UNP Q7RTN6
В	439	SER	-	expression tag	UNP Q7RTN6
В	440	LEU	-	expression tag	UNP Q7RTN6
В	441	LEU	-	expression tag	UNP Q7RTN6
В	442	LYS	-	expression tag	UNP Q7RTN6
В	443	GLN	-	expression tag	UNP Q7RTN6
В	444	ALA	-	expression tag	UNP Q7RTN6
В	445	GLY	-	expression tag	UNP Q7RTN6
В	446	ASP	-	expression tag	UNP Q7RTN6
В	447	VAL	-	expression tag	UNP Q7RTN6
В	448	GLU	-	expression tag	UNP Q7RTN6
В	449	GLU	-	expression tag	UNP Q7RTN6
В	450	ASN	-	expression tag	UNP Q7RTN6
В	451	PRO	-	expression tag	UNP Q7RTN6
В	452	GLY	_	expression tag	UNP Q7RTN6

• Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues		Ato	oms			AltConf
4	D	1	Total	С	N	О	Р	0
4	Ъ	1	27	10	5	10	2	U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Calcium-binding protein 39









4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	124780	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	45.8	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	4.145	Depositor
Minimum map value	-2.213	Depositor
Average map value	-0.001	Depositor
Map value standard deviation	0.102	Depositor
Recommended contour level	0.628	Depositor
Map size (Å)	213.76, 213.76, 213.76	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.835, 0.835, 0.835	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
WIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.62	0/2755	0.89	7/3707 (0.2%)	
2	С	0.64	0/2487	0.98	6/3358~(0.2%)	
3	В	0.63	0/2492	0.96	9/3383~(0.3%)	
All	All	0.63	0/7734	0.94	22/10448~(0.2%)	

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	С	304	ARG	NE-CZ-NH1	8.78	124.69	120.30
2	С	333	ARG	NE-CZ-NH1	8.61	124.61	120.30
2	С	74	ARG	NE-CZ-NH1	7.97	124.29	120.30
3	В	89	ARG	NE-CZ-NH1	7.75	124.18	120.30
3	В	101	ARG	NE-CZ-NH1	7.22	123.91	120.30
3	В	100	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	A	264	ARG	NE-CZ-NH1	6.87	123.73	120.30
3	В	134	ARG	NE-CZ-NH1	6.25	123.42	120.30
1	A	107	ARG	NE-CZ-NH1	6.23	123.42	120.30
3	В	215	ARG	NE-CZ-NH1	6.15	123.37	120.30
2	С	297	ARG	NE-CZ-NH1	5.99	123.30	120.30
3	В	384	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	A	313	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	A	240	ARG	NE-CZ-NH1	5.45	123.03	120.30
2	С	42	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	A	171	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	A	268	ARG	NE-CZ-NH1	5.23	122.91	120.30
2	С	331	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	A	108	ARG	NE-CZ-NH1	5.16	122.88	120.30
3	В	362	ARG	NE-CZ-NH1	5.12	122.86	120.30
3	В	395	ARG	NE-CZ-NH1	5.04	122.82	120.30

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
3	В	194	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2707	2751	2750	1	0
2	С	2430	2470	2469	0	0
3	В	2430	2425	2423	1	0
4	В	27	0	12	0	0
All	All	7594	7646	7654	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:141:LEU:HD22	3:B:123:LYS:HB3	1.86	0.58

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$326/362 \ (90\%)$	323 (99%)	3 (1%)	0	100	100
2	С	305/449 (68%)	300 (98%)	5 (2%)	0	100	100
3	В	$302/429 \ (70\%)$	293 (97%)	9 (3%)	0	100	100
All	All	933/1240 (75%)	916 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	302/327~(92%)	302 (100%)	0	100	100
2	С	265/388~(68%)	265 (100%)	0	100	100
3	В	271/378 (72%)	270 (100%)	1 (0%)	91	96
All	All	838/1093 (77%)	837 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	В	390	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	В	2001	-	24,29,29	1.20	2 (8%)	29,45,45	1.46	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	В	2001	-	-	1/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
4	В	2001	ADP	O4'-C1'	3.16	1.45	1.41
4	В	2001	ADP	C5-C4	-2.11	1.35	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
4	В	2001	ADP	C4-C5-N7	4.33	113.92	109.40
4	В	2001	ADP	O2B-PB-O3A	2.78	113.97	104.64
4	В	2001	ADP	N6-C6-N1	-2.35	113.70	118.57
4	В	2001	ADP	PA-O3A-PB	-2.25	125.11	132.83
4	В	2001	ADP	C2-N1-C6	-2.18	115.03	118.75
4	В	2001	ADP	O3B-PB-O3A	2.03	111.45	104.64



There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	2001	ADP	PA-O3A-PB-O1B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-43506. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map





The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

6.2.1 Primary map









X Index: 128


Y Index: 128

Z Index: 128

6.2.2 Raw map







X Index: 128

Y Index: 128


Z Index: 128


The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map









X Index: 131


Y Index: 131

Z Index: 112

6.3.2 Raw map







X Index: 115

Y Index: 109


Z Index: 112

The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map




The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.628. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $31~\mathrm{nm^3}$; this corresponds to an approximate mass of $28~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.350 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.350 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	2.86	-	-
Author-provided FSC curve	2.83	3.26	2.87
Unmasked-calculated*	3.42	3.94	3.50


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.42 differs from the reported value 2.86 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-43506 and PDB model 8VSU. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.628 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.628).



9.4 Atom inclusion (i)



At the recommended contour level, 74% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.628) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7340	0.5750
A	0.7250	0.5670
В	0.7370	0.5800
С	0.7590	0.5800



