



wwPDB X-ray Structure Validation Summary Report (i)

May 18, 2020 – 11:29 pm BST

PDB ID : 6VW8
Title : Formate Dehydrogenase FdsABG subcomplex FdsBG from C. necator
Authors : Young, T.
Deposited on : 2020-02-18
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see [references \(1\)](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 10508 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent formate dehydrogenase gamma subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C 1123	N 701	O 208	S 207	7	0	0
1	C	152	Total	C 1129	N 704	O 209	S 209	7	0	0

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	MET	-	initiating methionine	UNP Q0KDY3
A	-38	GLY	-	expression tag	UNP Q0KDY3
A	-37	GLY	-	expression tag	UNP Q0KDY3
A	-36	SER	-	expression tag	UNP Q0KDY3
A	-35	HIS	-	expression tag	UNP Q0KDY3
A	-34	HIS	-	expression tag	UNP Q0KDY3
A	-33	HIS	-	expression tag	UNP Q0KDY3
A	-32	HIS	-	expression tag	UNP Q0KDY3
A	-31	HIS	-	expression tag	UNP Q0KDY3
A	-30	HIS	-	expression tag	UNP Q0KDY3
A	-29	SER	-	expression tag	UNP Q0KDY3
A	-28	ARG	-	expression tag	UNP Q0KDY3
A	-27	ALA	-	expression tag	UNP Q0KDY3
A	-26	TRP	-	expression tag	UNP Q0KDY3
A	-25	ARG	-	expression tag	UNP Q0KDY3
A	-24	HIS	-	expression tag	UNP Q0KDY3
A	-23	PRO	-	expression tag	UNP Q0KDY3
A	-22	GLN	-	expression tag	UNP Q0KDY3
A	-21	GLY	-	expression tag	UNP Q0KDY3
A	-20	GLY	-	expression tag	UNP Q0KDY3
A	-19	SER	-	expression tag	UNP Q0KDY3
A	-18	HIS	-	expression tag	UNP Q0KDY3
A	-17	HIS	-	expression tag	UNP Q0KDY3
A	-16	HIS	-	expression tag	UNP Q0KDY3
A	-15	HIS	-	expression tag	UNP Q0KDY3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	HIS	-	expression tag	UNP Q0KDY3
A	-13	HIS	-	expression tag	UNP Q0KDY3
A	-12	GLY	-	expression tag	UNP Q0KDY3
A	-11	MET	-	expression tag	UNP Q0KDY3
A	-10	ARG	-	expression tag	UNP Q0KDY3
A	-9	ASP	-	expression tag	UNP Q0KDY3
A	-8	LEU	-	expression tag	UNP Q0KDY3
A	-7	TYR	-	expression tag	UNP Q0KDY3
A	-6	ASP	-	expression tag	UNP Q0KDY3
A	-5	ASP	-	expression tag	UNP Q0KDY3
A	-4	ASP	-	expression tag	UNP Q0KDY3
A	-3	ASP	-	expression tag	UNP Q0KDY3
A	-2	LYS	-	expression tag	UNP Q0KDY3
A	-1	ASP	-	expression tag	UNP Q0KDY3
A	0	PRO	-	expression tag	UNP Q0KDY3
C	-39	MET	-	initiating methionine	UNP Q0KDY3
C	-38	GLY	-	expression tag	UNP Q0KDY3
C	-37	GLY	-	expression tag	UNP Q0KDY3
C	-36	SER	-	expression tag	UNP Q0KDY3
C	-35	HIS	-	expression tag	UNP Q0KDY3
C	-34	HIS	-	expression tag	UNP Q0KDY3
C	-33	HIS	-	expression tag	UNP Q0KDY3
C	-32	HIS	-	expression tag	UNP Q0KDY3
C	-31	HIS	-	expression tag	UNP Q0KDY3
C	-30	HIS	-	expression tag	UNP Q0KDY3
C	-29	SER	-	expression tag	UNP Q0KDY3
C	-28	ARG	-	expression tag	UNP Q0KDY3
C	-27	ALA	-	expression tag	UNP Q0KDY3
C	-26	TRP	-	expression tag	UNP Q0KDY3
C	-25	ARG	-	expression tag	UNP Q0KDY3
C	-24	HIS	-	expression tag	UNP Q0KDY3
C	-23	PRO	-	expression tag	UNP Q0KDY3
C	-22	GLN	-	expression tag	UNP Q0KDY3
C	-21	GLY	-	expression tag	UNP Q0KDY3
C	-20	GLY	-	expression tag	UNP Q0KDY3
C	-19	SER	-	expression tag	UNP Q0KDY3
C	-18	HIS	-	expression tag	UNP Q0KDY3
C	-17	HIS	-	expression tag	UNP Q0KDY3
C	-16	HIS	-	expression tag	UNP Q0KDY3
C	-15	HIS	-	expression tag	UNP Q0KDY3
C	-14	HIS	-	expression tag	UNP Q0KDY3
C	-13	HIS	-	expression tag	UNP Q0KDY3

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-12	GLY	-	expression tag	UNP Q0KDY3
C	-11	MET	-	expression tag	UNP Q0KDY3
C	-10	ARG	-	expression tag	UNP Q0KDY3
C	-9	ASP	-	expression tag	UNP Q0KDY3
C	-8	LEU	-	expression tag	UNP Q0KDY3
C	-7	TYR	-	expression tag	UNP Q0KDY3
C	-6	ASP	-	expression tag	UNP Q0KDY3
C	-5	ASP	-	expression tag	UNP Q0KDY3
C	-4	ASP	-	expression tag	UNP Q0KDY3
C	-3	ASP	-	expression tag	UNP Q0KDY3
C	-2	LYS	-	expression tag	UNP Q0KDY3
C	-1	ASP	-	expression tag	UNP Q0KDY3
C	0	PRO	-	expression tag	UNP Q0KDY3

- Molecule 2 is a protein called NAD-dependent formate dehydrogenase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	513	Total	C	N	O	S	0	0	0
			3825	2407	676	720	22			
2	D	514	Total	C	N	O	S	0	0	0
			3836	2415	677	721	23			

- Molecule 3 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 4	Fe 2	S 2	0	0
3	C	1	Total 4	Fe 2	S 2	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 1	K 1		0	0
4	C	1	Total 1	K 1		0	0

- Molecule 5 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	B	1	Total 31	17	4	9	1	0	0
5	D	1	Total 31	17	4	9	1	0	0

- Molecule 6 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Fe S 8 4 4	0	0
6	D	1	Total Fe S 8 4 4	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total O S 5 4 1	0	0
7	B	1	Total O S 5 4 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	45	Total O 45 45	0	0
8	B	226	Total O 226 226	0	0
8	C	51	Total O 51 51	0	0
8	D	175	Total O 175 175	0	0

3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD-dependent formate dehydrogenase gamma subunit




- Molecule 1: NAD-dependent formate dehydrogenase gamma subunit




- Molecule 2: NAD-dependent formate dehydrogenase beta subunit



- Molecule 2: NAD-dependent formate dehydrogenase beta subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	132.66 Å 127.05 Å 99.84 Å 90.00° 123.50° 90.00°	Depositor
Resolution (Å)	48.53 – 2.30 48.53 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.53-2.30) 99.8 (48.53-2.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.05 (at 2.29 Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R , R_{free}	0.182 , 0.223 0.182 , 0.223	Depositor DCC
R_{free} test set	1996 reflections (3.27%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.285	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.4	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10508	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, SF4, K, FES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.42	0/1145	0.62	0/1561
1	C	0.48	0/1151	0.62	0/1569
2	B	0.51	0/3896	0.63	0/5277
2	D	0.46	0/3907	0.61	0/5291
All	All	0.48	0/10099	0.62	0/13698

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1123	0	1088	14	0
1	C	1129	0	1093	8	0
2	B	3825	0	3794	42	0
2	D	3836	0	3815	43	0
3	A	4	0	0	1	0
3	C	4	0	0	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	B	31	0	19	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	31	0	19	2	0
6	B	8	0	0	0	0
6	D	8	0	0	0	0
7	B	10	0	0	0	0
8	A	45	0	0	2	0
8	B	226	0	0	3	0
8	C	51	0	0	1	0
8	D	175	0	0	3	0
All	All	10508	0	9828	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:293:PRO:CD	2:B:294:PRO:HD3	1.78	1.13
2:B:293:PRO:HD2	2:B:294:PRO:CD	1.87	1.04
2:B:293:PRO:HD2	2:B:294:PRO:HD3	1.01	1.01
2:B:293:PRO:CD	2:B:294:PRO:CD	2.52	0.80
2:B:25:ARG:HH21	2:D:29:ARG:HH12	1.33	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	149/215 (69%)	145 (97%)	3 (2%)	1 (1%)	22 26
1	C	150/215 (70%)	146 (97%)	3 (2%)	1 (1%)	22 26
2	B	511/520 (98%)	490 (96%)	19 (4%)	2 (0%)	34 42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	D	512/520 (98%)	494 (96%)	17 (3%)	1 (0%)	47 58
All	All	1322/1470 (90%)	1275 (96%)	42 (3%)	5 (0%)	34 42

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	185	GLU
2	B	185	GLU
1	A	127	CYS
2	B	294	PRO
1	C	127	CYS

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	111/165 (67%)	109 (98%)	2 (2%)	59 75
1	C	112/165 (68%)	110 (98%)	2 (2%)	59 75
2	B	385/391 (98%)	379 (98%)	6 (2%)	62 78
2	D	387/391 (99%)	382 (99%)	5 (1%)	69 82
All	All	995/1112 (90%)	980 (98%)	15 (2%)	65 79

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	455	ARG
2	B	489	CYS
2	D	356	LYS
2	B	450	ARG
2	D	281	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SF4	B	602	2	0,12,12	0.00	-	-		
6	SF4	D	601	2	0,12,12	0.00	-	-		
3	FES	C	301	1	0,4,4	0.00	-	-		
7	SO4	B	603	-	4,4,4	0.14	0	6,6,6	0.35	0
3	FES	A	301	1	0,4,4	0.00	-	-		
7	SO4	B	604	-	4,4,4	0.19	0	6,6,6	0.14	0
5	FMN	B	601	-	31,33,33	4.15	12 (38%)	40,50,50	2.27	9 (22%)
5	FMN	D	600	-	31,33,33	4.22	10 (32%)	40,50,50	2.18	13 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SF4	B	602	2	-	-	0/6/5/5
6	SF4	D	601	2	-	-	0/6/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FES	C	301	1	-	-	0/1/1/1
3	FES	A	301	1	-	-	0/1/1/1
5	FMN	B	601	-	-	3/18/18/18	0/3/3/3
5	FMN	D	600	-	-	2/18/18/18	0/3/3/3

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	601	FMN	C10-N1	10.64	1.46	1.33
5	D	600	FMN	C10-N1	10.56	1.46	1.33
5	B	601	FMN	C9A-N10	9.23	1.51	1.38
5	D	600	FMN	C9A-N10	8.91	1.50	1.38
5	D	600	FMN	C4A-N5	8.73	1.45	1.33

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	601	FMN	C1'-N10-C9A	6.76	123.61	118.29
5	D	600	FMN	C1'-N10-C9A	5.98	123.00	118.29
5	B	601	FMN	C4-N3-C2	5.26	119.58	115.14
5	B	601	FMN	C7M-C7-C8	5.16	131.32	120.74
5	D	600	FMN	C4-N3-C2	5.10	119.45	115.14

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	601	FMN	C1'-C2'-C3'-C4'
5	B	601	FMN	C4'-C5'-O5'-P
5	D	600	FMN	C4'-C5'-O5'-P
5	B	601	FMN	C1'-C2'-C3'-O3'
5	D	600	FMN	N10-C1'-C2'-O2'

There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	301	FES	1	0
3	A	301	FES	1	0
5	B	601	FMN	2	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	600	FMN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/215 (70%)	0.27	13 (8%) 10 14	21, 34, 63, 85	0
1	C	152/215 (70%)	0.10	5 (3%) 46 53	19, 35, 50, 79	0
2	B	513/520 (98%)	-0.15	4 (0%) 86 89	16, 27, 45, 79	0
2	D	514/520 (98%)	-0.09	7 (1%) 75 80	19, 32, 49, 70	0
All	All	1330/1470 (90%)	-0.05	29 (2%) 62 69	16, 31, 52, 85	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	293	PRO	4.6
1	A	116	ALA	3.4
2	B	293	PRO	3.1
1	C	108	GLY	2.9
1	A	156	VAL	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no carbohydrates in this entry.


6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SO4	B	603	5/5	0.95	0.14	40,43,44,45	0
7	SO4	B	604	5/5	0.95	0.10	58,58,59,60	0
4	K	A	302	1/1	0.96	0.08	53,53,53,53	0
5	FMN	D	600	31/31	0.96	0.15	21,26,33,34	0
5	FMN	B	601	31/31	0.97	0.12	18,22,23,24	0
6	SF4	B	602	8/8	0.98	0.08	27,30,32,38	0
6	SF4	D	601	8/8	0.98	0.08	24,27,30,32	0
4	K	C	302	1/1	0.98	0.10	37,37,37,37	0
3	FES	A	301	4/4	0.99	0.13	26,26,28,28	0
3	FES	C	301	4/4	0.99	0.12	22,23,24,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.