



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 21, 2023 – 07:28 PM JST

PDB ID : 7VWP  
Title : Structure of the flavin-dependent monooxygenase FlsO1 from the biosynthesis of fluostatinsin  
Authors : Zhang, Y.; Yang, C.; Zhang, L.; Zhang, C.  
Deposited on : 2021-11-11  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	497	 4% (poor fit) 79% (0 outliers) 17% (1 outlier) .. (2 outliers)
1	B	497	 3% (poor fit) 83% (0 outliers) 15% (1 outlier) .. (2 outliers)
1	C	497	 4% (poor fit) 76% (0 outliers) 18% (1 outlier) . 5% (2 outliers)
1	D	497	 2% (poor fit) 85% (0 outliers) 11% (1 outlier) .. (2 outliers)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	505	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FlsO1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	484	3648	2298	662	675	13	0	1	0
1	A	480	3594	2266	649	666	13	0	0	0
1	B	491	3689	2317	669	690	13	0	1	0
1	C	471	3514	2213	638	650	13	0	1	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	108	ALA	GLN	engineered mutation	UNP A0A0P0I576
D	109	ALA	ARG	engineered mutation	UNP A0A0P0I576
A	108	ALA	GLN	engineered mutation	UNP A0A0P0I576
A	109	ALA	ARG	engineered mutation	UNP A0A0P0I576
B	108	ALA	GLN	engineered mutation	UNP A0A0P0I576
B	109	ALA	ARG	engineered mutation	UNP A0A0P0I576
C	108	ALA	GLN	engineered mutation	UNP A0A0P0I576
C	109	ALA	ARG	engineered mutation	UNP A0A0P0I576

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	301	Total	O	0	0
			301	301		
5	A	244	Total	O	0	0
			244	244		
5	B	297	Total	O	0	0
			297	297		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	C	173	Total 173	O 173	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FlsO1




- Molecule 1: FlsO1





- Molecule 1: FlsO1







● Molecule 1: FlsO1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.54Å 84.64Å 102.80Å 91.31° 107.86° 115.39°	Depositor
Resolution (Å)	13.18 – 2.30 13.18 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.5 (13.18-2.30) 99.0 (13.18-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0267, PHENIX 1.19.2-4158	Depositor
R, $R_{free}$	0.211 , 0.271 0.222 , 0.278	Depositor DCC
$R_{free}$ test set	4704 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.4	Xtrriage
Anisotropy	0.083	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	15743	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FAD, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.71	0/3667	0.85	0/4977
1	B	0.71	0/3766	0.87	3/5113 (0.1%)
1	C	0.69	0/3583	0.82	2/4860 (0.0%)
1	D	0.71	0/3725	0.87	4/5055 (0.1%)
All	All	0.70	0/14741	0.85	9/20005 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	393	ARG	NE-CZ-NH2	-7.33	116.64	120.30
1	B	331	ARG	NE-CZ-NH1	6.48	123.54	120.30
1	C	393	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	B	331	ARG	NE-CZ-NH2	-6.19	117.20	120.30
1	D	65	ARG	NE-CZ-NH2	-5.84	117.38	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3594	0	3522	54	0
1	B	3689	0	3605	59	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3514	0	3433	64	0
1	D	3648	0	3580	42	0
2	A	20	0	0	3	0
2	B	35	0	0	3	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
3	A	53	0	31	1	0
3	B	53	0	31	2	0
3	C	53	0	31	2	0
3	D	53	0	31	2	0
4	B	1	0	0	0	0
5	A	244	0	0	8	0
5	B	297	0	0	5	0
5	C	173	0	0	11	0
5	D	301	0	0	10	0
All	All	15743	0	14264	220	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 220 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:ARG:C	1:A:226:SER:N	2.14	1.00
1:C:317:LEU:O	1:C:320:THR:HG22	1.70	0.91
1:C:335:ASN:HB2	1:C:372:MET:HE2	1.56	0.86
1:B:440:ASP:OD2	1:B:497:ARG:HD2	1.82	0.80
1:B:62:LEU:HD22	1:B:105:MET:HE1	1.64	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/497 (94%)	453 (97%)	14 (3%)	1 (0%)	47	58
1	B	484/497 (97%)	470 (97%)	13 (3%)	1 (0%)	47	58
1	C	460/497 (93%)	426 (93%)	31 (7%)	3 (1%)	22	26
1	D	477/497 (96%)	457 (96%)	18 (4%)	2 (0%)	34	42
All	All	1889/1988 (95%)	1806 (96%)	76 (4%)	7 (0%)	34	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	159	SER
1	D	459	HIS
1	C	144	GLN
1	C	447	ALA
1	D	402	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/377 (95%)	344 (96%)	13 (4%)	35	49
1	B	368/377 (98%)	360 (98%)	8 (2%)	52	69
1	C	345/377 (92%)	337 (98%)	8 (2%)	50	67
1	D	364/377 (97%)	354 (97%)	10 (3%)	44	61
All	All	1434/1508 (95%)	1395 (97%)	39 (3%)	44	61

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	340	ARG
1	C	340	ARG
1	B	393	ARG
1	C	169	PHE
1	C	409	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	397	GLN
1	B	397	GLN
1	B	416	HIS
1	B	450	HIS
1	C	283	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	B	506	-	4,4,4	1.05	0	6,6,6	0.37	0
2	PO4	A	504	-	4,4,4	0.59	0	6,6,6	0.45	0
3	FAD	B	509	-	53,58,58	0.67	0	68,89,89	0.82	1 (1%)
2	PO4	B	507	-	4,4,4	0.78	0	6,6,6	0.55	0
2	PO4	C	501	-	4,4,4	0.70	0	6,6,6	0.37	0
3	FAD	D	503	-	53,58,58	0.70	0	68,89,89	0.80	0
3	FAD	A	505	-	53,58,58	0.64	0	68,89,89	0.91	6 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	B	502	-	4,4,4	0.62	0	6,6,6	0.53	0
2	PO4	B	505	-	4,4,4	0.51	0	6,6,6	0.67	0
2	PO4	D	502	-	4,4,4	0.98	0	6,6,6	0.49	0
2	PO4	B	503	-	4,4,4	1.51	1 (25%)	6,6,6	0.58	0
2	PO4	D	501	-	4,4,4	0.64	0	6,6,6	0.44	0
2	PO4	B	501	-	4,4,4	0.74	0	6,6,6	0.52	0
2	PO4	B	504	-	4,4,4	0.57	0	6,6,6	0.51	0
2	PO4	A	501	-	4,4,4	0.85	0	6,6,6	0.50	0
2	PO4	A	503	-	4,4,4	0.87	0	6,6,6	0.30	0
3	FAD	C	502	-	53,58,58	0.67	0	68,89,89	0.76	2 (2%)
2	PO4	A	502	-	4,4,4	0.85	0	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	C	502	-	-	1/30/50/50	0/6/6/6
3	FAD	B	509	-	-	2/30/50/50	0/6/6/6
3	FAD	A	505	-	-	1/30/50/50	0/6/6/6
3	FAD	D	503	-	-	1/30/50/50	0/6/6/6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	503	PO4	P-O1	2.69	1.57	1.50

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	505	FAD	O2A-PA-O1A	2.33	123.76	112.24
3	A	505	FAD	C4'-C3'-C2'	2.28	118.10	113.36
3	B	509	FAD	O2P-P-O1P	2.25	123.37	112.24
3	C	502	FAD	C5A-C6A-N6A	2.24	123.76	120.35
3	C	502	FAD	P-O3P-PA	-2.24	125.15	132.83

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	509	FAD	PA-O3P-P-O5'
3	A	505	FAD	O4B-C4B-C5B-O5B
3	D	503	FAD	O4B-C4B-C5B-O5B
3	C	502	FAD	O4B-C4B-C5B-O5B
3	B	509	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.


9 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	504	PO4	1	0
3	B	509	FAD	2	0
3	D	503	FAD	2	0
3	A	505	FAD	1	0
2	B	505	PO4	2	0
2	B	504	PO4	1	0
2	A	501	PO4	1	0
2	A	503	PO4	1	0
3	C	502	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	480/497 (96%)	-0.06	21 (4%) 34 41	12, 29, 62, 79	0
1	B	491/497 (98%)	-0.21	13 (2%) 56 63	8, 23, 49, 84	0
1	C	471/497 (94%)	0.34	20 (4%) 36 43	16, 40, 64, 76	0
1	D	484/497 (97%)	-0.22	8 (1%) 70 76	10, 24, 47, 78	0
All	All	1926/1988 (96%)	-0.04	62 (3%) 47 54	8, 29, 58, 84	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	454	PRO	5.6
1	B	451	THR	4.7
1	A	399	LEU	4.3
1	D	141	ASP	4.1
1	D	140	ALA	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	B	502	5/5	0.80	0.32	61,68,76,80	0
2	PO4	D	501	5/5	0.84	0.30	45,54,64,71	0
2	PO4	B	507	5/5	0.85	0.26	41,45,58,58	0
2	PO4	B	505	5/5	0.86	0.22	42,43,51,54	0
2	PO4	A	501	5/5	0.86	0.21	61,63,72,80	0
2	PO4	B	506	5/5	0.88	0.21	70,71,75,82	0
2	PO4	C	501	5/5	0.89	0.27	59,63,67,69	0
3	FAD	C	502	53/53	0.91	0.16	24,37,69,70	0
2	PO4	B	503	5/5	0.92	0.14	21,27,28,30	0
2	PO4	A	504	5/5	0.92	0.20	58,59,70,71	0
2	PO4	D	502	5/5	0.92	0.19	43,45,50,61	0
2	PO4	A	502	5/5	0.94	0.23	31,34,37,41	0
2	PO4	A	503	5/5	0.94	0.18	29,33,38,39	0
2	PO4	B	501	5/5	0.95	0.28	43,44,53,59	0
3	FAD	D	503	53/53	0.96	0.10	13,19,36,37	0
3	FAD	A	505	53/53	0.96	0.09	12,17,27,33	0
2	PO4	B	504	5/5	0.96	0.17	42,42,44,55	0
3	FAD	B	509	53/53	0.97	0.08	11,14,26,28	0
4	NA	B	508	1/1	0.97	0.09	30,30,30,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FAD C 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD D 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD A 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD B 509:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.