



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 11, 2024 – 09:54 PM EDT

PDB ID : 1W0M
Title : Triosephosphate isomerase from *Thermoproteus tenax*
Authors : Walden, H.; Taylor, G.; Lorentzen, E.; Pohl, E.; Lilie, H.; Schramm, A.; Knura, T.; Stubbe, K.; Tjaden, B.; Hensel, R.
Deposited on : 2004-06-08
Resolution : 2.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	226	82% 15% .
1	B	226	76% 20% .
1	C	226	80% 17% .
1	D	226	77% 20% .
1	E	226	78% 18% .
1	F	226	81% 15% .
1	G	226	72% 23% . .
1	H	226	77% 19% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 13752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIOSEPHOSPHATE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	226	1629	1030	291	306	2	0	0	1
1	B	226	1629	1030	291	306	2	0	0	1
1	C	226	1629	1030	291	306	2	0	0	1
1	D	226	1629	1030	291	306	2	0	0	1
1	E	226	1629	1030	291	306	2	0	0	1
1	F	225	1621	1024	290	305	2	0	0	1
1	G	224	1612	1019	289	302	2	0	0	1
1	H	226	1629	1030	291	306	2	0	0	1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0
2	E	1	Total O P 5 4 1	0	0
2	F	1	Total O P 5 4 1	0	0
2	G	1	Total O P 5 4 1	0	0
2	H	1	Total O P 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	102	Total O 102 102	0	0
3	B	97	Total O 97 97	0	0
3	C	139	Total O 139 139	0	0
3	D	110	Total O 110 110	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	89	Total 89	O 89	0	0
3	F	62	Total 62	O 62	0	0
3	G	47	Total 47	O 47	0	0
3	H	59	Total 59	O 59	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: TRIOSEPHOSPHATE ISOMERASE




- Molecule 1: TRIOSEPHOSPHATE ISOMERASE



- Molecule 1: TRIOSEPHOSPHATE ISOMERASE




- Molecule 1: TRIOSEPHOSPHATE ISOMERASE






- Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain E: 78% 18%




- Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain F: 81% 15%




- Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain G: 72% 23%



- Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain H: 77% 19%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	186.89Å 186.89Å 287.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.50	Depositor
% Data completeness (in resolution range)	96.5 (30.00-2.50)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.197 , 0.226	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	13752	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1653	0.73	0/2250
1	B	0.56	0/1653	0.74	0/2250
1	C	0.62	0/1653	0.75	0/2250
1	D	0.59	0/1653	0.73	0/2250
1	E	0.52	0/1653	0.78	4/2250 (0.2%)
1	F	0.50	0/1645	0.69	0/2239
1	G	0.42	0/1636	0.66	0/2227
1	H	0.46	0/1653	0.68	0/2250
All	All	0.53	0/13199	0.72	4/17966 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	3	LEU	C-N-CD	9.72	148.82	128.40
1	E	3	LEU	N-CA-C	6.05	127.33	111.00
1	E	3	LEU	CA-CB-CG	-5.88	101.78	115.30
1	E	3	LEU	C-N-CA	-5.58	98.57	122.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1702	48	0
1	B	1629	0	1702	51	0
1	C	1629	0	1702	46	0
1	D	1629	0	1702	46	0
1	E	1629	0	1702	43	0
1	F	1621	0	1691	38	0
1	G	1612	0	1685	58	0
1	H	1629	0	1702	45	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	5	0	0	0	0
2	H	5	0	0	0	0
3	A	102	0	0	2	0
3	B	97	0	0	6	0
3	C	139	0	0	6	0
3	D	110	0	0	5	0
3	E	89	0	0	2	0
3	F	62	0	0	2	0
3	G	47	0	0	4	0
3	H	59	0	0	5	0
All	All	13752	0	13588	352	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 352 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LEU:HB3	1:A:4:PRO:HD3	1.29	1.14
1:A:3:LEU:CB	1:A:4:PRO:HD3	1.83	1.07
1:B:3:LEU:CB	1:B:4:PRO:HD3	1.92	0.99
1:H:93:HIS:HD2	1:H:95:GLU:H	1.13	0.96
1:H:117:VAL:H	1:H:136:HIS:HD2	1.14	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	215 (96%)	7 (3%)	2 (1%)	17	31
1	B	224/226 (99%)	213 (95%)	9 (4%)	2 (1%)	17	31
1	C	224/226 (99%)	216 (96%)	6 (3%)	2 (1%)	17	31
1	D	224/226 (99%)	215 (96%)	7 (3%)	2 (1%)	17	31
1	E	224/226 (99%)	214 (96%)	8 (4%)	2 (1%)	17	31
1	F	223/226 (99%)	209 (94%)	12 (5%)	2 (1%)	17	31
1	G	222/226 (98%)	206 (93%)	13 (6%)	3 (1%)	11	20
1	H	224/226 (99%)	214 (96%)	8 (4%)	2 (1%)	17	31
All	All	1789/1808 (99%)	1702 (95%)	70 (4%)	17 (1%)	15	28

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	B	3	LEU
1	C	3	LEU
1	D	3	LEU
1	E	225	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/167 (99%)	157 (95%)	9 (5%)	22	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	166/167 (99%)	157 (95%)	9 (5%)	22	42
1	C	166/167 (99%)	156 (94%)	10 (6%)	19	37
1	D	166/167 (99%)	156 (94%)	10 (6%)	19	37
1	E	166/167 (99%)	159 (96%)	7 (4%)	30	54
1	F	165/167 (99%)	155 (94%)	10 (6%)	18	36
1	G	164/167 (98%)	156 (95%)	8 (5%)	25	47
1	H	166/167 (99%)	155 (93%)	11 (7%)	16	32
All	All	1325/1336 (99%)	1251 (94%)	74 (6%)	21	40

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	125	ARG
1	H	173	GLU
1	G	150	ARG
1	H	48	LEU
1	C	173	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	136	HIS
1	H	45	HIS
1	D	45	HIS
1	C	136	HIS
1	H	93	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	C	1226	-	4,4,4	1.45	0	6,6,6	0.47	0
2	PO4	A	1226	-	4,4,4	1.40	0	6,6,6	0.44	0
2	PO4	B	1226	-	4,4,4	1.48	0	6,6,6	0.46	0
2	PO4	E	1226	-	4,4,4	1.36	0	6,6,6	0.49	0
2	PO4	F	1225	-	4,4,4	1.35	0	6,6,6	0.46	0
2	PO4	D	1226	-	4,4,4	1.39	0	6,6,6	0.47	0
2	PO4	G	1224	-	4,4,4	1.60	0	6,6,6	0.49	0
2	PO4	H	1226	-	4,4,4	1.63	0	6,6,6	0.41	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.