



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 30, 2023 – 03:33 PM JST

PDB ID : 8W85
Title : HLA-DQ2.5-gamma2 gliadin peptide in complex with DQN0385AE01
Authors : Irie, M.; Tsushima, T.; Teranishi-Ikawa, Y.; Takahashi, N.; Ishii, S.; Okura, Y.; Fukami, T.A.; Torizawa, T.
Deposited on : 2023-08-31
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
1	E	223	
2	B	213	
2	F	213	
3	C	189	
3	G	189	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	222	 <p>2% 69% 14% 16%</p>
4	H	222	 <p>7% 67% 14% 20%</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DQN0385AE01 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	211	1559	988	261	303	7	0	0	0
1	E	126	875	555	141	174	5	0	0	0

- Molecule 2 is a protein called DQN0385AE01 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	211	1548	973	249	321	5	0	0	0
2	F	105	736	465	120	148	3	0	0	0

- Molecule 3 is a protein called HLA class II histocompatibility antigen, DQ alpha 1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	180	1383	896	225	260	2	0	0	0
3	G	179	1379	889	224	264	2	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	47	SER	CYS	engineered mutation	UNP P01909
C	184	LEU	-	expression tag	UNP P01909
C	185	GLU	-	expression tag	UNP P01909
C	186	VAL	-	expression tag	UNP P01909
C	187	LEU	-	expression tag	UNP P01909
C	188	PHE	-	expression tag	UNP P01909
C	189	GLN	-	expression tag	UNP P01909
G	47	SER	CYS	engineered mutation	UNP P01909

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	184	LEU	-	expression tag	UNP P01909
G	185	GLU	-	expression tag	UNP P01909
G	186	VAL	-	expression tag	UNP P01909
G	187	LEU	-	expression tag	UNP P01909
G	188	PHE	-	expression tag	UNP P01909
G	189	GLN	-	expression tag	UNP P01909

- Molecule 4 is a protein called MHC class II HLA-DQ-beta-1 - gamma2 gliadin peptide chimeric protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	186	1427	909	247	264	7	0	0	0
4	H	178	1360	869	231	253	7	0	0	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	986	GLY	-	linker	PDB ?
D	987	GLY	-	linker	PDB ?
D	988	GLY	-	linker	PDB ?
D	989	GLY	-	linker	PDB ?
D	990	SER	-	linker	PDB ?
D	991	ILE	-	linker	PDB ?
D	992	GLU	-	linker	PDB ?
D	993	GLY	-	linker	PDB ?
D	994	ARG	-	linker	PDB ?
D	995	GLY	-	linker	PDB ?
D	996	SER	-	linker	PDB ?
D	997	GLY	-	linker	PDB ?
D	998	GLY	-	linker	PDB ?
D	999	GLY	-	linker	PDB ?
D	1000	SER	-	linker	PDB ?
D	1191	LEU	-	expression tag	UNP O19712
D	1192	GLU	-	expression tag	UNP O19712
D	1193	VAL	-	expression tag	UNP O19712
D	1194	LEU	-	expression tag	UNP O19712
D	1195	PHE	-	expression tag	UNP O19712
D	1196	GLN	-	expression tag	UNP O19712
H	986	GLY	-	linker	PDB ?
H	987	GLY	-	linker	PDB ?

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	988	GLY	-	linker	PDB ?
H	989	GLY	-	linker	PDB ?
H	990	SER	-	linker	PDB ?
H	991	ILE	-	linker	PDB ?
H	992	GLU	-	linker	PDB ?
H	993	GLY	-	linker	PDB ?
H	994	ARG	-	linker	PDB ?
H	995	GLY	-	linker	PDB ?
H	996	SER	-	linker	PDB ?
H	997	GLY	-	linker	PDB ?
H	998	GLY	-	linker	PDB ?
H	999	GLY	-	linker	PDB ?
H	1000	SER	-	linker	PDB ?
H	1191	LEU	-	expression tag	UNP O19712
H	1192	GLU	-	expression tag	UNP O19712
H	1193	VAL	-	expression tag	UNP O19712
H	1194	LEU	-	expression tag	UNP O19712
H	1195	PHE	-	expression tag	UNP O19712
H	1196	GLN	-	expression tag	UNP O19712

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	C	1	14	8	1	5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	G	1	14	8	1	5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DQN0385AE01 Fab heavy chain




- Molecule 1: DQN0385AE01 Fab heavy chain




- Molecule 2: DQN0385AE01 Fab light chain




- Molecule 2: DQN0385AE01 Fab light chain




- Molecule 3: HLA class II histocompatibility antigen, DQ alpha 1 chain




- Molecule 3: HLA class II histocompatibility antigen, DQ alpha 1 chain




- Molecule 4: MHC class II HLA-DQ-beta-1 - gamma2 gliadin peptide chimeric protein



- Molecule 4: MHC class II HLA-DQ-beta-1 - gamma2 gliadin peptide chimeric protein





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	150.53Å 243.91Å 136.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.47 – 2.77 50.47 – 2.77	Depositor EDS
% Data completeness (in resolution range)	65.3 (50.47-2.77) 65.3 (50.47-2.77)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.77Å)	Xtrriage
Refinement program	BUSTER 2.11.8	Depositor
R, R_{free}	0.253 , 0.300 0.251 , 0.303	Depositor DCC
R_{free} test set	2093 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtrriage
Anisotropy	0.043	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10295	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1601	0.67	0/2197
1	E	0.36	0/898	0.57	0/1237
2	B	0.45	0/1584	0.71	1/2172 (0.0%)
2	F	0.40	0/755	0.60	0/1036
3	C	0.44	0/1425	0.65	0/1956
3	G	0.37	0/1421	0.60	0/1951
4	D	0.48	0/1459	0.65	0/1996
4	H	0.40	0/1389	0.59	0/1900
All	All	0.43	0/10532	0.64	1/14445 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	50	TYR	C-N-CA	5.79	136.19	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1559	0	1462	30	0
1	E	875	0	750	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1548	0	1419	24	0
2	F	736	0	639	12	0
3	C	1383	0	1299	23	0
3	G	1379	0	1280	26	0
4	D	1427	0	1344	26	0
4	H	1360	0	1270	18	0
5	C	14	0	13	1	0
5	G	14	0	13	0	0
All	All	10295	0	9489	148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:17:SER:HB3	1:E:85:ASN:HA	1.52	0.89
1:A:92:THR:HG23	1:A:115:THR:HA	1.62	0.80
3:G:113:ASN:H	3:G:142:LYS:NZ	1.80	0.78
1:A:6:GLU:HB3	1:A:112:THR:HB	1.67	0.77
2:B:53:THR:HG23	2:B:53:THR:O	1.87	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/223 (93%)	187 (90%)	17 (8%)	3 (1%)	11 19
1	E	122/223 (55%)	106 (87%)	16 (13%)	0	100 100
2	B	209/213 (98%)	185 (88%)	20 (10%)	4 (2%)	8 14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	103/213 (48%)	83 (81%)	17 (16%)	3 (3%)	4	6
3	C	178/189 (94%)	166 (93%)	12 (7%)	0	100	100
3	G	177/189 (94%)	164 (93%)	9 (5%)	4 (2%)	6	10
4	D	178/222 (80%)	169 (95%)	9 (5%)	0	100	100
4	H	168/222 (76%)	153 (91%)	14 (8%)	1 (1%)	25	42
All	All	1342/1694 (79%)	1213 (90%)	114 (8%)	15 (1%)	14	25

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	32	TRP
2	B	51	ALA
2	B	100	GLY
2	B	137	ASN
1	A	149	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/188 (89%)	152 (90%)	16 (10%)	8	15
1	E	82/188 (44%)	75 (92%)	7 (8%)	10	19
2	B	168/187 (90%)	158 (94%)	10 (6%)	19	33
2	F	71/187 (38%)	64 (90%)	7 (10%)	8	13
3	C	151/173 (87%)	145 (96%)	6 (4%)	31	51
3	G	151/173 (87%)	140 (93%)	11 (7%)	14	25
4	D	150/196 (76%)	145 (97%)	5 (3%)	38	58
4	H	142/196 (72%)	130 (92%)	12 (8%)	10	19
All	All	1083/1488 (73%)	1009 (93%)	74 (7%)	16	28

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	124	LEU
4	H	1139	THR
3	G	155	LEU
4	H	1050	VAL
2	B	165	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	179	HIS
4	H	8	GLN
4	D	1019	ASN
4	D	1174	HIS
3	G	7	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	C	1000	-	14,14,15	0.33	0	17,19,21	1.26	1 (5%)
5	NAG	G	1000	-	14,14,15	0.26	0	17,19,21	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	1000	-	-	2/6/23/26	0/1/1/1
5	NAG	G	1000	-	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1000	NAG	O5-C1-C2	-4.80	103.70	111.29

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	1000	NAG	C4-C5-C6-O6
5	C	1000	NAG	O5-C5-C6-O6
5	G	1000	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1000	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/223 (94%)	0.00	2 (0%) 84 89	44, 69, 91, 104	0
1	E	126/223 (56%)	0.35	10 (7%) 12 15	94, 115, 133, 144	0
2	B	211/213 (99%)	0.11	4 (1%) 66 75	52, 79, 102, 107	0
2	F	105/213 (49%)	1.01	19 (18%) 1 1	100, 121, 137, 141	0
3	C	180/189 (95%)	0.02	1 (0%) 89 92	46, 66, 88, 99	0
3	G	179/189 (94%)	0.22	4 (2%) 62 70	72, 93, 122, 139	0
4	D	186/222 (83%)	0.03	5 (2%) 54 63	50, 71, 94, 111	0
4	H	178/222 (80%)	0.55	15 (8%) 11 13	74, 95, 108, 116	0
All	All	1376/1694 (81%)	0.23	60 (4%) 34 41	44, 84, 124, 144	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	75	ILE	6.9
2	F	62	PHE	6.8
2	F	19	VAL	5.4
4	H	1163	MET	5.3
2	F	82	ASP	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	G	1000	14/15	0.76	0.17	170,170,171,171	0
5	NAG	C	1000	14/15	0.81	0.23	108,109,109,110	0

6.5 Other polymers [i](#)

There are no such residues in this entry.