



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 18, 2024 – 04:11 PM JST

PDB ID : 8WEV  
Title : Crystal structure of Feruoyl-CoA Synthetase complexed with AMP from Amycolatopsis thermoflava  
Authors : Seok, J.; Kim, K.-J.  
Deposited on : 2023-09-18  
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	491	 3% 87% 10% ..
1	B	491	 5% 81% 14% ..

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Feruloyl-CoA Synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	486	3632	2297	652	668	15	0	0	0
1	B	476	3566	2256	638	657	15	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ARG	GLN	conflict	UNP A0A3N2H3K2
A	48	GLU	ALA	conflict	UNP A0A3N2H3K2
A	224	PRO	ALA	conflict	UNP A0A3N2H3K2
A	239	VAL	MET	conflict	UNP A0A3N2H3K2
A	256	ARG	HIS	conflict	UNP A0A3N2H3K2
A	258	THR	ALA	conflict	UNP A0A3N2H3K2
A	448	ALA	GLY	conflict	UNP A0A3N2H3K2
B	47	ARG	GLN	conflict	UNP A0A3N2H3K2
B	48	GLU	ALA	conflict	UNP A0A3N2H3K2
B	224	PRO	ALA	conflict	UNP A0A3N2H3K2
B	239	VAL	MET	conflict	UNP A0A3N2H3K2
B	256	ARG	HIS	conflict	UNP A0A3N2H3K2
B	258	THR	ALA	conflict	UNP A0A3N2H3K2
B	448	ALA	GLY	conflict	UNP A0A3N2H3K2

- Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Mg	0	0
			3	3		
3	B	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		
4	B	38	Total	O	0	0
			38	38		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Feruloyl-CoA Synthetase



- Molecule 1: Feruloyl-CoA Synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.76Å 132.10Å 162.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.49 – 2.29 31.49 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.4 (31.49-2.29) 98.4 (31.49-2.29)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.85 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.199 , 0.261 0.203 , 0.260	Depositor DCC
$R_{free}$ test set	2680 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.9	Xtrriage
Anisotropy	0.402	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 34.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7342	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/3716	0.87	1/5065 (0.0%)
1	B	0.69	0/3646	0.85	0/4966
All	All	0.69	0/7362	0.86	1/10031 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	299	PRO	N-CA-CB	-5.11	96.98	102.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	450	LEU	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3632	0	3627	24	0
1	B	3566	0	3542	40	0
2	A	23	0	12	0	0
2	B	23	0	12	0	0
3	A	3	0	0	0	0
3	B	1	0	0	0	0
4	A	56	0	0	0	0
4	B	38	0	0	1	0
All	All	7342	0	7193	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:154:THR:O	1:B:154:THR:OG1	2.03	0.73
1:B:327:VAL:HG23	1:B:335:VAL:HG22	1.73	0.69
1:B:24:ARG:HE	1:B:189:ARG:NH1	1.89	0.69
1:B:483:LYS:HB3	1:B:485:GLU:HG2	1.75	0.68
1:B:71:LEU:C	1:B:71:LEU:HD23	2.16	0.66
1:B:298:ALA:N	1:B:299:PRO:HA	2.10	0.66
1:A:453:TRP:O	1:A:456:THR:HB	1.97	0.65
1:A:298:ALA:N	1:A:299:PRO:HA	2.13	0.64
1:B:24:ARG:HE	1:B:189:ARG:HH12	1.45	0.63
1:B:126:VAL:HG11	1:B:131:LEU:HD22	1.84	0.60
1:A:60:LEU:O	1:A:107:HIS:HA	2.02	0.59
1:B:473:LEU:HD23	1:B:482:LEU:HD11	1.88	0.55
1:B:475:THR:HG22	1:B:482:LEU:HG	1.90	0.54
1:B:205:VAL:HG21	1:B:240:PHE:HB2	1.89	0.54
1:B:71:LEU:HD23	1:B:71:LEU:O	2.08	0.54
1:A:485:GLU:HA	1:A:488:ARG:HG3	1.90	0.54
1:A:395:ILE:O	1:A:431:GLU:HB2	2.08	0.53
1:B:150:LEU:C	1:B:150:LEU:HD12	2.29	0.53
1:B:407:GLU:HG3	1:B:421:VAL:HG23	1.90	0.53
1:B:24:ARG:NE	1:B:189:ARG:HH12	2.05	0.53
1:A:200:ALA:HB1	1:A:240:PHE:CZ	2.45	0.51
1:B:71:LEU:C	1:B:71:LEU:CD2	2.79	0.51
1:B:34:GLU:O	1:B:38:ARG:HG3	2.10	0.50
1:A:483:LYS:C	1:A:485:GLU:H	2.16	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:VAL:HG21	1:A:478:SER:HB3	1.93	0.49
1:B:200:ALA:HB1	1:B:240:PHE:CZ	2.48	0.49
1:B:290:GLN:HE21	1:B:309:VAL:HG23	1.78	0.48
1:A:297:ALA:C	1:A:299:PRO:HA	2.33	0.48
1:A:327:VAL:HG23	1:A:335:VAL:HG22	1.96	0.48
1:B:438:VAL:HG12	1:B:470:THR:HG23	1.97	0.47
1:B:265:THR:HG22	1:B:267:LEU:HD12	1.97	0.47
1:B:297:ALA:C	1:B:299:PRO:HA	2.35	0.47
1:A:90:VAL:N	1:A:91:PRO:CD	2.78	0.47
1:A:358:PRO:HG3	1:B:141:LEU:HD12	1.96	0.46
1:A:267:LEU:HD23	1:A:289:VAL:HB	1.97	0.46
1:A:38:ARG:NH2	1:A:132:GLU:OE2	2.48	0.46
1:B:250:ALA:HA	1:B:255:TRP:CD2	2.51	0.45
1:B:57:VAL:HB	1:B:74:CYS:SG	2.57	0.45
1:B:152:MET:HE1	1:B:295:THR:HG23	1.98	0.44
1:A:241:GLY:O	1:A:268:CYS:HA	2.16	0.44
1:B:107:HIS:CG	1:B:115:VAL:HG21	2.53	0.44
1:B:49:ALA:HB2	1:B:131:LEU:HD11	2.00	0.44
1:A:436:VAL:HG21	1:A:489:ARG:HD3	1.99	0.44
1:B:189:ARG:NH2	1:B:234:GLU:O	2.45	0.44
1:B:273:VAL:HG22	1:B:290:GLN:OE1	2.18	0.44
1:A:375:VAL:HG22	1:A:390:LEU:HD13	2.00	0.43
1:B:25:HIS:HE1	1:B:220:SER:OG	2.02	0.43
1:A:482:LEU:HD13	1:A:482:LEU:HA	1.90	0.42
1:A:456:THR:HG22	1:A:457:ARG:HE	1.85	0.42
1:A:41:ARG:HD2	1:A:133:ALA:O	2.20	0.42
1:B:447:GLY:O	1:B:448:ALA:HB3	2.19	0.41
1:B:41:ARG:HG2	1:B:135:ASP:O	2.20	0.41
1:A:247:GLN:HG3	1:A:277:LEU:HD13	2.03	0.41
1:B:45:GLY:HA2	1:B:133:ALA:HB2	2.01	0.41
1:A:35:LEU:O	1:A:39:VAL:HG23	2.21	0.41
1:A:90:VAL:HG22	1:A:91:PRO:HD3	2.03	0.41
1:A:378:VAL:HG22	1:A:384:PHE:CD2	2.56	0.41
1:B:207:LEU:N	1:B:208:PRO:CD	2.84	0.41
1:B:433:GLY:HA3	4:B:615:HOH:O	2.21	0.41
1:B:292:TYR:HB3	1:B:302:LEU:HB2	2.04	0.40
1:B:344:VAL:HA	1:B:370:PHE:O	2.20	0.40
1:B:477:GLY:H	1:B:483:LYS:HZ1	1.68	0.40
1:B:318:VAL:HB	1:B:319:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	482/491 (98%)	464 (96%)	16 (3%)	2 (0%)	30	39
1	B	464/491 (94%)	436 (94%)	26 (6%)	2 (0%)	30	39
All	All	946/982 (96%)	900 (95%)	42 (4%)	4 (0%)	30	39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	298	ALA
1	B	298	ALA
1	B	474	PRO
1	A	479	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/374 (99%)	349 (94%)	22 (6%)	16	23
1	B	365/374 (98%)	345 (94%)	20 (6%)	18	26
All	All	736/748 (98%)	694 (94%)	42 (6%)	17	25

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	VAL
1	A	13	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	56	ARG
1	A	71	LEU
1	A	90	VAL
1	A	130	GLU
1	A	182	THR
1	A	273	VAL
1	A	282	LEU
1	A	299	PRO
1	A	331	SER
1	A	388	ASP
1	A	391	LYS
1	A	424	VAL
1	A	432	VAL
1	A	444	ARG
1	A	470	THR
1	A	471	ASP
1	A	478	SER
1	A	483	LYS
1	A	488	ARG
1	A	489	ARG
1	B	11	VAL
1	B	13	ARG
1	B	47	ARG
1	B	56	ARG
1	B	212	LYS
1	B	253	PRO
1	B	257	SER
1	B	276	ASP
1	B	299	PRO
1	B	364	VAL
1	B	432	VAL
1	B	450	LEU
1	B	456	THR
1	B	471	ASP
1	B	472	ARG
1	B	473	LEU
1	B	482	LEU
1	B	483	LYS
1	B	488	ARG
1	B	489	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	HIS
1	A	76	GLN
1	A	198	HIS
1	A	247	GLN
1	A	252	HIS
1	B	25	HIS
1	B	76	GLN
1	B	252	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	AMP	B	501	-	22,25,25	0.66	0	25,38,38	0.71	1 (4%)
2	AMP	A	501	-	22,25,25	0.63	0	25,38,38	0.76	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	B	501	-	-	3/6/26/26	0/3/3/3
2	AMP	A	501	-	-	0/6/26/26	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	AMP	C5-C6-N6	2.39	123.99	120.35
2	B	501	AMP	C5-C6-N6	2.07	123.50	120.35

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	AMP	C3'-C4'-C5'-O5'
2	B	501	AMP	O4'-C4'-C5'-O5'
2	B	501	AMP	C5'-O5'-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	486/491 (98%)	-0.01	17 (3%) 47 49	29, 48, 89, 127	0
1	B	476/491 (96%)	0.36	26 (5%) 32 33	35, 56, 98, 130	0
All	All	962/982 (97%)	0.18	43 (4%) 39 40	29, 52, 94, 130	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	490	PHE	5.3
1	B	450	LEU	5.2
1	B	313	ILE	4.6
1	B	484	GLY	4.3
1	A	284	ARG	4.1
1	A	154	THR	4.0
1	A	310	ALA	3.5
1	A	334	PRO	3.4
1	B	448	ALA	3.3
1	B	482	LEU	3.3
1	B	473	LEU	3.2
1	B	488	ARG	3.2
1	A	309	VAL	3.1
1	B	427	PRO	3.0
1	B	119	ALA	2.9
1	B	158	THR	2.8
1	A	484	GLY	2.8
1	B	306	ARG	2.8
1	A	441	ALA	2.8
1	B	155	SER	2.7
1	A	313	ILE	2.7
1	B	476	THR	2.6
1	B	331	SER	2.6
1	A	424	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	470	THR	2.5
1	B	131	LEU	2.5
1	A	489	ARG	2.4
1	B	414	PRO	2.4
1	B	449	GLU	2.4
1	A	485	GLU	2.4
1	B	477	GLY	2.4
1	A	120	GLY	2.3
1	A	442	GLY	2.3
1	B	116	ALA	2.2
1	B	475	THR	2.2
1	B	478	SER	2.2
1	B	453	TRP	2.2
1	A	438	VAL	2.2
1	B	309	VAL	2.2
1	A	444	ARG	2.1
1	B	485	GLU	2.0
1	A	337	PRO	2.0
1	B	439	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	AMP	B	501	23/23	0.94	0.07	50,62,66,67	0
3	MG	A	504	1/1	0.94	0.33	60,60,60,60	0
3	MG	A	502	1/1	0.95	0.39	36,36,36,36	0
3	MG	A	503	1/1	0.96	0.36	44,44,44,44	0


*Continued on next page...*



Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	AMP	A	501	23/23	0.96	0.06	43,48,52,55	0
3	MG	B	502	1/1	0.97	0.40	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around MG A 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around MG A 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









## 6.5 Other polymers [i](#)

There are no such residues in this entry.