



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2022 – 02:09 PM JST

PDB ID : 7WG1
Title : DVAA-KlAtel
Authors : Kim, M.K.; Kim, B.H.; Oh, S.-J.; Song, H.K.
Deposited on : 2021-12-27
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.30
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.30


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	
1	B	507	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16145 atoms, of which 7803 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arginyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	489	7854	2541	3894	645	757	17	0	0	0
1	B	491	7882	2548	3909	648	760	17	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ASP	-	expression tag	UNP Q6CXX6
A	-2	VAL	-	expression tag	UNP Q6CXX6
A	-1	ALA	-	expression tag	UNP Q6CXX6
A	0	ALA	-	expression tag	UNP Q6CXX6
B	-3	ASP	-	expression tag	UNP Q6CXX6
B	-2	VAL	-	expression tag	UNP Q6CXX6
B	-1	ALA	-	expression tag	UNP Q6CXX6
B	0	ALA	-	expression tag	UNP Q6CXX6

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	235	Total	O	0	0
			235	235		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	172	Total 172	O 172	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Arginyltransferase



- Molecule 1: Arginyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.09Å 107.74Å 161.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.16 – 2.19 48.16 – 2.19	Depositor EDS
% Data completeness (in resolution range)	98.9 (48.16-2.19) 98.9 (48.16-2.19)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 2.18Å)	Xtrriage
Refinement program	PHENIX (1.19.2_4158: ???)	Depositor
R, R_{free}	0.178 , 0.234 0.177 , 0.233	Depositor DCC
R_{free} test set	2000 reflections (3.21%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtrriage
Anisotropy	0.589	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16145	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/4056	0.68	0/5492
1	B	0.59	0/4069	0.69	0/5509
All	All	0.60	0/8125	0.68	0/11001

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3960	3894	3894	30	1
1	B	3973	3909	3911	43	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	235	0	0	2	0
3	B	172	0	0	5	0
All	All	8342	7803	7805	72	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:GLN:NE2	1:A:403:SER:O	2.18	0.77
1:B:388:LYS:HB3	1:B:391:LYS:HZ1	1.61	0.65
1:A:318:LYS:HE2	1:A:357:LYS:HE3	1.81	0.63
1:B:173:TYR:CD2	1:B:259:ILE:HG22	2.35	0.62
1:A:232:GLU:OE1	3:A:702:HOH:O	2.16	0.62
1:A:110:MET:HG3	1:A:115:LYS:HG3	1.81	0.61
1:A:52:VAL:HG13	1:A:405:TYR:HB3	1.83	0.60
1:A:189:GLU:OE1	1:B:109:THR:HA	2.04	0.58
1:B:4:THR:O	3:B:701:HOH:O	2.17	0.58
1:B:180:ASP:OD1	1:B:180:ASP:N	2.34	0.58
1:B:498:LYS:HE3	3:B:725:HOH:O	2.04	0.57
1:B:45:TYR:CD2	1:B:52:VAL:HG21	2.40	0.56
1:B:66:VAL:O	1:B:66:VAL:HG13	2.05	0.56
1:B:45:TYR:HD2	1:B:52:VAL:HG21	1.71	0.56
1:A:99:TYR:HB3	1:A:311:ILE:HG23	1.89	0.55
1:B:112:LYS:O	1:B:116:THR:HG23	2.06	0.55
1:A:374:PRO:O	1:A:402:ARG:NH1	2.40	0.55
1:B:153:PHE:O	1:B:154:LYS:HD3	2.06	0.55
1:A:381:ASN:HB3	1:A:439:MET:HE3	1.89	0.55
1:A:48:ASP:OD1	1:A:51:LYS:CE	2.57	0.52
1:B:399:ASP:OD1	1:B:427:ARG:NH2	2.42	0.52
1:A:51:LYS:O	1:A:408:ILE:N	2.42	0.52
1:B:382:ALA:HB1	1:B:430:ASN:HB3	1.92	0.52
1:B:117:SER:HA	1:B:120:ARG:HH11	1.75	0.51
1:A:252:GLY:HA3	1:A:288:TYR:O	2.10	0.51
1:A:108:LEU:HD21	1:A:110:MET:HE2	1.93	0.51
1:B:44:PHE:HD2	1:B:45:TYR:CE1	2.29	0.51
1:A:327:GLN:OE1	1:A:330:LYS:HE2	2.11	0.50
1:A:296:ASP:OD2	1:A:326:HIS:CE1	2.64	0.50
1:B:388:LYS:CB	1:B:391:LYS:HZ1	2.23	0.50
1:B:469:ILE:HD12	3:B:826:HOH:O	2.10	0.50
1:B:438:LEU:O	1:B:442:VAL:HG23	2.12	0.50
1:A:110:MET:CG	1:A:115:LYS:HG3	2.41	0.50
1:B:326:HIS:CE1	1:B:330:LYS:HD3	2.47	0.49
1:A:108:LEU:HD21	1:A:110:MET:CE	2.44	0.48
1:B:57:SER:OG	1:B:90:ASP:HB3	2.13	0.48
1:A:5:ASP:OD2	1:A:498:LYS:NZ	2.46	0.48
1:B:388:LYS:HA	1:B:391:LYS:HZ3	1.79	0.48
1:B:44:PHE:CD2	1:B:45:TYR:CE1	3.02	0.47
1:A:381:ASN:HB3	1:A:439:MET:CE	2.44	0.47
1:A:51:LYS:O	1:A:407:GLU:HA	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464:ILE:O	1:A:464:ILE:HG22	2.15	0.46
1:A:356:ASP:OD2	1:A:356:ASP:N	2.48	0.46
1:A:390:ARG:HD3	1:A:396:TYR:O	2.15	0.46
1:A:176:TYR:CD1	1:A:262:PRO:HB3	2.52	0.45
1:B:117:SER:HA	1:B:120:ARG:NH1	2.31	0.45
1:B:45:TYR:CD2	1:B:52:VAL:CG2	3.00	0.45
1:B:397:SER:O	1:B:427:ARG:NH2	2.50	0.45
1:B:252:GLY:HA3	1:B:288:TYR:O	2.16	0.45
1:A:223:ARG:NH2	3:A:715:HOH:O	2.51	0.44
1:B:195:THR:HB	1:B:196:PRO:HD2	2.00	0.44
1:A:48:ASP:OD1	1:A:51:LYS:HE2	2.17	0.43
1:B:39:PRO:HG2	1:B:369:ASP:OD2	2.18	0.43
1:B:97:ARG:HG2	1:B:333:GLU:O	2.18	0.43
1:B:99:TYR:HB3	1:B:311:ILE:HG23	2.01	0.43
1:B:235:TYR:HA	1:B:239:LYS:O	2.18	0.42
1:B:120:ARG:HG3	1:B:121:PHE:N	2.34	0.42
1:A:118:LEU:HD23	1:A:118:LEU:HA	1.91	0.42
1:B:218:LEU:HD22	1:B:222:GLU:HB3	2.01	0.42
1:B:24:GLY:O	1:B:300:LYS:NZ	2.50	0.41
1:B:45:TYR:CE2	1:B:52:VAL:CG2	3.04	0.41
1:B:66:VAL:O	1:B:66:VAL:CG1	2.68	0.41
1:B:278:LEU:HG	1:B:289:TYR:CD2	2.55	0.41
1:B:380:GLN:NE2	3:B:721:HOH:O	2.52	0.41
1:B:474:ILE:HD12	1:B:474:ILE:HA	1.88	0.41
1:A:458:VAL:HB	1:A:500:ILE:HG12	2.03	0.41
1:B:368:VAL:HG13	1:B:369:ASP:N	2.35	0.41
1:B:397:SER:HB2	1:B:427:ARG:HH21	1.86	0.41
1:A:351:ILE:HD12	1:A:436:VAL:HA	2.03	0.41
1:B:330:LYS:HD2	1:B:330:LYS:HA	1.90	0.41
1:A:32:SER:HB2	1:A:49:GLU:OE1	2.20	0.40
1:B:18:ASN:ND2	3:B:709:HOH:O	2.39	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:446:GLU:O	1:B:471:ARG:HH21[4_555]	1.56	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	483/507 (95%)	471 (98%)	12 (2%)	0	100	100
1	B	487/507 (96%)	481 (99%)	6 (1%)	0	100	100
All	All	970/1014 (96%)	952 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/461 (97%)	440 (98%)	7 (2%)	62	76
1	B	448/461 (97%)	439 (98%)	9 (2%)	55	69
All	All	895/922 (97%)	879 (98%)	16 (2%)	59	72

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	SER
1	A	57	SER
1	A	97	ARG
1	A	200	SER
1	A	206	GLU
1	A	300	LYS
1	A	357	LYS
1	B	42	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	97	ARG
1	B	105	GLU
1	B	179	SER
1	B	181	GLU
1	B	402	ARG
1	B	443	SER
1	B	447	SER
1	B	468	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	331	HIS
1	A	395	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/507 (96%)	0.06	23 (4%) 31 30	19, 32, 69, 111	0
1	B	491/507 (96%)	0.20	24 (4%) 29 28	20, 37, 73, 124	0
All	All	980/1014 (96%)	0.13	47 (4%) 30 29	19, 34, 72, 124	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	354	ALA	7.1
1	B	352	THR	6.9
1	A	409	PRO	5.9
1	B	355	SER	5.5
1	A	408	ILE	4.7
1	B	343	ALA	4.5
1	B	351	ILE	4.5
1	A	352	THR	4.4
1	B	356	ASP	4.2
1	B	342	VAL	4.1
1	B	341	THR	4.0
1	A	351	ILE	3.6
1	B	346	ASP	3.6
1	A	0	ALA	3.5
1	B	54	LEU	3.5
1	B	422	LYS	3.4
1	A	341	THR	3.3
1	A	356	ASP	3.1
1	A	401	GLN	3.1
1	A	342	VAL	3.0
1	B	408	ILE	3.0
1	A	129	GLU	2.9
1	B	344	SER	2.9
1	A	406	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	463	LYS	2.8
1	A	50	ASP	2.8
1	A	423	LYS	2.8
1	B	32	SER	2.8
1	B	128	PRO	2.7
1	B	52	VAL	2.7
1	A	48	ASP	2.7
1	B	44	PHE	2.6
1	A	51	LYS	2.6
1	A	128	PRO	2.6
1	B	129	GLU	2.5
1	B	353	SER	2.5
1	B	406	LYS	2.5
1	A	52	VAL	2.5
1	A	29	LYS	2.4
1	A	463	LYS	2.3
1	A	130	PHE	2.3
1	A	407	GLU	2.3
1	A	180	ASP	2.2
1	A	344	SER	2.1
1	B	365	ILE	2.1
1	B	131	LYS	2.0
1	B	181	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

2	ZN	B	601	1/1	0.97	0.13	46,46,46,46	0
2	ZN	A	601	1/1	0.99	0.11	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.