



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 21, 2023 – 08:07 PM JST

PDB ID : 7WKX
Title : IL-17A in complex with the humanized antibody HB0017
Authors : Xu, J.; Zhu, X.; He, Y.
Deposited on : 2022-01-12
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3617 (2.84-2.80)
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	
1	B	217	
2	C	219	
2	D	219	
3	E	151	
3	F	151	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of HB0017 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	217	Total 1633	C 1039	N 263	O 325	S 6	0	0	0
1	B	212	Total 1595	C 1018	N 255	O 316	S 6	0	0	0

- Molecule 2 is a protein called Light chain of HB0017 Fab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	218	Total 1692	C 1066	N 281	O 340	S 5	0	2	0
2	D	216	Total 1663	C 1048	N 278	O 332	S 5	0	0	0

- Molecule 3 is a protein called Interleukin-17A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	95	Total 773	C 481	N 144	O 142	S 6	0	0	0
3	F	99	Total 802	C 499	N 149	O 147	S 7	0	0	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	5	MET	-	initiating methionine	UNP Q16552
E	6	GLY	-	expression tag	UNP Q16552
E	7	HIS	-	expression tag	UNP Q16552
E	8	HIS	-	expression tag	UNP Q16552
E	9	HIS	-	expression tag	UNP Q16552
E	10	HIS	-	expression tag	UNP Q16552
E	11	HIS	-	expression tag	UNP Q16552
E	12	HIS	-	expression tag	UNP Q16552

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	13	GLU	-	expression tag	UNP Q16552
E	14	ASN	-	expression tag	UNP Q16552
E	15	LEU	-	expression tag	UNP Q16552
E	16	TYR	-	expression tag	UNP Q16552
E	17	PHE	-	expression tag	UNP Q16552
E	18	GLN	-	expression tag	UNP Q16552
E	19	GLY	-	expression tag	UNP Q16552
F	5	MET	-	initiating methionine	UNP Q16552
F	6	GLY	-	expression tag	UNP Q16552
F	7	HIS	-	expression tag	UNP Q16552
F	8	HIS	-	expression tag	UNP Q16552
F	9	HIS	-	expression tag	UNP Q16552
F	10	HIS	-	expression tag	UNP Q16552
F	11	HIS	-	expression tag	UNP Q16552
F	12	HIS	-	expression tag	UNP Q16552
F	13	GLU	-	expression tag	UNP Q16552
F	14	ASN	-	expression tag	UNP Q16552
F	15	LEU	-	expression tag	UNP Q16552
F	16	TYR	-	expression tag	UNP Q16552
F	17	PHE	-	expression tag	UNP Q16552
F	18	GLN	-	expression tag	UNP Q16552
F	19	GLY	-	expression tag	UNP Q16552

- Molecule 4 is MALONIC ACID (three-letter code: MLA) (formula: C₃H₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 3 4	0	0
4	B	1	Total C O 7 3 4	0	0
4	C	1	Total C O 7 3 4	0	0
4	C	1	Total C O 7 3 4	0	0
4	C	1	Total C O 7 3 4	0	0
4	E	1	Total C O 7 3 4	0	0
4	F	1	Total C O 7 3 4	0	0

- Molecule 5 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	7	Total O 7 7	0	0
6	B	3	Total O 3 3	0	0
6	C	21	Total O 21 21	0	0
6	D	5	Total O 5 5	0	0
6	E	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heavy chain of HB0017 Fab




- Molecule 1: Heavy chain of HB0017 Fab




- Molecule 2: Light chain of HB0017 Fab




- Molecule 2: Light chain of HB0017 Fab



- Molecule 3: Interleukin-17A



- Molecule 3: Interleukin-17A



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	301.14Å 50.83Å 117.22Å 90.00° 101.27° 90.00°	Depositor
Resolution (Å)	45.20 – 2.81 45.16 – 2.81	Depositor EDS
% Data completeness (in resolution range)	95.1 (45.20-2.81) 95.1 (45.16-2.81)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.205 , 0.254 0.220 , 0.252	Depositor DCC
R_{free} test set	1997 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	62.3	Xtrriage
Anisotropy	0.125	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8260	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLA, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/1678	0.79	0/2290
1	B	0.62	0/1639	0.80	0/2237
2	C	0.66	0/1741	0.80	0/2370
2	D	0.62	0/1706	0.75	0/2325
3	E	0.59	0/791	0.72	0/1075
3	F	0.59	0/822	0.73	0/1120
All	All	0.63	0/8377	0.77	0/11417

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1633	0	1569	30	0
1	B	1595	0	1522	54	0
2	C	1692	0	1635	18	0
2	D	1663	0	1603	32	0
3	E	773	0	743	32	0
3	F	802	0	773	30	0
4	A	7	0	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	7	0	2	0	0
4	C	21	0	6	1	0
4	E	7	0	2	0	0
4	F	7	0	2	1	0
5	A	4	0	3	1	0
5	B	4	0	3	0	0
5	C	4	0	3	0	0
5	D	4	0	3	0	0
6	A	7	0	0	0	0
6	B	3	0	0	0	0
6	C	21	0	0	0	0
6	D	5	0	0	0	0
6	E	1	0	0	0	0
All	All	8260	0	7871	169	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 169 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:115:ILE:HD12	3:F:115:ILE:HD12	1.50	0.94
3:E:52:HIS:HB3	3:F:136:GLU:HG2	1.58	0.84
1:B:76:VAL:HG13	1:B:78:THR:HG22	1.67	0.77
1:A:154:VAL:CG2	1:A:182:LEU:HD21	2.17	0.74
1:A:18:VAL:HG23	1:A:86:LEU:HD11	1.68	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/217 (99%)	203 (94%)	12 (6%)	0	100	100
1	B	208/217 (96%)	192 (92%)	13 (6%)	3 (1%)	11	32
2	C	218/219 (100%)	210 (96%)	7 (3%)	1 (0%)	29	59
2	D	214/219 (98%)	203 (95%)	11 (5%)	0	100	100
3	E	89/151 (59%)	88 (99%)	1 (1%)	0	100	100
3	F	95/151 (63%)	91 (96%)	4 (4%)	0	100	100
All	All	1039/1174 (88%)	987 (95%)	48 (5%)	4 (0%)	34	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	7	SER
1	B	191	SER
1	B	177	SER
2	C	143	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/181 (100%)	180 (99%)	1 (1%)	86	95
1	B	175/181 (97%)	171 (98%)	4 (2%)	50	80
2	C	196/195 (100%)	195 (100%)	1 (0%)	88	96
2	D	192/195 (98%)	192 (100%)	0	100	100
3	E	90/140 (64%)	89 (99%)	1 (1%)	73	91
3	F	94/140 (67%)	94 (100%)	0	100	100
All	All	928/1032 (90%)	921 (99%)	7 (1%)	81	94

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	153	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	182	LEU
3	E	46	MET
2	C	113	ARG
1	B	89	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	47	GLN
2	D	215	ASN
2	C	59	HIS
3	E	111	ASN
2	D	14	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACY	B	302	-	3,3,3	1.16	0	3,3,3	0.68	0
4	MLA	C	304	-	6,6,6	1.46	1 (16%)	7,7,7	1.69	2 (28%)
4	MLA	C	303	-	6,6,6	1.50	0	7,7,7	1.29	1 (14%)
5	ACY	A	302	-	3,3,3	1.42	1 (33%)	3,3,3	0.93	0
5	ACY	C	301	-	3,3,3	1.32	0	3,3,3	1.05	0
5	ACY	D	301	-	3,3,3	1.13	0	3,3,3	0.92	0
4	MLA	E	201	-	6,6,6	1.53	0	7,7,7	1.26	0
4	MLA	F	201	-	6,6,6	1.66	1 (16%)	7,7,7	1.24	1 (14%)
4	MLA	C	302	-	6,6,6	1.64	2 (33%)	7,7,7	1.34	2 (28%)
4	MLA	B	301	-	6,6,6	2.57	2 (33%)	7,7,7	1.02	0
4	MLA	A	301	-	6,6,6	1.58	1 (16%)	7,7,7	1.66	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MLA	C	304	-	-	2/4/4/4	-
4	MLA	C	303	-	-	4/4/4/4	-
4	MLA	E	201	-	-	1/4/4/4	-
4	MLA	F	201	-	-	2/4/4/4	-
4	MLA	C	302	-	-	2/4/4/4	-
4	MLA	B	301	-	-	0/4/4/4	-
4	MLA	A	301	-	-	0/4/4/4	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	301	MLA	C2-C3	4.76	1.58	1.51
4	B	301	MLA	C2-C1	3.21	1.56	1.51
4	F	201	MLA	C2-C1	2.82	1.55	1.51
4	C	302	MLA	C2-C3	2.72	1.55	1.51
4	A	301	MLA	C2-C1	2.65	1.55	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	304	MLA	O3A-C3-C2	-3.00	113.33	122.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	MLA	O1B-C1-C2	-2.56	114.61	122.08
4	C	304	MLA	C3-C2-C1	-2.46	104.25	112.87
4	A	301	MLA	O3A-C3-C2	-2.30	115.35	122.08
4	C	302	MLA	O1B-C1-C2	-2.22	115.58	122.08

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	201	MLA	C1-C2-C3-O3B
4	F	201	MLA	C1-C2-C3-O3A
4	C	303	MLA	O1B-C1-C2-C3
4	C	303	MLA	O1A-C1-C2-C3
4	C	302	MLA	C1-C2-C3-O3A

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	302	ACY	1	0
4	F	201	MLA	1	0
4	C	302	MLA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/217 (100%)	-0.14	2 (0%) 84 80	35, 55, 82, 123	0
1	B	212/217 (97%)	-0.02	3 (1%) 75 69	47, 77, 106, 125	0
2	C	218/219 (99%)	-0.22	0 100 100	34, 47, 74, 103	0
2	D	216/219 (98%)	-0.15	1 (0%) 91 88	47, 72, 108, 134	0
3	E	95/151 (62%)	0.28	8 (8%) 11 5	53, 80, 121, 136	0
3	F	99/151 (65%)	0.24	3 (3%) 50 40	47, 75, 112, 128	0
All	All	1057/1174 (90%)	-0.06	17 (1%) 72 65	34, 65, 106, 136	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	128	HIS	5.8
1	A	136	SER	4.2
3	E	53	ASN	3.9
3	F	43	ARG	3.4
1	A	137	GLY	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACY	D	301	4/4	0.68	0.24	86,91,91,97	0
5	ACY	A	302	4/4	0.70	0.31	69,75,82,93	0
4	MLA	F	201	7/7	0.80	0.20	95,100,106,106	0
5	ACY	C	301	4/4	0.81	0.31	68,74,75,81	0
4	MLA	E	201	7/7	0.86	0.20	85,95,99,102	0
4	MLA	C	304	7/7	0.92	0.21	63,69,78,84	0
5	ACY	B	302	4/4	0.92	0.16	75,77,77,79	0
4	MLA	B	301	7/7	0.92	0.30	63,68,80,83	0
4	MLA	C	302	7/7	0.92	0.29	61,70,87,91	0
4	MLA	C	303	7/7	0.95	0.25	68,80,90,96	0
4	MLA	A	301	7/7	0.96	0.22	46,53,65,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.