



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 24, 2024 – 02:45 PM EDT

PDB ID : 5WL2  
Title : VH1-69 germline antibody with CDR H3 sequence of CR9114  
Authors : Wilson, I.A.; Lang, S.  
Deposited on : 2017-07-25  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	230	 80% 15% . .
1	H	230	 2% 83% 11% 5%
2	B	216	 % 81% 17% .
2	L	216	 2% 79% 17% . .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain of VH1-69 germline antibody with CDR H3 sequence of CR9114.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	219	Total 1622	C 1020	N 271	O 324	S 7	0	0	0
1	A	221	Total 1640	C 1033	N 273	O 327	S 7	0	0	0


- Molecule 2 is a protein called Germline-reverted light chain of CR9114.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	211	Total 1572	C 984	N 261	O 323	S 4	0	1	0
2	B	211	Total 1572	C 984	N 261	O 323	S 4	0	1	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	95	Total 95	O 95	0	0
3	L	46	Total 46	O 46	0	0
3	A	98	Total 98	O 98	0	0
3	B	57	Total 57	O 57	0	0





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.12Å 71.12Å 185.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.66 – 2.00 43.66 – 1.99	Depositor EDS
% Data completeness (in resolution range)	96.8 (43.66-2.00) 96.8 (43.66-1.99)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.74 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.200 , 0.236 0.200 , 0.235	Depositor DCC
$R_{free}$ test set	3653 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.2	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.021 for -h,-k,l 0.487 for h,-h-k,-l 0.024 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6702	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.58	2/1680 (0.1%)	0.70	3/2290 (0.1%)
1	H	0.58	2/1660 (0.1%)	0.71	3/2261 (0.1%)
2	B	0.43	0/1612	0.64	0/2204
2	L	0.48	0/1612	0.76	3/2204 (0.1%)
All	All	0.52	4/6564 (0.1%)	0.70	9/8959 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	113	GLN	C-N	-8.62	1.17	1.33
1	H	112	GLY	C-N	8.57	1.53	1.34
1	A	112	GLY	C-N	8.05	1.52	1.34
1	H	113	GLN	C-N	-7.84	1.19	1.33

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	191	ARG	NE-CZ-NH1	-13.85	113.38	120.30
1	A	112	GLY	C-N-CA	-8.07	101.53	121.70
1	H	112	GLY	C-N-CA	-7.68	102.51	121.70
1	A	67	ARG	NE-CZ-NH2	-6.57	117.02	120.30
2	L	191	ARG	NE-CZ-NH2	6.31	123.46	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	108	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1640	0	1598	29	0
1	H	1622	0	1582	17	0
2	B	1572	0	1527	28	0
2	L	1572	0	1527	36	0
3	A	98	0	0	4	0
3	B	57	0	0	7	1
3	H	95	0	0	1	1
3	L	46	0	0	6	0
All	All	6702	0	6234	107	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:153:ASP:HA	2:L:191:ARG:HH12	1.03	1.10
2:L:153:ASP:HA	2:L:191:ARG:NH1	1.74	1.03
2:L:162:GLU:OE2	3:L:301:HOH:O	1.83	0.96
1:A:217:LYS:NZ	2:B:125:GLU:OE2	2.00	0.95
1:A:28:THR:HG22	1:A:30:SER:H	1.35	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:381:HOH:O	3:B:356:HOH:O[3_555]	2.13	0.07



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/230 (95%)	212 (97%)	6 (3%)	1 (0%)	29	23
1	H	215/230 (94%)	212 (99%)	3 (1%)	0	100	100
2	B	210/216 (97%)	204 (97%)	6 (3%)	0	100	100
2	L	210/216 (97%)	204 (97%)	6 (3%)	0	100	100
All	All	854/892 (96%)	832 (97%)	21 (2%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	30	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/192 (95%)	180 (98%)	3 (2%)	62	67
1	H	181/192 (94%)	179 (99%)	2 (1%)	73	78
2	B	177/181 (98%)	174 (98%)	3 (2%)	60	65
2	L	177/181 (98%)	171 (97%)	6 (3%)	37	36
All	All	718/746 (96%)	704 (98%)	14 (2%)	55	61

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	191	ARG
1	A	67	ARG
2	B	177	SER
2	B	46	LEU
2	B	110	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	163	ASN
2	L	190	HIS
1	A	163	ASN
2	B	78	GLN
2	B	110	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	H	1
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	113:GLN	C	114:GLY	N	1.18
1	A	113:GLN	C	114:GLY	N	1.17

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/230 (96%)	-0.24	1 (0%) 91 90	35, 46, 84, 111	0
1	H	219/230 (95%)	-0.19	5 (2%) 60 59	34, 46, 82, 102	0
2	B	211/216 (97%)	0.09	3 (1%) 75 74	35, 62, 90, 96	0
2	L	211/216 (97%)	0.07	4 (1%) 66 65	35, 62, 89, 96	0
All	All	862/892 (96%)	-0.07	13 (1%) 73 72	34, 52, 88, 111	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	211	THR	3.9
1	H	139	THR	2.6
1	H	27	GLY	2.6
2	B	28	GLY	2.6
1	H	137	LYS	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.