



Full wwPDB EM Validation Report ⓘ

Nov 13, 2022 – 05:21 PM EST

PDB ID : 6WL1
EMDB ID : EMD-21815
Title : Cryo-EM of Form 1 related peptide filament, 36-31-3
Authors : Wang, F.; Gnewou, O.M.; Modlin, C.; Egelman, E.H.; Conticello, V.P.
Deposited on : 2020-04-17
Resolution : 4.00 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.


























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	36	100%
1	B	36	100%
1	C	36	100%
1	D	36	100%
1	E	36	100%
1	F	36	100%
1	G	36	100%
1	H	36	100%
1	I	36	100%










Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	36	6%  100%
1	K	36	8%  100%
1	L	36	8%  100%
1	M	36	6%  100%
1	N	36	8%  100%
1	O	36	8%  100%
1	P	36	8%  100%
1	Q	36	6%  100%
1	R	36	8%  100%
1	S	36	8%  100%
1	T	36	8%  100%
1	U	36	6%  100%
1	V	36	8%  100%
1	W	36	6%  100%
1	X	36	6%  100%
1	Y	36	8%  100%
1	Z	36	8%  100%
1	a	36	8%  100%
1	b	36	8%  100%
1	c	36	8%  100%
1	d	36	8%  100%
1	e	36	6%  100%
1	f	36	8%  100%
1	g	36	6%  100%
1	h	36	6%  100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	i	36	 8% 100%
1	j	36	 8% 100%
1	k	36	 8% 100%
1	l	36	 8% 100%
1	m	36	 6% 100%
1	n	36	 8% 100%
1	o	36	 6% 100%
1	p	36	 8% 100%
1	q	36	 8% 100%
1	r	36	 6% 100%
1	s	36	 6% 100%
1	t	36	 8% 100%
1	u	36	 6% 100%
1	v	36	 6% 100%
1	w	36	 8% 100%
1	x	36	 8% 100%
1	y	36	 6% 100%
1	z	36	 6% 100%

2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 14612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called peptide 36-31-3.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
1	A	36	281	179	50	52	0	0
1	B	36	281	179	50	52	0	0
1	C	36	281	179	50	52	0	0
1	D	36	281	179	50	52	0	0
1	E	36	281	179	50	52	0	0
1	F	36	281	179	50	52	0	0
1	G	36	281	179	50	52	0	0
1	H	36	281	179	50	52	0	0
1	I	36	281	179	50	52	0	0
1	J	36	281	179	50	52	0	0
1	K	36	281	179	50	52	0	0
1	L	36	281	179	50	52	0	0
1	M	36	281	179	50	52	0	0
1	N	36	281	179	50	52	0	0
1	O	36	281	179	50	52	0	0
1	P	36	281	179	50	52	0	0
1	Q	36	281	179	50	52	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf	Trace
1	R	36	Total 281	C 179	N 50	O 52	0	0
1	S	36	Total 281	C 179	N 50	O 52	0	0
1	T	36	Total 281	C 179	N 50	O 52	0	0
1	U	36	Total 281	C 179	N 50	O 52	0	0
1	V	36	Total 281	C 179	N 50	O 52	0	0
1	W	36	Total 281	C 179	N 50	O 52	0	0
1	X	36	Total 281	C 179	N 50	O 52	0	0
1	Y	36	Total 281	C 179	N 50	O 52	0	0
1	Z	36	Total 281	C 179	N 50	O 52	0	0
1	a	36	Total 281	C 179	N 50	O 52	0	0
1	b	36	Total 281	C 179	N 50	O 52	0	0
1	c	36	Total 281	C 179	N 50	O 52	0	0
1	d	36	Total 281	C 179	N 50	O 52	0	0
1	e	36	Total 281	C 179	N 50	O 52	0	0
1	f	36	Total 281	C 179	N 50	O 52	0	0
1	g	36	Total 281	C 179	N 50	O 52	0	0
1	h	36	Total 281	C 179	N 50	O 52	0	0
1	i	36	Total 281	C 179	N 50	O 52	0	0
1	j	36	Total 281	C 179	N 50	O 52	0	0
1	k	36	Total 281	C 179	N 50	O 52	0	0
1	l	36	Total 281	C 179	N 50	O 52	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf	Trace
1	m	36	Total	C	N	O	0	0
			281	179	50	52		
1	n	36	Total	C	N	O	0	0
			281	179	50	52		
1	o	36	Total	C	N	O	0	0
			281	179	50	52		
1	p	36	Total	C	N	O	0	0
			281	179	50	52		
1	q	36	Total	C	N	O	0	0
			281	179	50	52		
1	r	36	Total	C	N	O	0	0
			281	179	50	52		
1	s	36	Total	C	N	O	0	0
			281	179	50	52		
1	t	36	Total	C	N	O	0	0
			281	179	50	52		
1	u	36	Total	C	N	O	0	0
			281	179	50	52		
1	v	36	Total	C	N	O	0	0
			281	179	50	52		
1	w	36	Total	C	N	O	0	0
			281	179	50	52		
1	x	36	Total	C	N	O	0	0
			281	179	50	52		
1	y	36	Total	C	N	O	0	0
			281	179	50	52		
1	z	36	Total	C	N	O	0	0
			281	179	50	52		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3





- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3






- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3







- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3




- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



- Molecule 1: peptide 36-31-3



4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=124°, rise=2.5 Å, axial sym=C1	Depositor
Number of segments used	66079	Depositor
Resolution determination method	OTHER	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.002	Depositor
Minimum map value	-0.001	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.000462	Depositor
Map size (Å)	211.99998, 211.99998, 211.99998	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/282	0.48	0/378
1	B	0.36	0/282	0.48	0/378
1	C	0.36	0/282	0.48	0/378
1	D	0.36	0/282	0.48	0/378
1	E	0.36	0/282	0.48	0/378
1	F	0.36	0/282	0.48	0/378
1	G	0.36	0/282	0.48	0/378
1	H	0.36	0/282	0.48	0/378
1	I	0.36	0/282	0.48	0/378
1	J	0.36	0/282	0.48	0/378
1	K	0.36	0/282	0.48	0/378
1	L	0.36	0/282	0.48	0/378
1	M	0.36	0/282	0.48	0/378
1	N	0.36	0/282	0.48	0/378
1	O	0.36	0/282	0.48	0/378
1	P	0.36	0/282	0.48	0/378
1	Q	0.36	0/282	0.48	0/378
1	R	0.36	0/282	0.48	0/378
1	S	0.36	0/282	0.48	0/378
1	T	0.36	0/282	0.48	0/378
1	U	0.36	0/282	0.48	0/378
1	V	0.36	0/282	0.48	0/378
1	W	0.36	0/282	0.48	0/378
1	X	0.36	0/282	0.48	0/378
1	Y	0.36	0/282	0.48	0/378
1	Z	0.36	0/282	0.48	0/378
1	a	0.36	0/282	0.48	0/378
1	b	0.36	0/282	0.48	0/378
1	c	0.36	0/282	0.48	0/378
1	d	0.36	0/282	0.48	0/378
1	e	0.36	0/282	0.48	0/378
1	f	0.36	0/282	0.48	0/378
1	g	0.36	0/282	0.48	0/378
1	h	0.36	0/282	0.48	0/378

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	i	0.36	0/282	0.48	0/378
1	j	0.36	0/282	0.48	0/378
1	k	0.36	0/282	0.48	0/378
1	l	0.36	0/282	0.48	0/378
1	m	0.36	0/282	0.48	0/378
1	n	0.36	0/282	0.48	0/378
1	o	0.36	0/282	0.48	0/378
1	p	0.36	0/282	0.48	0/378
1	q	0.36	0/282	0.48	0/378
1	r	0.36	0/282	0.48	0/378
1	s	0.36	0/282	0.48	0/378
1	t	0.36	0/282	0.48	0/378
1	u	0.36	0/282	0.48	0/378
1	v	0.36	0/282	0.48	0/378
1	w	0.36	0/282	0.48	0/378
1	x	0.36	0/282	0.48	0/378
1	y	0.36	0/282	0.48	0/378
1	z	0.36	0/282	0.48	0/378
All	All	0.36	0/14664	0.48	0/19656

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	34/36 (94%)	34 (100%)	0	0	100	100
1	B	34/36 (94%)	34 (100%)	0	0	100	100
1	C	34/36 (94%)	34 (100%)	0	0	100	100
1	D	34/36 (94%)	34 (100%)	0	0	100	100
1	E	34/36 (94%)	34 (100%)	0	0	100	100
1	F	34/36 (94%)	34 (100%)	0	0	100	100
1	G	34/36 (94%)	34 (100%)	0	0	100	100
1	H	34/36 (94%)	34 (100%)	0	0	100	100
1	I	34/36 (94%)	34 (100%)	0	0	100	100
1	J	34/36 (94%)	34 (100%)	0	0	100	100
1	K	34/36 (94%)	34 (100%)	0	0	100	100
1	L	34/36 (94%)	34 (100%)	0	0	100	100
1	M	34/36 (94%)	34 (100%)	0	0	100	100
1	N	34/36 (94%)	34 (100%)	0	0	100	100
1	O	34/36 (94%)	34 (100%)	0	0	100	100
1	P	34/36 (94%)	34 (100%)	0	0	100	100
1	Q	34/36 (94%)	34 (100%)	0	0	100	100
1	R	34/36 (94%)	34 (100%)	0	0	100	100
1	S	34/36 (94%)	34 (100%)	0	0	100	100
1	T	34/36 (94%)	34 (100%)	0	0	100	100
1	U	34/36 (94%)	34 (100%)	0	0	100	100
1	V	34/36 (94%)	34 (100%)	0	0	100	100
1	W	34/36 (94%)	34 (100%)	0	0	100	100
1	X	34/36 (94%)	34 (100%)	0	0	100	100
1	Y	34/36 (94%)	34 (100%)	0	0	100	100
1	Z	34/36 (94%)	34 (100%)	0	0	100	100
1	a	34/36 (94%)	34 (100%)	0	0	100	100
1	b	34/36 (94%)	34 (100%)	0	0	100	100
1	c	34/36 (94%)	34 (100%)	0	0	100	100
1	d	34/36 (94%)	34 (100%)	0	0	100	100
1	e	34/36 (94%)	34 (100%)	0	0	100	100
1	f	34/36 (94%)	34 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	g	34/36 (94%)	34 (100%)	0	0	100	100
1	h	34/36 (94%)	34 (100%)	0	0	100	100
1	i	34/36 (94%)	34 (100%)	0	0	100	100
1	j	34/36 (94%)	34 (100%)	0	0	100	100
1	k	34/36 (94%)	34 (100%)	0	0	100	100
1	l	34/36 (94%)	34 (100%)	0	0	100	100
1	m	34/36 (94%)	34 (100%)	0	0	100	100
1	n	34/36 (94%)	34 (100%)	0	0	100	100
1	o	34/36 (94%)	34 (100%)	0	0	100	100
1	p	34/36 (94%)	34 (100%)	0	0	100	100
1	q	34/36 (94%)	34 (100%)	0	0	100	100
1	r	34/36 (94%)	34 (100%)	0	0	100	100
1	s	34/36 (94%)	34 (100%)	0	0	100	100
1	t	34/36 (94%)	34 (100%)	0	0	100	100
1	u	34/36 (94%)	34 (100%)	0	0	100	100
1	v	34/36 (94%)	34 (100%)	0	0	100	100
1	w	34/36 (94%)	34 (100%)	0	0	100	100
1	x	34/36 (94%)	34 (100%)	0	0	100	100
1	y	34/36 (94%)	34 (100%)	0	0	100	100
1	z	34/36 (94%)	34 (100%)	0	0	100	100
All	All	1768/1872 (94%)	1768 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	26/26 (100%)	26 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	26/26 (100%)	26 (100%)	0	100	100
1	C	26/26 (100%)	26 (100%)	0	100	100
1	D	26/26 (100%)	26 (100%)	0	100	100
1	E	26/26 (100%)	26 (100%)	0	100	100
1	F	26/26 (100%)	26 (100%)	0	100	100
1	G	26/26 (100%)	26 (100%)	0	100	100
1	H	26/26 (100%)	26 (100%)	0	100	100
1	I	26/26 (100%)	26 (100%)	0	100	100
1	J	26/26 (100%)	26 (100%)	0	100	100
1	K	26/26 (100%)	26 (100%)	0	100	100
1	L	26/26 (100%)	26 (100%)	0	100	100
1	M	26/26 (100%)	26 (100%)	0	100	100
1	N	26/26 (100%)	26 (100%)	0	100	100
1	O	26/26 (100%)	26 (100%)	0	100	100
1	P	26/26 (100%)	26 (100%)	0	100	100
1	Q	26/26 (100%)	26 (100%)	0	100	100
1	R	26/26 (100%)	26 (100%)	0	100	100
1	S	26/26 (100%)	26 (100%)	0	100	100
1	T	26/26 (100%)	26 (100%)	0	100	100
1	U	26/26 (100%)	26 (100%)	0	100	100
1	V	26/26 (100%)	26 (100%)	0	100	100
1	W	26/26 (100%)	26 (100%)	0	100	100
1	X	26/26 (100%)	26 (100%)	0	100	100
1	Y	26/26 (100%)	26 (100%)	0	100	100
1	Z	26/26 (100%)	26 (100%)	0	100	100
1	a	26/26 (100%)	26 (100%)	0	100	100
1	b	26/26 (100%)	26 (100%)	0	100	100
1	c	26/26 (100%)	26 (100%)	0	100	100
1	d	26/26 (100%)	26 (100%)	0	100	100
1	e	26/26 (100%)	26 (100%)	0	100	100
1	f	26/26 (100%)	26 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	g	26/26 (100%)	26 (100%)	0	100	100
1	h	26/26 (100%)	26 (100%)	0	100	100
1	i	26/26 (100%)	26 (100%)	0	100	100
1	j	26/26 (100%)	26 (100%)	0	100	100
1	k	26/26 (100%)	26 (100%)	0	100	100
1	l	26/26 (100%)	26 (100%)	0	100	100
1	m	26/26 (100%)	26 (100%)	0	100	100
1	n	26/26 (100%)	26 (100%)	0	100	100
1	o	26/26 (100%)	26 (100%)	0	100	100
1	p	26/26 (100%)	26 (100%)	0	100	100
1	q	26/26 (100%)	26 (100%)	0	100	100
1	r	26/26 (100%)	26 (100%)	0	100	100
1	s	26/26 (100%)	26 (100%)	0	100	100
1	t	26/26 (100%)	26 (100%)	0	100	100
1	u	26/26 (100%)	26 (100%)	0	100	100
1	v	26/26 (100%)	26 (100%)	0	100	100
1	w	26/26 (100%)	26 (100%)	0	100	100
1	x	26/26 (100%)	26 (100%)	0	100	100
1	y	26/26 (100%)	26 (100%)	0	100	100
1	z	26/26 (100%)	26 (100%)	0	100	100
All	All	1352/1352 (100%)	1352 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21815. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 121



Y Index: 121



Z Index: 18


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.000462. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 101 nm³; this corresponds to an approximate mass of 91 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.250\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-21815 and PDB model 6WL1. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.000462 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.000462).

9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary





































The table lists the average atom inclusion at the recommended contour level (0.000462) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5920	0.4630
A	0.5978	0.4610
B	0.6014	0.4630
C	0.5906	0.4660
D	0.5978	0.4600
E	0.5906	0.4620
F	0.5906	0.4600
G	0.5833	0.4630
H	0.5978	0.4610
I	0.5942	0.4620
J	0.5978	0.4630
K	0.5906	0.4640
L	0.5906	0.4610
M	0.5942	0.4640
N	0.5870	0.4590
O	0.5870	0.4630
P	0.5942	0.4670
Q	0.5942	0.4590
R	0.5906	0.4600
S	0.5906	0.4670
T	0.5942	0.4610
U	0.5978	0.4590
V	0.5942	0.4680
W	0.5906	0.4600
X	0.5942	0.4700
Y	0.5906	0.4630
Z	0.5942	0.4640
a	0.5942	0.4580
b	0.5978	0.4640
c	0.5978	0.4610
d	0.5833	0.4660
e	0.5978	0.4680
f	0.5833	0.4590
g	0.5870	0.4600
h	0.5870	0.4690



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
i	 0.5833	 0.4660
j	 0.5797	 0.4620
k	 0.5797	 0.4650
l	 0.5870	 0.4610
m	 0.5870	 0.4610
n	 0.6051	 0.4660
o	 0.5942	 0.4610
p	 0.5942	 0.4600
q	 0.5906	 0.4620
r	 0.5978	 0.4600
s	 0.6051	 0.4560
t	 0.5906	 0.4660
u	 0.5942	 0.4610
v	 0.5942	 0.4590
w	 0.5906	 0.4590
x	 0.5833	 0.4640
y	 0.5906	 0.4610
z	 0.5942	 0.4680