



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2023 – 01:13 am GMT

PDB ID : 2WME
Title : Crystallographic structure of betaine aldehyde dehydrogenase from *Pseudomonas aeruginosa*
Authors : Gonzalez-Segura, L.; Rudino-Pinera, E.; Munoz-Clares, R.A.; Horjales, E.
Deposited on : 2009-06-30
Resolution : 2.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	88% 11% .
1	B	490	91% 8% .
1	D	490	91% 8% .
1	F	490	85% 13% .
1	G	490	88% 10% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	490	
2	C	490	
2	E	490	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	1501	-	-	X	-
6	GOL	D	1496	-	-	X	-
6	GOL	E	1496	-	-	X	X
6	GOL	G	1495	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 33548 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETAINE ALDEHYDE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	489	Total 3824	C 2398	N 673	O 739	S 14	0	10	0
1	B	489	Total 3800	C 2386	N 664	O 736	S 14	0	7	0
1	D	489	Total 3744	C 2351	N 657	O 722	S 14	0	0	0
1	F	489	Total 3789	C 2378	N 666	O 731	S 14	0	5	0
1	G	489	Total 3753	C 2356	N 658	O 725	S 14	0	1	0
1	H	489	Total 3753	C 2356	N 658	O 725	S 14	0	1	0

- Molecule 2 is a protein called BETAINE ALDEHYDE DEHYDROGENASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	489	Total 3768	C 2364	N 660	O 729	S 15	0	3	0
2	E	489	Total 3767	C 2364	N 660	O 729	S 14	0	3	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	E	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	F	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	G	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	H	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		
4	D	1	Total	C	O	S	0	1
			8	4	2	2		
4	F	1	Total	C	O	S	0	0
			4	2	1	1		
4	G	1	Total	C	O	S	0	1
			8	4	2	2		

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	K	0	0
			2	2		
5	B	2	Total	K	0	0
			2	2		
5	C	2	Total	K	0	0
			2	2		
5	D	2	Total	K	0	0
			2	2		
5	E	2	Total	K	0	0
			2	2		
5	F	2	Total	K	0	0
			2	2		
5	G	2	Total	K	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	2	Total 2 2	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	C	1	Total C O 6 3 3	0	0
6	C	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0
6	D	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	E	1	Total C O 6 3 3	0	0
6	F	1	Total C O 6 3 3	0	0
6	G	1	Total C O 6 3 3	0	0
6	G	1	Total C O 6 3 3	0	0
6	G	1	Total C O 6 3 3	0	0
6	H	1	Total C O 6 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	417	Total O 417 417	0	0
7	B	405	Total O 405 405	0	0
7	C	367	Total O 367 367	0	0
7	D	362	Total O 362 362	0	0
7	E	357	Total O 357 357	0	0
7	F	347	Total O 347 347	0	0
7	G	360	Total O 360 360	0	0
7	H	317	Total O 317 317	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE




- Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE




- Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE





- Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE









• Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE




• Molecule 1: BETAINE ALDEHYDE DEHYDROGENASE



• Molecule 2: BETAINE ALDEHYDE DEHYDROGENASE



• Molecule 2: BETAINE ALDEHYDE DEHYDROGENASE

Chain E:  88% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	334.95Å 133.01Å 101.81Å 90.00° 94.94° 90.00°	Depositor
Resolution (Å)	42.68 – 2.10 42.66 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.2 (42.68-2.10) 95.1 (42.66-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.166 , 0.211 0.167 , 0.223	Depositor DCC
R_{free} test set	12459 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtrriage
Anisotropy	0.067	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	33548	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, GOL, K, NAP, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	0/3892	0.91	11/5273 (0.2%)
1	B	1.00	2/3868 (0.1%)	0.87	9/5242 (0.2%)
1	D	0.98	0/3812	0.82	5/5166 (0.1%)
1	F	0.92	0/3857	0.79	4/5225 (0.1%)
1	G	0.95	3/3821 (0.1%)	0.82	3/5178 (0.1%)
1	H	0.85	1/3821 (0.0%)	0.79	2/5178 (0.0%)
2	C	0.95	1/3821 (0.0%)	0.84	8/5177 (0.2%)
2	E	0.95	1/3827 (0.0%)	0.87	12/5185 (0.2%)
All	All	0.96	8/30719 (0.0%)	0.84	54/41624 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	248	GLU	CB-CG	6.22	1.64	1.52
1	G	412	GLU	CG-CD	6.12	1.61	1.51
1	H	127	GLN	CB-CG	5.33	1.67	1.52
2	E	44	GLU	CG-CD	5.31	1.59	1.51
1	B	440	TRP	CB-CG	5.28	1.59	1.50

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	475	ARG	NE-CZ-NH1	12.35	126.47	120.30
1	B	475	ARG	NE-CZ-NH1	11.50	126.05	120.30
1	A	475	ARG	NE-CZ-NH2	-10.35	115.12	120.30
2	E	3	ARG	NE-CZ-NH1	10.27	125.43	120.30
1	A	407	ARG	NE-CZ-NH1	10.27	125.43	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3824	0	3787	46	0
1	B	3800	0	3761	29	0
1	D	3744	0	3714	27	0
1	F	3789	0	3758	53	0
1	G	3753	0	3720	40	0
1	H	3753	0	3720	70	0
2	C	3768	0	3732	31	0
2	E	3767	0	3732	41	0
3	A	31	0	11	9	0
3	B	31	0	11	4	0
3	C	31	0	11	0	0
3	D	31	0	11	7	0
3	E	31	0	11	2	0
3	F	31	0	11	3	0
3	G	31	0	11	1	0
3	H	31	0	11	2	0
4	A	4	0	6	3	0
4	B	4	0	5	1	0
4	D	8	0	11	0	0
4	F	4	0	6	2	0
4	G	8	0	12	3	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
5	D	2	0	0	0	0
5	E	2	0	0	0	0
5	F	2	0	0	0	0
5	G	2	0	0	0	0
5	H	2	0	0	0	0
6	A	36	0	48	6	0
6	B	18	0	24	3	0
6	C	18	0	24	2	0
6	D	12	0	16	7	0
6	E	12	0	16	6	0
6	F	6	0	8	1	0
6	G	18	0	24	4	0
6	H	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	417	0	0	2	0
7	B	405	0	0	6	0
7	C	367	0	0	9	0
7	D	362	0	0	10	0
7	E	357	0	0	8	0
7	F	347	0	0	8	0
7	G	360	0	0	6	0
7	H	317	0	0	17	0
All	All	33548	0	30220	348	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 348 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:297:SER:HB3	7:C:2223:HOH:O	1.32	1.25
1:G:73:ARG:HD2	7:G:2072:HOH:O	1.34	1.25
1:G:43:ARG:HG2	1:G:43:ARG:HH11	1.07	1.09
1:A:286:CYS:SG	4:A:1492:BME:S2	2.46	1.07
1:G:335:GLU:HG2	7:G:2256:HOH:O	1.55	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/490 (101%)	480 (97%)	17 (3%)	0	100	100
1	B	494/490 (101%)	480 (97%)	14 (3%)	0	100	100
1	D	487/490 (99%)	470 (96%)	17 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	492/490 (100%)	481 (98%)	11 (2%)	0	100	100
1	G	488/490 (100%)	474 (97%)	13 (3%)	1 (0%)	47	49
1	H	488/490 (100%)	462 (95%)	25 (5%)	1 (0%)	47	49
2	C	488/490 (100%)	473 (97%)	14 (3%)	1 (0%)	47	49
2	E	489/490 (100%)	468 (96%)	21 (4%)	0	100	100
All	All	3923/3920 (100%)	3788 (97%)	132 (3%)	3 (0%)	51	54

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	363	PHE
2	C	414	GLY
1	G	414	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/391 (102%)	390 (98%)	10 (2%)	47	52
1	B	397/391 (102%)	385 (97%)	12 (3%)	41	44
1	D	390/391 (100%)	383 (98%)	7 (2%)	59	65
1	F	395/391 (101%)	382 (97%)	13 (3%)	38	40
1	G	391/391 (100%)	381 (97%)	10 (3%)	46	50
1	H	391/391 (100%)	375 (96%)	16 (4%)	30	31
2	C	391/390 (100%)	380 (97%)	11 (3%)	43	47
2	E	392/390 (100%)	379 (97%)	13 (3%)	38	40
All	All	3147/3126 (101%)	3055 (97%)	92 (3%)	42	46

5 of 92 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	210	ARG
1	G	322	ASN
1	F	335	GLU
1	F	475	ARG
1	G	475	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	284	GLN
1	F	465	ASN
2	E	295	HIS
1	F	219	HIS
1	G	83	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSO	C	286[B]	2	3,6,7	0.76	0	0,6,8	-	-
2	CSO	E	286	2	3,6,7	0.93	0	0,6,8	-	-
2	CSO	C	286[A]	2	3,6,7	0.84	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	C	286[B]	2	-	0/1/5/7	-
2	CSO	E	286	2	-	0/1/5/7	-
2	CSO	C	286[A]	2	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	286[B]	CSO	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 52 ligands modelled in this entry, 16 are monoatomic - leaving 36 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	E	1491	-	27,33,52	1.13	2 (7%)	35,52,80	1.45	6 (17%)
6	GOL	D	1495	-	5,5,5	0.30	0	5,5,5	0.26	0
6	GOL	C	1497	-	5,5,5	0.47	0	5,5,5	0.75	0
4	BME	D	1492[B]	-	3,3,3	0.63	0	1,2,2	0.06	0
6	GOL	E	1495	-	5,5,5	0.33	0	5,5,5	0.46	0
6	GOL	E	1496	-	5,5,5	0.46	0	5,5,5	0.79	0
4	BME	G	1492[A]	-	3,3,3	0.51	0	1,2,2	0.56	0
6	GOL	B	1497	-	5,5,5	0.38	0	5,5,5	0.83	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	F	1491	-	27,33,52	1.05	2 (7%)	35,52,80	1.65	6 (17%)
6	GOL	A	1497	-	5,5,5	0.48	0	5,5,5	0.35	0
3	NAP	H	1491	-	27,33,52	0.92	1 (3%)	35,52,80	1.28	2 (5%)
4	BME	D	1492[A]	-	3,3,3	0.21	0	1,2,2	0.93	0
6	GOL	A	1498	-	5,5,5	0.53	0	5,5,5	0.60	0
6	GOL	G	1495	-	5,5,5	0.48	0	5,5,5	0.79	0
6	GOL	A	1499	-	5,5,5	0.21	0	5,5,5	0.86	0
4	BME	A	1492	-	3,3,3	0.29	0	1,2,2	1.09	0
6	GOL	C	1496	-	5,5,5	0.78	0	5,5,5	0.57	0
6	GOL	B	1496	-	5,5,5	1.06	0	5,5,5	1.39	1 (20%)
6	GOL	H	1495	-	5,5,5	0.61	0	5,5,5	0.64	0
6	GOL	C	1495	-	5,5,5	0.63	0	5,5,5	0.56	0
6	GOL	D	1496	-	5,5,5	0.67	0	5,5,5	0.51	0
4	BME	F	1492	-	3,3,3	0.79	0	1,2,2	0.65	0
6	GOL	G	1497	-	5,5,5	0.53	0	5,5,5	0.44	0
3	NAP	B	1491	-	27,33,52	1.06	2 (7%)	35,52,80	1.37	5 (14%)
3	NAP	A	1491	-	27,33,52	1.06	2 (7%)	35,52,80	1.49	5 (14%)
6	GOL	B	1495	-	5,5,5	0.63	0	5,5,5	0.59	0
6	GOL	A	1496	-	5,5,5	0.73	0	5,5,5	0.97	0
6	GOL	A	1500	-	5,5,5	0.89	0	5,5,5	1.38	1 (20%)
3	NAP	G	1491	-	27,33,52	1.05	2 (7%)	35,52,80	1.60	5 (14%)
6	GOL	F	1495	-	5,5,5	0.37	0	5,5,5	0.60	0
6	GOL	G	1496	-	5,5,5	0.39	0	5,5,5	0.75	0
3	NAP	D	1491	-	27,33,52	0.96	1 (3%)	35,52,80	1.36	4 (11%)
4	BME	G	1492[B]	-	3,3,3	0.51	0	1,2,2	0.13	0
4	BME	B	1492	-	3,3,3	0.54	0	1,2,2	0.57	0
3	NAP	C	1491	-	27,33,52	1.12	2 (7%)	35,52,80	1.57	6 (17%)
6	GOL	A	1501	-	5,5,5	0.56	0	5,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	E	1491	-	-	3/17/37/67	0/3/3/5
6	GOL	D	1495	-	-	4/4/4/4	-
6	GOL	C	1497	-	-	4/4/4/4	-
4	BME	D	1492[B]	-	-	0/1/1/1	-
6	GOL	E	1495	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	E	1496	-	-	2/4/4/4	-
4	BME	G	1492[A]	-	-	0/1/1/1	-
6	GOL	B	1497	-	-	2/4/4/4	-
3	NAP	F	1491	-	-	4/17/37/67	0/3/3/5
6	GOL	A	1497	-	-	2/4/4/4	-
3	NAP	H	1491	-	-	2/17/37/67	0/3/3/5
4	BME	D	1492[A]	-	-	1/1/1/1	-
6	GOL	A	1498	-	-	2/4/4/4	-
6	GOL	G	1495	-	-	2/4/4/4	-
6	GOL	A	1499	-	-	0/4/4/4	-
4	BME	A	1492	-	-	0/1/1/1	-
6	GOL	C	1496	-	-	2/4/4/4	-
6	GOL	B	1496	-	-	4/4/4/4	-
6	GOL	H	1495	-	-	2/4/4/4	-
6	GOL	C	1495	-	-	2/4/4/4	-
6	GOL	D	1496	-	-	2/4/4/4	-
4	BME	F	1492	-	-	1/1/1/1	-
6	GOL	G	1497	-	-	2/4/4/4	-
3	NAP	B	1491	-	-	2/17/37/67	0/3/3/5
3	NAP	A	1491	-	-	4/17/37/67	0/3/3/5
6	GOL	B	1495	-	-	4/4/4/4	-
6	GOL	A	1496	-	-	2/4/4/4	-
6	GOL	A	1500	-	-	2/4/4/4	-
3	NAP	G	1491	-	-	0/17/37/67	0/3/3/5
6	GOL	F	1495	-	-	2/4/4/4	-
6	GOL	G	1496	-	-	4/4/4/4	-
3	NAP	D	1491	-	-	10/17/37/67	0/3/3/5
4	BME	G	1492[B]	-	-	1/1/1/1	-
4	BME	B	1492	-	-	1/1/1/1	-
3	NAP	C	1491	-	-	7/17/37/67	0/3/3/5
6	GOL	A	1501	-	-	0/4/4/4	-

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1491	NAP	O4B-C1B	3.30	1.45	1.41
3	C	1491	NAP	O4B-C1B	3.01	1.45	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	1491	NAP	P2B-O2B	2.82	1.64	1.59
3	A	1491	NAP	P2B-O2B	2.80	1.64	1.59
3	G	1491	NAP	O4B-C1B	2.76	1.44	1.41

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1491	NAP	N3A-C2A-N1A	-5.60	119.92	128.68
3	D	1491	NAP	N3A-C2A-N1A	-5.53	120.03	128.68
3	F	1491	NAP	N3A-C2A-N1A	-5.27	120.45	128.68
3	G	1491	NAP	N3A-C2A-N1A	-5.16	120.61	128.68
3	C	1491	NAP	N3A-C2A-N1A	-4.94	120.96	128.68

There are no chirality outliers.

5 of 84 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1491	NAP	C5B-O5B-PA-O3
3	C	1491	NAP	PA-O3-PN-O5D
3	E	1491	NAP	PA-O3-PN-O2N
4	D	1492[A]	BME	O1-C1-C2-S2
4	F	1492	BME	O1-C1-C2-S2

There are no ring outliers.

25 monomers are involved in 61 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1491	NAP	2	0
6	D	1495	GOL	1	0
6	C	1497	GOL	1	0
6	E	1495	GOL	1	0
6	E	1496	GOL	5	0
4	G	1492[A]	BME	1	0
3	F	1491	NAP	3	0
3	H	1491	NAP	2	0
6	G	1495	GOL	4	0
4	A	1492	BME	3	0
6	C	1496	GOL	1	0
6	B	1496	GOL	1	0
6	H	1495	GOL	1	0
6	D	1496	GOL	6	0


Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	1492	BME	2	0
3	B	1491	NAP	4	0
3	A	1491	NAP	9	0
6	B	1495	GOL	3	0
3	G	1491	NAP	1	0
6	F	1495	GOL	1	0
6	G	1496	GOL	1	0
3	D	1491	NAP	7	0
4	G	1492[B]	BME	2	0
4	B	1492	BME	1	0
6	A	1501	GOL	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/490 (99%)	-0.21	1 (0%) 95 95	9, 20, 25, 36	0
1	B	489/490 (99%)	-0.17	0 100 100	10, 20, 26, 36	0
1	D	489/490 (99%)	-0.12	3 (0%) 89 91	10, 20, 27, 32	0
1	F	489/490 (99%)	0.07	12 (2%) 57 62	12, 21, 28, 34	0
1	G	489/490 (99%)	-0.14	2 (0%) 92 93	12, 20, 27, 35	0
1	H	489/490 (99%)	0.95	85 (17%) 1 1	15, 23, 31, 37	0
2	C	488/490 (99%)	-0.09	5 (1%) 82 85	11, 20, 27, 36	0
2	E	488/490 (99%)	-0.19	1 (0%) 95 95	10, 20, 26, 33	0
All	All	3910/3920 (99%)	0.01	109 (2%) 53 59	9, 20, 28, 37	0

The worst 5 of 109 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	362	ALA	5.8
1	H	359	THR	5.5
1	H	363	PHE	5.5
1	H	98	LEU	4.5
1	H	382	THR	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	C	286[A]	7/8	0.91	0.21	15,22,23,25	7

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	C	286[B]	7/8	0.91	0.21	22,23,25,26	7
2	CSO	E	286	7/8	0.93	0.12	19,22,26,30	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	C	1496	6/6	0.77	0.20	37,38,40,42	0
3	NAP	H	1491	31/48	0.79	0.28	24,33,43,44	31
6	GOL	C	1497	6/6	0.79	0.18	31,34,35,41	0
6	GOL	E	1496	6/6	0.79	0.46	41,45,46,47	0
6	GOL	H	1495	6/6	0.79	0.22	41,46,46,47	0
6	GOL	G	1497	6/6	0.82	0.21	33,34,35,36	0
6	GOL	A	1500	6/6	0.82	0.17	25,29,30,33	0
6	GOL	A	1499	6/6	0.84	0.18	33,35,38,41	0
4	BME	G	1492[B]	4/4	0.85	0.27	37,38,39,41	4
6	GOL	D	1496	6/6	0.85	0.16	27,30,30,33	0
4	BME	D	1492[A]	4/4	0.85	0.25	28,29,30,34	4
4	BME	D	1492[B]	4/4	0.85	0.25	18,19,20,21	4
4	BME	G	1492[A]	4/4	0.85	0.27	21,23,23,24	4
6	GOL	C	1495	6/6	0.86	0.19	42,47,48,50	0
6	GOL	B	1496	6/6	0.87	0.15	30,31,32,33	0
6	GOL	A	1501	6/6	0.88	0.19	34,41,43,43	0
6	GOL	B	1497	6/6	0.89	0.22	27,38,42,44	0
4	BME	F	1492	4/4	0.89	0.18	39,42,44,48	0
6	GOL	A	1496	6/6	0.89	0.17	21,27,28,30	0
3	NAP	A	1491	31/48	0.89	0.20	21,28,45,46	31
3	NAP	G	1491	31/48	0.90	0.21	26,31,44,48	31
6	GOL	E	1495	6/6	0.90	0.15	45,46,46,47	0
3	NAP	D	1491	31/48	0.90	0.19	24,28,44,46	31
6	GOL	B	1495	6/6	0.90	0.14	37,38,40,42	0
3	NAP	F	1491	31/48	0.90	0.29	26,37,48,51	0
6	GOL	G	1495	6/6	0.91	0.13	35,39,39,40	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	G	1496	6/6	0.92	0.13	35,36,38,39	0
6	GOL	A	1498	6/6	0.92	0.24	35,36,37,37	0
6	GOL	A	1497	6/6	0.92	0.18	23,28,32,32	0
3	NAP	E	1491	31/48	0.93	0.13	13,19,42,44	31
6	GOL	D	1495	6/6	0.93	0.18	36,42,43,45	0
3	NAP	B	1491	31/48	0.93	0.15	19,23,39,41	31
3	NAP	C	1491	31/48	0.94	0.14	19,25,46,50	0
4	BME	A	1492	4/4	0.94	0.17	26,26,28,33	0
4	BME	B	1492	4/4	0.94	0.14	31,35,35,36	0
6	GOL	F	1495	6/6	0.94	0.11	26,27,31,33	0
5	K	H	1494	1/1	0.95	0.13	32,32,32,32	0
5	K	H	1493	1/1	0.97	0.17	32,32,32,32	0
5	K	D	1494	1/1	0.98	0.04	18,18,18,18	0
5	K	G	1494	1/1	0.99	0.10	23,23,23,23	0
5	K	A	1494	1/1	0.99	0.03	17,17,17,17	0
5	K	B	1493	1/1	0.99	0.07	17,17,17,17	0
5	K	B	1494	1/1	0.99	0.03	16,16,16,16	0
5	K	C	1493	1/1	0.99	0.06	22,22,22,22	0
5	K	C	1494	1/1	0.99	0.05	20,20,20,20	0
5	K	D	1493	1/1	0.99	0.08	23,23,23,23	0
5	K	A	1493	1/1	0.99	0.06	20,20,20,20	0
5	K	E	1493	1/1	0.99	0.05	19,19,19,19	0
5	K	F	1493	1/1	0.99	0.09	26,26,26,26	0
5	K	F	1494	1/1	0.99	0.10	21,21,21,21	0
5	K	G	1493	1/1	0.99	0.08	17,17,17,17	0
5	K	E	1494	1/1	1.00	0.07	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAP H 1491:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP A 1491:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP G 1491:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around NAP D 1491:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)











Electron density around NAP B 1491:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.