



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 27, 2020 – 02:14 PM BST

PDB ID : 6WPG
Title : Structural Basis of Salicylic Acid Perception by Arabidopsis NPR Proteins
Authors : Wang, W.; Withers, J.; Li, H.; Zwack, P.J.; Rusnac, D.V.; Shi, H.; Liu, L.; Yan, S.; Hinds, T.R.; Guttman, M.; Dong, X.; Zheng, N.
Deposited on : 2020-04-27
Resolution : 2.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	
1	B	177	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1961 atoms, of which 10 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulatory protein NPR4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	119	981	617	166	185	13	0	0	0
1	B	106	874	552	151	159	12	0	0	0

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-38	PHE	-	expression tag	UNP Q5ICL9
A	-37	CYS	-	expression tag	UNP Q5ICL9
A	-36	LEU	-	expression tag	UNP Q5ICL9
A	-35	THR	-	expression tag	UNP Q5ICL9
A	-34	LEU	-	expression tag	UNP Q5ICL9
A	-33	ARG	-	expression tag	UNP Q5ICL9
A	-32	ARG	-	expression tag	UNP Q5ICL9
A	-31	ARG	-	expression tag	UNP Q5ICL9
A	-30	TYR	-	expression tag	UNP Q5ICL9
A	-29	THR	-	expression tag	UNP Q5ICL9
A	-28	MET	-	expression tag	UNP Q5ICL9
A	-27	GLY	-	expression tag	UNP Q5ICL9
A	-26	SER	-	expression tag	UNP Q5ICL9
A	-25	SER	-	expression tag	UNP Q5ICL9
A	-24	HIS	-	expression tag	UNP Q5ICL9
A	-23	HIS	-	expression tag	UNP Q5ICL9
A	-22	HIS	-	expression tag	UNP Q5ICL9
A	-21	HIS	-	expression tag	UNP Q5ICL9
A	-20	HIS	-	expression tag	UNP Q5ICL9
A	-19	HIS	-	expression tag	UNP Q5ICL9
A	-18	SER	-	expression tag	UNP Q5ICL9
A	-17	SER	-	expression tag	UNP Q5ICL9
A	-16	GLY	-	expression tag	UNP Q5ICL9
A	-15	LEU	-	expression tag	UNP Q5ICL9
A	-14	VAL	-	expression tag	UNP Q5ICL9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	PRO	-	expression tag	UNP Q5ICL9
A	-12	ARG	-	expression tag	UNP Q5ICL9
A	-11	GLY	-	expression tag	UNP Q5ICL9
A	-10	SER	-	expression tag	UNP Q5ICL9
A	-9	HIS	-	expression tag	UNP Q5ICL9
A	-8	SER	-	expression tag	UNP Q5ICL9
A	-7	GLU	-	expression tag	UNP Q5ICL9
A	-6	ASN	-	expression tag	UNP Q5ICL9
A	-5	LEU	-	expression tag	UNP Q5ICL9
A	-4	TYR	-	expression tag	UNP Q5ICL9
A	-3	PHE	-	expression tag	UNP Q5ICL9
A	-2	GLN	-	expression tag	UNP Q5ICL9
A	-1	SER	-	expression tag	UNP Q5ICL9
A	0	MET	-	expression tag	UNP Q5ICL9
A	?	-	LEU	deletion	UNP Q5ICL9
A	?	-	THR	deletion	UNP Q5ICL9
A	?	-	PRO	deletion	UNP Q5ICL9
A	?	-	PRO	deletion	UNP Q5ICL9
A	?	-	PRO	deletion	UNP Q5ICL9
A	?	-	SER	deletion	UNP Q5ICL9
B	334	PHE	-	expression tag	UNP Q5ICL9
B	335	CYS	-	expression tag	UNP Q5ICL9
B	336	LEU	-	expression tag	UNP Q5ICL9
B	337	THR	-	expression tag	UNP Q5ICL9
B	338	LEU	-	expression tag	UNP Q5ICL9
B	339	ARG	-	expression tag	UNP Q5ICL9
B	340	ARG	-	expression tag	UNP Q5ICL9
B	341	ARG	-	expression tag	UNP Q5ICL9
B	342	TYR	-	expression tag	UNP Q5ICL9
B	343	THR	-	expression tag	UNP Q5ICL9
B	344	MET	-	expression tag	UNP Q5ICL9
B	345	GLY	-	expression tag	UNP Q5ICL9
B	346	SER	-	expression tag	UNP Q5ICL9
B	347	SER	-	expression tag	UNP Q5ICL9
B	348	HIS	-	expression tag	UNP Q5ICL9
B	349	HIS	-	expression tag	UNP Q5ICL9
B	350	HIS	-	expression tag	UNP Q5ICL9
B	351	HIS	-	expression tag	UNP Q5ICL9
B	352	HIS	-	expression tag	UNP Q5ICL9
B	353	HIS	-	expression tag	UNP Q5ICL9
B	354	SER	-	expression tag	UNP Q5ICL9
B	355	SER	-	expression tag	UNP Q5ICL9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	356	GLY	-	expression tag	UNP Q5ICL9
B	357	LEU	-	expression tag	UNP Q5ICL9
B	358	VAL	-	expression tag	UNP Q5ICL9
B	359	PRO	-	expression tag	UNP Q5ICL9
B	360	ARG	-	expression tag	UNP Q5ICL9
B	361	GLY	-	expression tag	UNP Q5ICL9
B	362	SER	-	expression tag	UNP Q5ICL9
B	363	HIS	-	expression tag	UNP Q5ICL9
B	364	SER	-	expression tag	UNP Q5ICL9
B	365	GLU	-	expression tag	UNP Q5ICL9
B	366	ASN	-	expression tag	UNP Q5ICL9
B	367	LEU	-	expression tag	UNP Q5ICL9
B	368	TYR	-	expression tag	UNP Q5ICL9
B	369	PHE	-	expression tag	UNP Q5ICL9
B	370	GLN	-	expression tag	UNP Q5ICL9
B	371	SER	-	expression tag	UNP Q5ICL9
B	372	MET	-	expression tag	UNP Q5ICL9
B	?	-	LEU	deletion	UNP Q5ICL9
B	?	-	THR	deletion	UNP Q5ICL9
B	?	-	PRO	deletion	UNP Q5ICL9
B	?	-	PRO	deletion	UNP Q5ICL9
B	?	-	PRO	deletion	UNP Q5ICL9
B	?	-	SER	deletion	UNP Q5ICL9

- Molecule 2 is 2-HYDROXYBENZOIC ACID (three-letter code: SAL) (formula: C₇H₆O₃) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			15	7	5	3		
2	B	1	Total	C	H	O	0	0
			15	7	5	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	42	Total	O	0	0
			42	42		
3	B	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Regulatory protein NPR4



- Molecule 1: Regulatory protein NPR4



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.29Å 88.29Å 138.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.42 – 2.28 44.15 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.42-2.28) 99.7 (44.15-2.28)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.205 , 0.224 0.205 , 0.224	Depositor DCC
R_{free} test set	1430 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	54.4	Xtrriage
Anisotropy	0.638	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 74.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1961	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/997	0.53	0/1337
1	B	0.48	0/888	0.55	0/1190
All	All	0.48	0/1885	0.54	0/2527

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	981	0	973	23	0
1	B	874	0	883	24	0
2	A	10	5	5	0	0
2	B	10	5	5	0	0
3	A	42	0	0	4	0
3	B	34	0	0	5	0
All	All	1951	10	1866	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:493:ARG:HD2	3:A:702:HOH:O	1.62	0.97
1:A:399:PRO:HG3	1:A:497:PRO:HB3	1.47	0.94
1:A:0:MET:SD	3:B:727:HOH:O	2.37	0.81
1:A:389:ARG:HG2	3:A:710:HOH:O	1.89	0.73
1:B:429:ALA:HA	3:B:727:HOH:O	1.91	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	113/177 (64%)	113 (100%)	0	0	100	100
1	B	100/177 (56%)	98 (98%)	2 (2%)	0	100	100
All	All	213/354 (60%)	211 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/162 (68%)	109 (99%)	1 (1%)	78	88
1	B	98/162 (60%)	97 (99%)	1 (1%)	76	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	208/324 (64%)	206 (99%)	2 (1%)	76	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	474	GLN
1	B	504	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SAL	A	601	-	8,10,10	2.36	2 (25%)	9,13,13	0.50	0
2	SAL	B	601	-	8,10,10	2.19	2 (25%)	9,13,13	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAL	A	601	-	-	0/0/4/4	0/1/1/1
2	SAL	B	601	-	-	0/0/4/4	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	SAL	C1-C1'	6.15	1.53	1.47
2	B	601	SAL	C1-C1'	5.67	1.52	1.47
2	A	601	SAL	O2-C2	2.30	1.41	1.36
2	B	601	SAL	O2-C2	2.16	1.40	1.36

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	119/177 (67%)	0.45	7 (5%) 22 27	52, 68, 130, 189	0
1	B	106/177 (59%)	0.67	9 (8%) 10 14	51, 74, 125, 134	0
All	All	225/354 (63%)	0.56	16 (7%) 16 19	51, 69, 127, 189	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	372	MET	5.0
1	B	400	THR	3.8
1	B	371	SER	3.4
1	A	399	PRO	3.1
1	B	374	PRO	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SAL	B	601	10/10	0.98	0.14	46,55,65,66	0
2	SAL	A	601	10/10	0.99	0.16	52,56,68,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.