



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:29 pm BST

PDB ID : 3WQC  
Title : D-threo-3-hydroxyaspartate dehydratase from *Delftia* sp. HT23  
Authors : Yasutake, Y.; Matsumoto, Y.; Wada, M.  
Deposited on : 2014-01-25  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	 % 87% 9% ••
1	B	390	 4% 88% 9% •

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-threo-3-hydroxyaspartate dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	379	2867	1782	535	534	16	0	7	0
1	B	389	2954	1834	556	546	18	0	7	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	EXPRESSION TAG	UNP B2DFG5
A	-8	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-7	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-6	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-5	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-4	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-3	HIS	-	EXPRESSION TAG	UNP B2DFG5
A	-2	ALA	-	EXPRESSION TAG	UNP B2DFG5
A	-1	MET	-	EXPRESSION TAG	UNP B2DFG5
A	0	SER	-	EXPRESSION TAG	UNP B2DFG5
B	-9	GLY	-	EXPRESSION TAG	UNP B2DFG5
B	-8	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-7	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-6	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-5	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-4	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-3	HIS	-	EXPRESSION TAG	UNP B2DFG5
B	-2	ALA	-	EXPRESSION TAG	UNP B2DFG5
B	-1	MET	-	EXPRESSION TAG	UNP B2DFG5
B	0	SER	-	EXPRESSION TAG	UNP B2DFG5

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0
2	B	1	15	8	1	5	1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	3	3	3	0	0
3	A	2	2	2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0
5	A	1	Total Cl 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	333	Total O 333 333	0	0
6	B	317	Total O 317 317	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-threo-3-hydroxyaspartate dehydratase



- Molecule 1: D-threo-3-hydroxyaspartate dehydratase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.69Å 157.69Å 158.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.63 – 1.50 45.63 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.63-1.50) 99.9 (45.63-1.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.95 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.121 , 0.152 0.119 , 0.150	Depositor DCC
$R_{free}$ test set	7908 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.0	Xtriage
Anisotropy	0.159	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 52.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -l,-k,-h 0.003 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6526	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, CL, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.25	7/2935 (0.2%)	1.18	16/3980 (0.4%)
1	B	1.27	12/3025 (0.4%)	1.19	25/4100 (0.6%)
All	All	1.26	19/5960 (0.3%)	1.19	41/8080 (0.5%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	183	GLU	CD-OE1	12.59	1.39	1.25
1	A	18	ASP	CB-CG	10.19	1.73	1.51
1	A	19	ARG	CZ-NH1	6.93	1.42	1.33
1	A	60	SER	CB-OG	-6.90	1.33	1.42
1	B	30	ARG	CZ-NH2	6.71	1.41	1.33
1	B	183	GLU	CD-OE2	6.69	1.33	1.25
1	B	370	SER	CB-OG	-6.59	1.33	1.42
1	B	204	ARG	CD-NE	-6.42	1.35	1.46
1	B	71	GLU	CD-OE1	6.10	1.32	1.25
1	A	375	GLU	CD-OE1	-6.06	1.19	1.25
1	B	374	TRP	CD2-CE2	5.85	1.48	1.41
1	A	287	ARG	CZ-NH2	5.59	1.40	1.33
1	B	147	ASP	CB-CG	-5.45	1.40	1.51
1	B	100[A]	ARG	CZ-NH2	5.20	1.39	1.33
1	B	100[B]	ARG	CZ-NH2	5.20	1.39	1.33
1	A	132	TRP	CD2-CE2	5.18	1.47	1.41
1	A	284	SER	CA-CB	5.16	1.60	1.52
1	B	225	SER	CB-OG	-5.13	1.35	1.42
1	B	18	ASP	CB-CG	5.11	1.62	1.51

All (41) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	ARG	NE-CZ-NH1	20.11	130.36	120.30
1	B	30	ARG	NE-CZ-NH1	-15.70	112.45	120.30
1	A	19	ARG	NE-CZ-NH2	-11.83	114.39	120.30
1	B	204	ARG	NE-CZ-NH2	-10.55	115.02	120.30
1	A	55	ARG	NE-CZ-NH2	9.54	125.07	120.30
1	B	287	ARG	NE-CZ-NH1	-9.04	115.78	120.30
1	A	338	ARG	CG-CD-NE	-8.80	93.32	111.80
1	B	370	SER	N-CA-CB	-8.62	97.58	110.50
1	B	30	ARG	NE-CZ-NH2	8.39	124.49	120.30
1	A	147	ASP	CB-CG-OD1	8.32	125.78	118.30
1	B	204	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	B	147	ASP	CB-CG-OD2	-8.10	111.01	118.30
1	B	231	THR	OG1-CB-CG2	-7.72	92.24	110.00
1	B	18	ASP	CB-CG-OD1	7.27	124.84	118.30
1	B	16	ASP	CB-CG-OD1	7.16	124.74	118.30
1	B	193	ARG	NE-CZ-NH2	-6.85	116.88	120.30
1	B	106	LYS	CD-CE-NZ	-6.77	96.14	111.70
1	A	287	ARG	NE-CZ-NH1	-6.55	117.03	120.30
1	A	16	ASP	CB-CG-OD2	6.50	124.15	118.30
1	B	83	TYR	CB-CG-CD2	-6.26	117.25	121.00
1	B	153	LEU	CB-CG-CD1	6.22	121.57	111.00
1	A	16	ASP	CB-CG-OD1	6.11	123.80	118.30
1	A	16	ASP	OD1-CG-OD2	-5.94	112.02	123.30
1	A	338	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	B	22	ARG	CG-CD-NE	-5.66	99.92	111.80
1	A	147	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	287	ARG	NE-CZ-NH2	5.55	123.07	120.30
1	B	226[A]	ARG	NE-CZ-NH2	5.53	123.06	120.30
1	B	226[B]	ARG	NE-CZ-NH2	5.53	123.06	120.30
1	B	101	ARG	CG-CD-NE	5.45	123.25	111.80
1	B	83	TYR	CB-CG-CD1	5.42	124.25	121.00
1	B	377	LEU	CB-CG-CD2	-5.42	101.79	111.00
1	B	338	ARG	CG-CD-NE	-5.34	100.58	111.80
1	A	344	ARG	CG-CD-NE	-5.29	100.69	111.80
1	B	344	ARG	CG-CD-NE	-5.25	100.78	111.80
1	A	285	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	A	376	ARG	NE-CZ-NH2	5.15	122.88	120.30
1	B	37	ARG	NE-CZ-NH2	5.14	122.87	120.30
1	A	338	ARG	CD-NE-CZ	5.03	130.64	123.60
1	B	226[A]	ARG	NE-CZ-NH1	-5.01	117.80	120.30
1	B	226[B]	ARG	NE-CZ-NH1	-5.01	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2867	0	2876	19	0
1	B	2954	0	2946	27	0
2	A	15	0	6	0	0
2	B	15	0	6	0	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
4	A	12	0	15	3	0
4	B	6	0	8	0	0
5	A	1	0	0	0	0
5	B	1	0	0	1	0
6	A	333	0	0	7	0
6	B	317	0	0	4	0
All	All	6526	0	5857	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:ARG:HH11	1:A:30:ARG:HG3	1.08	1.16
1:B:100[B]:ARG:NH1	1:B:100[B]:ARG:HG2	1.63	1.05
1:B:100[B]:ARG:CG	1:B:100[B]:ARG:HH11	1.66	1.03
1:B:100[B]:ARG:HH11	1:B:100[B]:ARG:HG2	0.86	1.02
1:A:30:ARG:NH1	1:A:30:ARG:HG3	1.81	0.88
1:B:291:ARG:HD2	1:B:291:ARG:H	1.37	0.88
1:B:18:ASP:HB3	1:B:22:ARG:NH2	1.90	0.86
1:B:68[B]:LYS:HD3	6:B:810:HOH:O	1.78	0.84
1:A:231[B]:THR:HG22	6:A:798:HOH:O	1.84	0.78
1:A:338:ARG:HD2	6:A:652:HOH:O	1.87	0.75
1:A:30:ARG:CG	1:A:30:ARG:HH11	1.95	0.74
1:B:100[B]:ARG:CG	1:B:100[B]:ARG:NH1	2.36	0.72

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231[B]:THR:CG2	6:A:798:HOH:O	2.40	0.69
1:B:242:ASP:OD1	1:B:245[B]:MET:HG2	1.94	0.68
1:A:51:ALA:HA	1:A:54:GLN:HE21	1.59	0.68
1:B:291:ARG:H	1:B:291:ARG:CD	2.08	0.67
1:B:51:ALA:HA	1:B:54:GLN:HE21	1.62	0.63
1:A:242:ASP:OD1	1:A:245[B]:MET:HG2	2.04	0.58
1:A:43:LYS:NZ	6:A:698:HOH:O	2.38	0.56
4:A:404:GOL:H11	5:B:401:CL:CL	2.43	0.56
1:B:68[B]:LYS:CD	6:B:810:HOH:O	2.44	0.56
1:B:18:ASP:HB3	1:B:22:ARG:HH22	1.69	0.55
1:B:368:ASP:OD1	1:B:370:SER:CB	2.58	0.52
1:A:326:ARG:HD2	1:A:332:GLU:OE1	2.11	0.50
1:B:245[B]:MET:HA	1:B:245[B]:MET:HE3	1.93	0.49
1:B:368:ASP:OD1	1:B:370:SER:HB2	2.14	0.47
4:A:404:GOL:H12	6:A:560:HOH:O	2.15	0.46
1:B:100[B]:ARG:NH1	6:B:799:HOH:O	2.49	0.45
1:A:92:LEU:HD21	1:A:118:ILE:HG12	1.98	0.45
1:A:68:LYS:HG2	1:A:377:LEU:HD21	2.00	0.44
1:A:291:ARG:H	1:A:291:ARG:HG2	1.58	0.44
1:A:291:ARG:HH11	1:A:291:ARG:HG3	1.82	0.44
1:B:368:ASP:OD1	1:B:370:SER:HB3	2.17	0.44
1:A:180:ASP:OD2	1:A:293:LYS:HE3	2.18	0.44
1:B:288:GLY:O	1:B:291:ARG:HD2	2.18	0.44
4:A:403:GOL:C1	6:A:556:HOH:O	2.66	0.44
1:B:219:PRO:HG3	1:B:245[B]:MET:HE1	2.00	0.43
1:A:11:PRO:HA	1:A:261:LEU:O	2.20	0.42
1:B:72:GLN:OE1	1:B:377:LEU:HD13	2.19	0.42
1:B:67:LEU:HD23	1:B:67:LEU:HA	1.93	0.41
1:B:132:TRP:CH2	1:B:212:VAL:HG11	2.56	0.41
1:A:324:LEU:C	1:A:324:LEU:HD23	2.41	0.41
1:B:338:ARG:HD3	1:B:338:ARG:HA	1.88	0.41
1:A:94[A]:GLN:HG3	6:A:637:HOH:O	2.20	0.41
1:B:326:ARG:HD3	1:B:330:ALA:HB3	2.03	0.41
1:B:100[A]:ARG:HD2	6:B:817:HOH:O	2.21	0.41
1:B:275:ILE:HD12	1:B:323:ILE:CD1	2.51	0.41
1:A:28:GLN:HG2	1:A:57:ALA:O	2.21	0.40
1:B:324:LEU:C	1:B:324:LEU:HD23	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/390 (98%)	376 (98%)	6 (2%)	2 (0%)	29	9
1	B	394/390 (101%)	385 (98%)	8 (2%)	1 (0%)	41	18
All	All	778/780 (100%)	761 (98%)	14 (2%)	3 (0%)	34	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	B	141	ARG
1	A	236	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/289 (99%)	284 (99%)	3 (1%)	76	57
1	B	296/289 (102%)	293 (99%)	3 (1%)	76	57
All	All	583/578 (101%)	577 (99%)	6 (1%)	73	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	30	ARG
1	A	291	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	153	LEU
1	B	291	ARG
1	B	334	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	B	54	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PLP	A	401	1	15,15,16	2.32	5 (33%)	20,22,23	2.22	8 (40%)
4	GOL	A	403	-	5,5,5	0.75	0	5,5,5	2.45	3 (60%)
4	GOL	A	404	-	5,5,5	0.36	0	5,5,5	1.68	2 (40%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PLP	B	402	1	15,15,16	1.94	4 (26%)	20,22,23	1.38	4 (20%)
4	GOL	B	404	-	5,5,5	0.72	0	5,5,5	1.71	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	401	1	-	0/6/6/8	0/1/1/1
4	GOL	A	403	-	-	0/4/4/4	-
4	GOL	A	404	-	-	2/4/4/4	-
2	PLP	B	402	1	-	0/6/6/8	0/1/1/1
4	GOL	B	404	-	-	1/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PLP	C5-C4	4.96	1.46	1.40
2	A	401	PLP	C3-C2	4.72	1.45	1.40
2	B	402	PLP	C4A-C4	-4.14	1.43	1.51
2	B	402	PLP	C3-C2	3.88	1.44	1.40
2	A	401	PLP	C4A-C4	-2.77	1.45	1.51
2	A	401	PLP	O3-C3	-2.75	1.30	1.37
2	A	401	PLP	C6-C5	2.50	1.42	1.37
2	B	402	PLP	C5-C4	2.37	1.43	1.40
2	B	402	PLP	C3-C4	2.08	1.44	1.40

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PLP	C5-C6-N1	-4.12	116.96	123.82
2	A	401	PLP	C6-N1-C2	3.98	126.54	119.17
2	A	401	PLP	C4A-C4-C5	3.86	124.91	120.94
2	A	401	PLP	C6-C5-C4	3.72	121.09	118.16
4	A	403	GOL	C3-C2-C1	3.22	124.23	111.70
4	A	403	GOL	O3-C3-C2	-3.20	94.88	110.20
2	A	401	PLP	C2A-C2-N1	3.16	123.84	117.67
2	B	402	PLP	C6-N1-C2	2.90	124.54	119.17
2	A	401	PLP	C2A-C2-C3	-2.81	117.42	120.89
4	A	403	GOL	O2-C2-C3	-2.79	96.85	109.12

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PLP	O3P-P-O2P	2.42	116.87	107.64
2	B	402	PLP	O3P-P-O2P	2.24	116.21	107.64
4	A	404	GOL	O2-C2-C3	-2.24	99.25	109.12
4	A	404	GOL	O1-C1-C2	2.21	120.79	110.20
4	B	404	GOL	O3-C3-C2	-2.17	99.79	110.20
4	B	404	GOL	C3-C2-C1	-2.15	103.34	111.70
2	B	402	PLP	C5-C6-N1	-2.12	120.29	123.82
2	A	401	PLP	C4A-C4-C3	-2.04	117.03	120.50
2	B	402	PLP	O3-C3-C4	2.00	123.38	118.10

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	GOL	C1-C2-C3-O3
4	A	404	GOL	O2-C2-C3-O3
4	B	404	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	GOL	1	0
4	A	404	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	379/390 (97%)	-0.48	2 (0%) 91 93	9, 17, 37, 68	0
1	B	389/390 (99%)	-0.34	14 (3%) 42 47	10, 17, 43, 72	0
All	All	768/780 (98%)	-0.41	16 (2%) 63 68	9, 17, 38, 72	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-8	HIS	5.4
1	B	-5	HIS	4.7
1	B	-4	HIS	4.4
1	B	331	ALA	4.4
1	A	2	GLN	4.2
1	B	-2	ALA	3.7
1	B	-1	MET	3.7
1	B	-7	HIS	3.4
1	B	330	ALA	3.3
1	B	-6	HIS	3.1
1	B	333	ALA	2.9
1	B	0	SER	2.8
1	B	334	ASP	2.5
1	B	1	MET	2.3
1	B	-3	HIS	2.3
1	A	147	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	A	406	1/1	0.80	0.16	28,28,28,28	1
4	GOL	A	403	6/6	0.93	0.12	28,44,48,52	0
4	GOL	A	404	6/6	0.95	0.13	26,35,45,54	0
4	GOL	B	404	6/6	0.95	0.09	31,36,44,49	0
3	MG	B	406	1/1	0.98	0.25	21,21,21,21	1
2	PLP	B	402	15/16	0.99	0.04	11,13,19,21	0
2	PLP	A	401	15/16	0.99	0.06	12,15,19,21	0
5	CL	A	405	1/1	1.00	0.05	19,19,19,19	1
3	MG	B	405	1/1	1.00	0.20	37,37,37,37	0
3	MG	A	402	1/1	1.00	0.07	14,14,14,14	0
5	CL	B	401	1/1	1.00	0.08	19,19,19,19	1
3	MG	B	403	1/1	1.00	0.09	14,14,14,14	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.