



Full wwPDB EM Validation Report ⓘ

Apr 9, 2024 – 12:12 PM JST

PDB ID : 8WRL
EMDB ID : EMD-37779
Title : XBB.1.5 RBD in complex with ACE2
Authors : Feng, L.L.; Feng, L.L.
Deposited on : 2023-10-15
Resolution : 3.36 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	594	
2	B	238	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Processed angiotensin-converting enzyme 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	594	4845	3099	803	914	29	0	0

- Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	196	1558	1006	262	282	8	0	0

There are 41 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	300	MET	-	initiating methionine	UNP P0DTC2
B	301	PRO	-	expression tag	UNP P0DTC2
B	302	LEU	-	expression tag	UNP P0DTC2
B	303	LEU	-	expression tag	UNP P0DTC2
B	304	LEU	-	expression tag	UNP P0DTC2
B	305	LEU	-	expression tag	UNP P0DTC2
B	306	LEU	-	expression tag	UNP P0DTC2
B	307	PRO	-	expression tag	UNP P0DTC2
B	308	LEU	-	expression tag	UNP P0DTC2
B	309	LEU	-	expression tag	UNP P0DTC2
B	310	TRP	-	expression tag	UNP P0DTC2
B	311	ALA	-	expression tag	UNP P0DTC2
B	312	GLY	-	expression tag	UNP P0DTC2
B	313	ALA	-	expression tag	UNP P0DTC2
B	314	LEU	-	expression tag	UNP P0DTC2
B	315	ALA	-	expression tag	UNP P0DTC2
B	316	MET	-	expression tag	UNP P0DTC2
B	317	ALA	-	expression tag	UNP P0DTC2
B	318	ALA	-	expression tag	UNP P0DTC2
B	339	HIS	GLY	variant	UNP P0DTC2
B	346	THR	ARG	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	368	ILE	LEU	variant	UNP P0DTC2
B	371	PHE	SER	variant	UNP P0DTC2
B	373	PRO	SER	variant	UNP P0DTC2
B	375	PHE	SER	variant	UNP P0DTC2
B	376	ALA	THR	variant	UNP P0DTC2
B	405	ASN	ASP	variant	UNP P0DTC2
B	408	SER	ARG	variant	UNP P0DTC2
B	417	ASN	LYS	variant	UNP P0DTC2
B	440	LYS	ASN	variant	UNP P0DTC2
B	445	PRO	VAL	variant	UNP P0DTC2
B	446	SER	GLY	variant	UNP P0DTC2
B	460	LYS	ASN	variant	UNP P0DTC2
B	477	ASN	SER	variant	UNP P0DTC2
B	478	LYS	THR	variant	UNP P0DTC2
B	484	ALA	GLU	variant	UNP P0DTC2
B	486	PRO	PHE	variant	UNP P0DTC2
B	490	SER	PHE	variant	UNP P0DTC2
B	498	ARG	GLN	variant	UNP P0DTC2
B	501	TYR	ASN	variant	UNP P0DTC2
B	505	HIS	TYR	variant	UNP P0DTC2

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf	
			Total	C	N		O
3	A	1	14	8	1	5	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
3	A	1	Total 14	8	1	5	0
3	A	1	Total 14	8	1	5	0
3	A	1	Total 14	8	1	5	0
3	B	1	Total 14	8	1	5	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Processed angiotensin-converting enzyme 2



- Molecule 2: Spike protein S1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	226753	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	52	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	4.343	Depositor
Minimum map value	-3.445	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.083	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	206.848, 206.848, 206.848	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.808, 0.808, 0.808	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	1/4981 (0.0%)	0.57	1/6767 (0.0%)
2	B	0.31	0/1607	0.56	0/2188
All	All	0.34	1/6588 (0.0%)	0.56	1/8955 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	331	SER	CA-CB	-5.24	1.45	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	497	TYR	CB-CA-C	-5.16	100.08	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	169	ARG	Sidechain
1	A	357	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4845	0	4621	91	0
2	B	1558	0	1493	15	0
3	A	56	0	52	0	0
3	B	14	0	13	0	0
All	All	6473	0	6179	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:TRP:HE1	1:A:502:SER:HB2	1.36	0.90
1:A:472:GLN:HG2	1:A:475:LYS:HD2	1.68	0.76
1:A:379:ILE:O	1:A:383:MET:HG3	1.92	0.70
1:A:446:ILE:HG21	1:A:523:PHE:HE2	1.61	0.65
1:A:474:MET:HE3	1:A:498:CYS:HA	1.80	0.63
1:A:433:GLU:O	1:A:436:ILE:HG22	1.99	0.63
1:A:83:TYR:O	1:A:101:GLN:NE2	2.34	0.60
1:A:293:VAL:HG12	1:A:366:MET:HE2	1.83	0.60
1:A:398:GLU:HG3	1:A:514:ARG:HG2	1.84	0.59
2:B:454:ARG:NH2	2:B:469:SER:O	2.36	0.59
1:A:402:GLU:HB2	1:A:518:ARG:HG3	1.85	0.59
2:B:457:ARG:NH1	2:B:467:ASP:OD2	2.36	0.58
1:A:184:VAL:HG22	1:A:464:PHE:HE1	1.69	0.57
1:A:243:TYR:CZ	1:A:247:LYS:HD2	2.40	0.56
1:A:591:LEU:O	1:A:592:PHE:C	2.45	0.56
1:A:81:GLN:NE2	1:A:101:GLN:O	2.39	0.55
1:A:308:PHE:O	1:A:328:TRP:HH2	1.89	0.54
2:B:411:ALA:HB3	2:B:414:GLN:HG3	1.89	0.54
1:A:169:ARG:HH22	1:A:271:TRP:HD1	1.55	0.54
1:A:491:VAL:HB	1:A:493:HIS:HE1	1.73	0.53
1:A:326:GLY:O	1:A:330:ASN:ND2	2.40	0.53
1:A:491:VAL:HB	1:A:493:HIS:CE1	2.43	0.53
1:A:245:ARG:HG3	1:A:256:ILE:HD11	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:358:ILE:HD11	1:A:379:ILE:HG21	1.90	0.52
1:A:284:PRO:HG2	1:A:437:ASN:HA	1.92	0.51
1:A:173:GLY:HA2	1:A:176:LEU:HD12	1.93	0.51
1:A:410:LEU:HD21	1:A:442:GLN:NE2	2.26	0.51
1:A:439:LEU:CD1	1:A:590:PRO:HB2	2.40	0.51
1:A:462:MET:HE2	1:A:468:ILE:HD11	1.93	0.51
1:A:243:TYR:O	1:A:247:LYS:HG3	2.11	0.50
1:A:524:GLN:HG2	1:A:583:PRO:HG2	1.92	0.50
1:A:201:ASP:OD1	1:A:204:ARG:NH2	2.43	0.50
1:A:152:MET:O	1:A:161:ARG:NH2	2.45	0.50
1:A:404:VAL:O	1:A:408:MET:HG2	2.13	0.49
1:A:474:MET:SD	1:A:474:MET:N	2.82	0.49
1:A:397:ASN:HD21	1:A:521:TYR:HE2	1.61	0.49
1:A:520:LEU:HD23	1:A:579:MET:HB3	1.95	0.48
1:A:307:ILE:HG23	1:A:369:PHE:HD1	1.78	0.48
1:A:468:ILE:HG22	1:A:473:TRP:HD1	1.79	0.48
1:A:476:LYS:O	1:A:477:TRP:C	2.52	0.48
2:B:422:ASN:HD21	2:B:454:ARG:H	1.60	0.48
1:A:327:PHE:CD2	1:A:327:PHE:C	2.86	0.48
1:A:49:ASN:HA	1:A:52:THR:HG22	1.96	0.48
1:A:459:TRP:CH2	1:A:500:PRO:HG2	2.48	0.48
1:A:285:PHE:HE2	1:A:436:ILE:HG21	1.79	0.47
1:A:477:TRP:CZ2	1:A:500:PRO:HA	2.49	0.47
1:A:501:ALA:O	1:A:502:SER:CB	2.62	0.47
1:A:346:PRO:HB2	1:A:375:GLU:HG3	1.96	0.47
2:B:420:ASP:OD2	2:B:460:LYS:NZ	2.46	0.47
1:A:252:TYR:CZ	1:A:266:LEU:HD11	2.49	0.47
1:A:586:ASN:HA	1:A:589:GLU:HB2	1.97	0.47
1:A:529:LEU:HD11	1:A:554:LEU:HD13	1.96	0.46
1:A:302:TRP:CH2	1:A:310:GLU:HG3	2.50	0.46
1:A:399:GLY:HA3	1:A:517:THR:HG22	1.97	0.46
1:A:499:ASP:O	1:A:500:PRO:C	2.53	0.46
1:A:439:LEU:HD12	1:A:590:PRO:HB2	1.97	0.46
1:A:327:PHE:CE1	1:A:358:ILE:HG13	2.51	0.46
1:A:204:ARG:HE	1:A:219:ARG:HB3	1.81	0.46
2:B:422:ASN:ND2	2:B:454:ARG:H	2.14	0.45
1:A:284:PRO:HB3	1:A:594:TRP:CH2	2.51	0.45
1:A:586:ASN:OD1	1:A:587:TYR:N	2.49	0.45
1:A:494:ASP:OD1	1:A:494:ASP:N	2.38	0.45
1:A:115:ARG:NH1	1:A:118:THR:OG1	2.50	0.45
1:A:351:LEU:HD11	1:A:357:ARG:HD2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:ASP:OD2	1:A:219:ARG:NH1	2.49	0.45
1:A:45:LEU:O	1:A:49:ASN:ND2	2.50	0.45
1:A:363:LYS:HE3	1:A:363:LYS:HB3	1.82	0.45
1:A:327:PHE:CD2	1:A:327:PHE:O	2.70	0.45
1:A:108:LEU:HD22	1:A:112:LYS:HE3	1.99	0.44
1:A:188:ASN:OD1	1:A:192:ARG:NH1	2.50	0.44
1:A:293:VAL:CG1	1:A:366:MET:HE2	2.48	0.44
1:A:498:CYS:C	1:A:500:PRO:HD2	2.38	0.43
1:A:293:VAL:HG12	1:A:366:MET:HB2	2.00	0.43
1:A:468:ILE:CD1	1:A:476:LYS:HG2	2.48	0.43
2:B:461:LEU:HD22	2:B:465:GLU:HB3	2.00	0.43
1:A:201:ASP:O	1:A:219:ARG:NH1	2.52	0.43
2:B:422:ASN:OD1	2:B:422:ASN:N	2.50	0.43
1:A:195:HIS:N	1:A:195:HIS:ND1	2.67	0.43
2:B:442:ASP:OD2	2:B:509:ARG:NE	2.48	0.43
1:A:530:CYS:HB2	1:A:542:CYS:HB3	1.63	0.42
1:A:31:LYS:HD3	2:B:456:PHE:HE2	1.84	0.42
1:A:21:ILE:HA	1:A:24:GLN:HG2	2.01	0.42
1:A:601:ASN:OD1	1:A:601:ASN:N	2.49	0.42
1:A:589:GLU:HB3	1:A:590:PRO:HD3	2.02	0.42
2:B:432:CYS:HB2	2:B:513:LEU:HB2	2.02	0.42
2:B:433:VAL:O	2:B:433:VAL:HG12	2.20	0.42
1:A:47:SER:O	1:A:51:ASN:ND2	2.53	0.41
1:A:248:LEU:HD12	1:A:262:LEU:HD12	2.01	0.41
1:A:148:LEU:HD23	1:A:151:ILE:HD11	2.02	0.41
1:A:519:THR:HA	1:A:522:GLN:HG2	2.02	0.41
2:B:341:VAL:HG11	2:B:397:ALA:HB1	2.01	0.41
1:A:232:GLU:HB3	1:A:581:VAL:HG21	2.03	0.41
1:A:376:MET:O	1:A:379:ILE:HG22	2.19	0.41
1:A:144:LEU:HA	1:A:148:LEU:HB2	2.01	0.41
2:B:374:PHE:HD1	2:B:436:TRP:HB3	1.85	0.41
1:A:29:LEU:HD21	1:A:97:LEU:HG	2.03	0.41
1:A:334:THR:HG22	1:A:335:ASP:OD1	2.20	0.41
1:A:552:GLN:NE2	1:A:556:ASN:OD1	2.53	0.41
1:A:227:GLU:O	1:A:231:GLU:HG2	2.21	0.41
2:B:519:HIS:O	2:B:519:HIS:CG	2.74	0.41
1:A:356:PHE:HB3	1:A:379:ILE:HD11	2.03	0.40
1:A:458:LYS:HZ1	1:A:484:ILE:HD11	1.86	0.40
1:A:184:VAL:HG21	1:A:473:TRP:CH2	2.56	0.40
1:A:287:GLN:H	1:A:287:GLN:HG3	1.60	0.40
1:A:118:THR:O	1:A:122:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	592/594 (100%)	558 (94%)	31 (5%)	3 (0%)	29	63
2	B	194/238 (82%)	171 (88%)	23 (12%)	0	100	100
All	All	786/832 (94%)	729 (93%)	54 (7%)	3 (0%)	38	68

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	502	SER
1	A	360	MET
1	A	496	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	525/525 (100%)	505 (96%)	20 (4%)	33	63
2	B	169/205 (82%)	163 (96%)	6 (4%)	35	64
All	All	694/730 (95%)	668 (96%)	26 (4%)	37	63

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	177	ARG
1	A	195	HIS
1	A	201	ASP
1	A	217	TYR
1	A	287	GLN
1	A	306	ARG
1	A	327	PHE
1	A	347	THR
1	A	381	TYR
1	A	431	ASP
1	A	473	TRP
1	A	474	MET
1	A	477	TRP
1	A	494	ASP
1	A	500	PRO
1	A	530	CYS
1	A	547	SER
1	A	571	GLU
1	A	580	ASN
1	A	587	TYR
2	B	377	PHE
2	B	422	ASN
2	B	443	SER
2	B	453	TYR
2	B	489	TYR
2	B	495	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	422	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	601	2	14,14,15	0.25	0	17,19,21	0.45	0
3	NAG	A	701	1	14,14,15	0.26	0	17,19,21	0.48	0
3	NAG	A	703	1	14,14,15	0.41	0	17,19,21	0.38	0
3	NAG	A	704	1	14,14,15	0.36	0	17,19,21	0.36	0
3	NAG	A	702	1	14,14,15	0.32	0	17,19,21	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	601	2	-	1/6/23/26	0/1/1/1
3	NAG	A	701	1	-	0/6/23/26	0/1/1/1
3	NAG	A	703	1	-	2/6/23/26	0/1/1/1
3	NAG	A	704	1	-	2/6/23/26	0/1/1/1
3	NAG	A	702	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.



All (7) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	702	NAG	O5-C5-C6-O6
3	A	704	NAG	O5-C5-C6-O6
3	A	702	NAG	C4-C5-C6-O6
3	A	704	NAG	C4-C5-C6-O6
3	A	703	NAG	C4-C5-C6-O6
3	B	601	NAG	O5-C5-C6-O6
3	A	703	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-37779. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)

6.1.1 Primary map




6.1.2 Raw map




The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 136




Y Index: 112




Z Index: 115


6.3.2 Raw map



X Index: 134



Y Index: 134




Z Index: 159


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 50 nm³; this corresponds to an approximate mass of 45 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.298 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.298 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.36	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.72	4.00	3.78

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.72 differs from the reported value 3.36 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-37779 and PDB model 8WRL. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).





9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7210	 0.4520
A	 0.7020	 0.4390
B	 0.7790	 0.4920

