



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 03:25 PM BST

PDB ID : 6WT4
Title : Structure of a bacterial STING receptor from Flavobacteriaceae sp. in complex with 3',3'-cGAMP
Authors : Morehouse, B.R.; Govande, A.A.; Millman, A.; Keszei, A.F.A.; Lowey, B.; Ofir, G.; Shao, S.; Sorek, R.; Kranzusch, P.J.
Deposited on : 2020-05-01
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	
1	B	162	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2563 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacterial STING.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	147	Total	C	N	O	S	0	1	0
			1211	787	198	223	3			
1	B	140	Total	C	N	O	S	0	1	0
			1154	751	189	211	3			

- Molecule 2 is 2-amino-9-[(2R,3R,3aS,5R,7aR,9R,10R,10aS,12R,14aR)-9-(6-amino-9H-purin-9-yl)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecin-2-yl]-1,9-dihydro-6H-purin-6-one (three-letter code: 4BW) (formula: C₂₀H₂₄N₁₀O₁₃P₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	1
			90	40	20	26	4		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	57	Total	O	0	0
			57	57		
4	B	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bacterial STING



- Molecule 1: Bacterial STING



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	121.85Å 121.85Å 48.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.08 – 1.78 44.79 – 1.78	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.08-1.78) 99.6 (44.79-1.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 1.78Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.199 , 0.230 0.201 , 0.235	Depositor DCC
R_{free} test set	2005 reflections (5.90%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtrriage
Anisotropy	0.251	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2563	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, 4BW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1233	0.44	0/1659
1	B	0.27	0/1176	0.43	0/1584
All	All	0.27	0/2409	0.44	0/3243

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1211	0	1242	12	0
1	B	1154	0	1179	9	0
2	A	90	0	43	5	0
3	B	10	0	0	0	0
4	A	57	0	0	2	0
4	B	41	0	0	0	0
All	All	2563	0	2464	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301[A]:4BW:CBO	2:A:301[A]:4BW:OAX	1.67	1.07
2:A:301[B]:4BW:CBO	2:A:301[B]:4BW:OAX	1.67	1.05
1:A:153[A]:ARG:HH22	1:B:153[A]:ARG:HH22	1.33	0.73
2:A:301[B]:4BW:H23	1:B:153[B]:ARG:NH2	1.87	0.72
1:A:230:LYS:NZ	4:A:401:HOH:O	2.23	0.71
1:B:97:ILE:HD12	1:B:100:ILE:HD11	1.81	0.62
1:A:153[A]:ARG:HH22	1:B:153[A]:ARG:NH2	1.99	0.59
1:A:153[A]:ARG:NH2	1:B:153[A]:ARG:HH22	2.01	0.55
1:A:194:GLU:HA	1:A:197:LYS:HE2	1.92	0.51
2:A:301[B]:4BW:H23	1:B:153[B]:ARG:HH22	1.55	0.50
1:A:114:LYS:HZ3	1:A:116:LYS:HE3	1.81	0.46
1:A:193:SER:HB3	1:A:196:LYS:HG2	2.00	0.43
1:A:116:LYS:NZ	4:A:405:HOH:O	2.52	0.42
1:B:113:TYR:HB2	1:B:168:PHE:CZ	2.54	0.42
1:A:199:LYS:HD2	1:A:199:LYS:HA	1.82	0.42
1:B:199:LYS:HB2	1:B:199:LYS:HE2	1.84	0.41
1:A:121:LEU:HD11	1:A:205:ILE:HD11	2.03	0.41
1:A:175:LEU:HD13	1:B:151:ARG:HA	2.03	0.41
1:A:153[B]:ARG:NH2	2:A:301[B]:4BW:OAC	2.47	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/162 (89%)	141 (98%)	3 (2%)	0	100	100
1	B	137/162 (85%)	133 (97%)	4 (3%)	0	100	100
All	All	281/324 (87%)	274 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/149 (92%)	133 (97%)	4 (3%)	42	25
1	B	130/149 (87%)	127 (98%)	3 (2%)	50	34
All	All	267/298 (90%)	260 (97%)	7 (3%)	46	29

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	151	ARG
1	A	170	PHE
1	A	206	GLU
1	B	111	CYS
1	B	116	LYS
1	B	157	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	301	-	4,4,4	0.13	0	6,6,6	0.11	0
3	SO4	B	302	-	4,4,4	0.14	0	6,6,6	0.07	0
2	4BW	A	301[B]	-	44,51,51	5.26	19 (43%)	52,80,80	1.68	10 (19%)
2	4BW	A	301[A]	-	44,51,51	5.28	19 (43%)	52,80,80	1.67	10 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4BW	A	301[B]	-	-	0/22/62/62	0/6/7/7
2	4BW	A	301[A]	-	-	1/22/62/62	0/6/7/7

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301[A]	4BW	OAX-CBO	19.13	1.67	1.41
2	A	301[B]	4BW	OAX-CBO	18.78	1.67	1.41
2	A	301[A]	4BW	CBI-CBO	-17.66	1.27	1.53
2	A	301[B]	4BW	CBI-CBO	-17.54	1.27	1.53
2	A	301[A]	4BW	C2'-C3'	-12.77	1.24	1.52
2	A	301[B]	4BW	C2'-C3'	-12.72	1.24	1.52
2	A	301[A]	4BW	O4'-C4'	-7.68	1.27	1.45
2	A	301[B]	4BW	O4'-C4'	-7.59	1.28	1.45
2	A	301[B]	4BW	O4'-C1'	7.50	1.51	1.41
2	A	301[A]	4BW	O4'-C1'	6.84	1.50	1.41
2	A	301[B]	4BW	CBG-NAS	6.31	1.45	1.35
2	A	301[B]	4BW	OAX-CBK	-6.27	1.31	1.45
2	A	301[A]	4BW	CBG-NAS	6.15	1.45	1.35
2	A	301[A]	4BW	OAX-CBK	-6.04	1.31	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301[A]	4BW	CBC-CBE	5.64	1.51	1.41
2	A	301[A]	4BW	CBA-NAA	5.60	1.45	1.33
2	A	301[B]	4BW	CBA-NAA	5.53	1.45	1.33
2	A	301[B]	4BW	CBC-CBE	5.45	1.50	1.41
2	A	301[B]	4BW	CBA-NAT	5.39	1.45	1.35
2	A	301[A]	4BW	CBA-NAT	5.26	1.44	1.35
2	A	301[B]	4BW	CBC-NAT	4.77	1.41	1.33
2	A	301[A]	4BW	CBC-NAT	4.75	1.41	1.33
2	A	301[A]	4BW	O2'-C2'	3.98	1.52	1.43
2	A	301[A]	4BW	C3'-C4'	3.90	1.63	1.52
2	A	301[B]	4BW	C3'-C4'	3.87	1.63	1.52
2	A	301[B]	4BW	O2'-C2'	3.78	1.51	1.43
2	A	301[B]	4BW	OAZ-CBM	-3.43	1.31	1.44
2	A	301[A]	4BW	CBI-CBM	3.43	1.60	1.52
2	A	301[A]	4BW	OAZ-CBM	-3.23	1.32	1.44
2	A	301[B]	4BW	CBI-CBM	3.16	1.60	1.52
2	A	301[A]	4BW	C6-N6	2.80	1.44	1.34
2	A	301[B]	4BW	C6-N6	2.65	1.43	1.34
2	A	301[B]	4BW	OAG-CBI	2.50	1.48	1.43
2	A	301[A]	4BW	OAG-CBI	2.33	1.48	1.43
2	A	301[B]	4BW	CBA-NAS	2.25	1.45	1.34
2	A	301[A]	4BW	CBA-NAS	2.25	1.45	1.34
2	A	301[A]	4BW	PBS-OAZ	-2.23	1.54	1.60
2	A	301[B]	4BW	PBS-OAZ	-2.03	1.55	1.60

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301[B]	4BW	NAS-CBA-NAT	-5.50	119.88	127.22
2	A	301[A]	4BW	NAS-CBA-NAT	-5.38	120.05	127.22
2	A	301[A]	4BW	N3-C2-N1	-4.83	121.13	128.68
2	A	301[B]	4BW	N3-C2-N1	-4.75	121.25	128.68
2	A	301[B]	4BW	CBA-NAS-CBG	4.30	120.27	115.36
2	A	301[A]	4BW	CBA-NAS-CBG	3.90	119.82	115.36
2	A	301[B]	4BW	C4-C5-N7	-3.04	106.23	109.40
2	A	301[A]	4BW	CBE-CBC-NAT	-2.69	119.75	123.43
2	A	301[A]	4BW	CBG-CBE-NAR	-2.65	106.64	109.40
2	A	301[A]	4BW	CBC-NAT-CBA	2.62	120.09	115.93
2	A	301[A]	4BW	C4-C5-N7	-2.48	106.82	109.40
2	A	301[B]	4BW	CBC-NAT-CBA	2.43	119.78	115.93
2	A	301[B]	4BW	CBE-CBC-NAT	-2.40	120.14	123.43
2	A	301[B]	4BW	CBG-CBE-NAR	-2.35	106.95	109.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301[B]	4BW	C2-N1-C6	2.33	122.75	118.75
2	A	301[A]	4BW	C2-N1-C6	2.31	122.71	118.75
2	A	301[A]	4BW	NAA-CBA-NAT	2.17	120.62	117.25
2	A	301[A]	4BW	CBM-CBI-CBO	2.14	104.63	99.89
2	A	301[B]	4BW	NAA-CBA-NAT	2.08	120.48	117.25
2	A	301[B]	4BW	CBM-CBI-CBO	2.05	104.43	99.89

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301[A]	4BW	C5'-O5'-PBS-OAI

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301[B]	4BW	4	0
2	A	301[A]	4BW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/162 (90%)	1.23	21 (14%) 2 2	30, 44, 86, 106	0
1	B	140/162 (86%)	1.75	36 (25%) 0 0	31, 52, 94, 111	0
All	All	287/324 (88%)	1.48	57 (19%) 1 0	30, 48, 88, 111	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	118	PHE	13.5
1	A	106	VAL	7.6
1	B	116	LYS	6.9
1	A	150	ALA	6.3
1	A	235	LEU	6.2
1	B	121	LEU	5.5
1	B	117	HIS	5.3
1	B	199	LYS	5.3
1	B	180	TYR	5.2
1	B	124	THR	5.2
1	B	178	TYR	5.2
1	B	119	ASP	5.0
1	B	200	PHE	4.4
1	B	232	LEU	4.4
1	B	148	LYS	4.1
1	B	201	VAL	4.0
1	A	232	LEU	3.9
1	A	194	GLU	3.8
1	A	146	SER	3.8
1	B	123	SER	3.7
1	B	126	ILE	3.7
1	A	151	ARG	3.6
1	A	104	GLY	3.6
1	B	128	MET	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	197	LYS	3.5
1	B	125	ASN	3.5
1	A	195	LEU	3.4
1	B	225	ILE	3.3
1	B	115	PRO	3.2
1	B	120	GLU	3.2
1	B	113	TYR	3.1
1	B	216	ILE	3.1
1	B	153[A]	ARG	3.0
1	B	131	ALA	2.9
1	A	199	LYS	2.9
1	A	153[A]	ARG	2.8
1	B	155	ILE	2.8
1	A	178	TYR	2.8
1	A	145	LEU	2.7
1	B	110	ILE	2.7
1	A	148	LYS	2.6
1	A	165	HIS	2.6
1	A	107	ASN	2.6
1	B	221	LEU	2.5
1	A	181	VAL	2.5
1	B	203	LEU	2.5
1	B	181	VAL	2.4
1	B	132	GLU	2.4
1	B	135	ASN	2.4
1	B	143	ILE	2.4
1	A	147	LEU	2.3
1	B	156	LEU	2.3
1	A	193	SER	2.3
1	A	109	LYS	2.2
1	B	147	LEU	2.1
1	B	127	ASP	2.1
1	B	158	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	B	302	5/5	0.93	0.22	74,79,82,86	0
2	4BW	A	301[B]	45/45	0.94	0.15	24,30,36,37	45
2	4BW	A	301[A]	45/45	0.94	0.15	23,30,36,39	45
3	SO4	B	301	5/5	0.97	0.13	56,62,64,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4BW A 301 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.