



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 05:25 PM EDT

PDB ID : 6WV2
Title : Crystal Structure of Streptococcal Bacteriophage Hyaluronidase: Presence of a Prokaryotic Collagen and Elucidation of Catalytic Mechanism
Authors : Deivanayagam, C.; Schormann, N.
Deposited on : 2020-05-05
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	 5% 85% 8% 7%
1	B	293	 5% 84% 9% 8%
1	C	293	 5% 82% 11% 8%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6383 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hyaluronan Lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	273	Total 2044	C 1266	N 360	O 411	S 7	0	0	0
1	B	271	Total 2028	C 1257	N 358	O 406	S 7	0	0	0
1	C	271	Total 2028	C 1257	N 358	O 406	S 7	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	372	LEU	-	expression tag	UNP P15316
A	373	GLU	-	expression tag	UNP P15316
A	374	HIS	-	expression tag	UNP P15316
A	375	HIS	-	expression tag	UNP P15316
A	376	HIS	-	expression tag	UNP P15316
A	377	HIS	-	expression tag	UNP P15316
A	378	HIS	-	expression tag	UNP P15316
A	379	HIS	-	expression tag	UNP P15316
B	372	LEU	-	expression tag	UNP P15316
B	373	GLU	-	expression tag	UNP P15316
B	374	HIS	-	expression tag	UNP P15316
B	375	HIS	-	expression tag	UNP P15316
B	376	HIS	-	expression tag	UNP P15316
B	377	HIS	-	expression tag	UNP P15316
B	378	HIS	-	expression tag	UNP P15316
B	379	HIS	-	expression tag	UNP P15316
C	372	LEU	-	expression tag	UNP P15316
C	373	GLU	-	expression tag	UNP P15316
C	374	HIS	-	expression tag	UNP P15316
C	375	HIS	-	expression tag	UNP P15316
C	376	HIS	-	expression tag	UNP P15316
C	377	HIS	-	expression tag	UNP P15316
C	378	HIS	-	expression tag	UNP P15316

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	379	HIS	-	expression tag	UNP P15316

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	99	Total O 99 99	0	0
3	B	89	Total O 89 89	0	0
3	C	94	Total O 94 94	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hyaluronan Lyase



- Molecule 1: Hyaluronan Lyase



- Molecule 1: Hyaluronan Lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.45Å 86.79Å 168.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.61 – 2.21 38.61 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.61-2.21) 99.4 (38.61-2.21)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.79 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.213 , 0.237 0.213 , 0.237	Depositor DCC
R_{free} test set	2396 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	31.4	Xtrriage
Anisotropy	0.548	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6383	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/2071	0.54	0/2781
1	B	0.32	0/2055	0.51	0/2759
1	C	0.41	0/2055	0.52	0/2759
All	All	0.35	0/6181	0.52	0/8299

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	2071	30	0
1	B	2028	0	2058	29	0
1	C	2028	0	2058	46	0
2	A	1	0	0	0	0
3	A	99	0	0	1	0
3	B	89	0	0	0	0
3	C	94	0	0	0	0
All	All	6383	0	6187	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:377:HIS:CE1	1:C:377:HIS:CD2	2.21	1.28
1:A:375:HIS:CE1	1:C:375:HIS:CE1	2.33	1.17
1:C:375:HIS:CE1	1:C:377:HIS:CE1	2.49	0.99
1:B:377:HIS:HE1	1:C:377:HIS:CD2	1.71	0.99
1:C:311:ARG:CZ	1:C:313:ARG:NH2	2.32	0.92
1:B:377:HIS:CE1	1:C:377:HIS:HD2	1.84	0.86
1:A:117:LYS:HD3	1:B:117:LYS:HB3	1.57	0.84
1:A:375:HIS:HE1	1:C:375:HIS:CE1	1.95	0.81
1:B:377:HIS:CE1	1:C:377:HIS:NE2	2.48	0.81
1:A:375:HIS:CE1	1:C:375:HIS:HE1	1.96	0.79
1:A:375:HIS:NE2	1:C:375:HIS:CE1	2.51	0.78
1:A:120:LYS:HE2	1:C:118:ALA:HA	1.66	0.76
1:A:375:HIS:NE2	1:C:375:HIS:HE1	1.85	0.73
1:C:311:ARG:NH1	1:C:313:ARG:NH2	2.35	0.73
1:B:121:SER:O	1:C:126:LYS:NZ	2.20	0.73
1:C:313:ARG:NE	1:C:318:ASP:OD1	2.23	0.71
1:A:368:ILE:HG21	1:C:368:ILE:HD13	1.73	0.70
1:A:113:LEU:HD11	1:B:110:ILE:HG23	1.75	0.69
1:C:375:HIS:HE1	1:C:377:HIS:CE1	2.13	0.67
1:C:311:ARG:CZ	1:C:313:ARG:HH21	2.10	0.64
1:B:377:HIS:NE2	1:C:377:HIS:NE2	2.47	0.62
1:A:120:LYS:HE2	1:C:119:ASP:H	1.63	0.62
1:A:117:LYS:HE3	1:B:114:GLU:HA	1.81	0.61
1:A:209:VAL:HG22	1:C:220:ILE:HA	1.83	0.60
1:B:368:ILE:HD13	1:C:368:ILE:HG21	1.84	0.60
1:B:109:LYS:O	1:B:112:LYS:N	2.36	0.58
1:A:214:LYS:HE2	1:B:201:THR:HA	1.85	0.58
1:C:113:LEU:O	1:C:117:LYS:HB2	2.03	0.58
1:B:343:LYS:HE2	1:C:338:GLY:HA2	1.84	0.58
1:A:115:SER:O	1:B:120:LYS:HE2	2.05	0.57
1:A:336:VAL:HG23	1:C:340:LEU:HD11	1.86	0.57
1:C:313:ARG:HG2	1:C:318:ASP:OD1	2.06	0.54
1:A:340:LEU:HD11	1:B:336:VAL:HG23	1.89	0.54
1:C:375:HIS:CE1	1:C:377:HIS:NE2	2.76	0.53
1:A:110:ILE:HA	1:A:113:LEU:HD12	1.92	0.52
1:C:375:HIS:NE2	1:C:377:HIS:CE1	2.77	0.52
1:A:338:GLY:HA2	1:C:343:LYS:HE3	1.92	0.51
1:C:264:ASN:ND2	1:C:266:ASN:O	2.44	0.51
1:B:113:LEU:O	1:B:117:LYS:HB2	2.14	0.48
1:B:237:THR:HG23	1:B:237:THR:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:377:HIS:NE2	1:C:377:HIS:CD2	2.78	0.47
1:B:314:ASN:O	1:B:315:LYS:HB2	2.15	0.47
1:A:311:ARG:HD3	1:B:298:TYR:CD1	2.50	0.47
1:A:109:LYS:NZ	1:A:112:LYS:HD2	2.30	0.46
1:A:237:THR:HG22	1:B:221:VAL:HG22	1.98	0.46
1:C:315:LYS:O	1:C:316:ASN:HB2	2.16	0.46
1:C:127:ALA:O	1:C:131:ILE:HG12	2.16	0.45
1:B:308:LYS:HD3	1:B:321:TYR:CG	2.52	0.45
1:A:117:LYS:HD3	1:B:117:LYS:CB	2.37	0.45
1:B:340:LEU:HD11	1:C:336:VAL:HG23	1.98	0.45
1:A:221:VAL:HG22	1:C:237:THR:HG22	1.99	0.44
1:B:276:ALA:O	1:C:256:GLY:HA2	2.19	0.43
1:A:207:GLN:HE21	1:A:209:VAL:CG1	2.31	0.43
3:A:573:HOH:O	1:C:360:LYS:HD2	2.18	0.43
1:A:276:ALA:O	1:B:256:GLY:HA2	2.18	0.42
1:C:375:HIS:CE1	1:C:377:HIS:HE1	2.23	0.42
1:C:311:ARG:HD3	1:C:313:ARG:NE	2.34	0.42
1:A:233:ALA:O	1:B:217:ALA:HB3	2.20	0.42
1:A:249:ARG:NH1	1:C:263:GLU:OE1	2.53	0.42
1:B:237:THR:CG2	1:C:221:VAL:HG22	2.49	0.42
1:B:230:PHE:CE1	1:C:255:LEU:HD11	2.56	0.41
1:A:282:ILE:HD11	1:A:296:GLY:HA2	2.01	0.41
1:C:313:ARG:HG2	1:C:318:ASP:HA	2.02	0.41
1:C:282:ILE:HD11	1:C:296:GLY:HA2	2.02	0.41
1:A:256:GLY:HA2	1:C:276:ALA:O	2.21	0.41
1:A:120:LYS:HG2	1:C:118:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/293 (92%)	267 (98%)	4 (2%)	0	100	100
1	B	269/293 (92%)	263 (98%)	6 (2%)	0	100	100
1	C	269/293 (92%)	265 (98%)	4 (2%)	0	100	100
All	All	809/879 (92%)	795 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/240 (93%)	223 (100%)	0	100	100
1	B	221/240 (92%)	221 (100%)	0	100	100
1	C	221/240 (92%)	221 (100%)	0	100	100
All	All	665/720 (92%)	665 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	375	HIS
1	C	377	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/293 (93%)	0.06	14 (5%) 28 26	22, 36, 84, 136	0
1	B	271/293 (92%)	0.03	15 (5%) 25 23	22, 35, 86, 115	0
1	C	271/293 (92%)	-0.03	14 (5%) 27 25	22, 36, 85, 111	0
All	All	815/879 (92%)	0.02	43 (5%) 26 24	22, 36, 85, 136	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	376	HIS	6.5
1	B	110	ILE	5.9
1	B	112	LYS	5.3
1	A	106	THR	4.8
1	A	110	ILE	4.0
1	C	111	THR	3.9
1	A	372	LEU	3.8
1	C	112	LYS	3.8
1	A	376	HIS	3.8
1	B	111	THR	3.6
1	B	376	HIS	3.3
1	B	114	GLU	3.2
1	C	375	HIS	3.1
1	C	123	VAL	2.9
1	C	113	LEU	2.9
1	A	113	LEU	2.9
1	C	114	GLU	2.9
1	B	377	HIS	2.8
1	A	111	THR	2.8
1	C	109	LYS	2.7
1	A	112	LYS	2.6
1	C	110	ILE	2.6
1	B	109	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	113	LEU	2.5
1	A	368	ILE	2.5
1	C	115	SER	2.4
1	A	377	HIS	2.4
1	B	374	HIS	2.3
1	B	115	SER	2.3
1	A	157	ILE	2.3
1	A	154	LYS	2.3
1	B	375	HIS	2.2
1	A	270	LYS	2.2
1	B	230	PHE	2.2
1	C	230	PHE	2.1
1	B	124	TYR	2.1
1	A	370	LYS	2.1
1	B	118	ALA	2.1
1	A	373	GLU	2.1
1	C	116	SER	2.1
1	C	121	SER	2.0
1	B	122	ALA	2.0
1	C	377	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NI	A	401	1/1	0.73	0.07	65,65,65,65	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.