



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 11:37 PM EDT

PDB ID : 6WXA  
Title : Crystal structure of truncated Streptococcal bacteriophage hyaluronidase complexed with unsaturated hyaluronan hexa-saccharides  
Authors : Deivanayagam, C.; Schormann, N.  
Deposited on : 2020-05-10  
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	 5% 91% 6%
1	B	293	 4% 92% 6%
1	C	293	 3% 92% 6%
2	D	6	 83% 17%
2	G	6	 67% 33%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	5	 100%
4	F	4	 75%  25%
4	H	4	 75%  25%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 6656 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hyaluronan lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	Total 2075	C 1284	N 365	O 419	S 7	0	1	0
1	B	275	Total 2062	C 1277	N 363	O 415	S 7	0	0	0
1	C	274	Total 2053	C 1271	N 361	O 414	S 7	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	372	LEU	-	expression tag	UNP P15316
A	373	GLU	-	expression tag	UNP P15316
A	374	HIS	-	expression tag	UNP P15316
A	375	HIS	-	expression tag	UNP P15316
A	376	HIS	-	expression tag	UNP P15316
A	377	HIS	-	expression tag	UNP P15316
A	378	HIS	-	expression tag	UNP P15316
A	379	HIS	-	expression tag	UNP P15316
B	372	LEU	-	expression tag	UNP P15316
B	373	GLU	-	expression tag	UNP P15316
B	374	HIS	-	expression tag	UNP P15316
B	375	HIS	-	expression tag	UNP P15316
B	376	HIS	-	expression tag	UNP P15316
B	377	HIS	-	expression tag	UNP P15316
B	378	HIS	-	expression tag	UNP P15316
B	379	HIS	-	expression tag	UNP P15316
C	372	LEU	-	expression tag	UNP P15316
C	373	GLU	-	expression tag	UNP P15316
C	374	HIS	-	expression tag	UNP P15316
C	375	HIS	-	expression tag	UNP P15316
C	376	HIS	-	expression tag	UNP P15316
C	377	HIS	-	expression tag	UNP P15316
C	378	HIS	-	expression tag	UNP P15316

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	379	HIS	-	expression tag	UNP P15316

- Molecule 2 is an oligosaccharide called 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	6	78	42	3	33	0	0	0
2	G	6	78	42	3	33	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	E	5	67	36	3	28	0	0	0

- Molecule 4 is an oligosaccharide called 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	4	52	28	2	22	0	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	H	4	52	28	2	22	0	0	0

- Molecule 5 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ni	0	0
			1	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	39	Total	O	0	0
			39	39		
6	B	47	Total	O	0	0
			47	47		
6	C	52	Total	O	0	0
			52	52		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hyaluronan lyase

Chain A: 





- Molecule 1: Hyaluronan lyase

Chain B: 




- Molecule 1: Hyaluronan lyase

Chain C: 




- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 



- Molecule 2: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%




- Molecule 4: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  75% 25%



- Molecule 4: 4-deoxy-alpha-L-threo-hex-4-enopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-beta-D-glucopyranuronic acid-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  75% 25%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.30Å 86.49Å 168.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.91 – 2.30 44.91 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (44.91-2.30) 99.6 (44.91-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.20 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.210 , 0.243 0.213 , 0.249	Depositor DCC
$R_{free}$ test set	2231 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.2	Xtrriage
Anisotropy	0.357	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 27.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6656	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, NAG, BDP, GCD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/2102	0.80	0/2822
1	B	0.71	0/2089	0.80	0/2804
1	C	0.73	0/2080	0.80	0/2793
All	All	0.72	0/6271	0.80	0/8419

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2075	0	2095	14	0
1	B	2062	0	2090	5	0
1	C	2053	0	2077	7	0
2	D	78	0	55	0	0
2	G	78	0	55	0	0
3	E	67	0	51	0	0
4	F	52	0	37	0	0
4	H	52	0	37	0	0
5	B	1	0	0	0	0
6	A	39	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	47	0	0	0	0
6	C	52	0	0	0	0
All	All	6656	0	6497	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:LYS:HE3	1:A:273:GLU:OE1	1.78	0.83
1:A:270:LYS:HE3	1:A:273:GLU:CD	2.00	0.81
1:A:178:ALA:HB3	1:C:181:MET:HE3	1.63	0.80
1:B:373:GLU:HG2	1:B:373:GLU:O	1.86	0.76
1:A:270:LYS:HG2	1:A:270:LYS:O	1.99	0.61
1:A:178:ALA:CB	1:C:181:MET:HE3	2.32	0.57
1:A:270:LYS:HE3	1:A:273:GLU:OE2	2.04	0.57
1:A:178:ALA:HB3	1:C:181:MET:CE	2.33	0.56
1:A:109:LYS:O	1:A:113:LEU:HD23	2.05	0.56
1:A:144:VAL:HG23	1:C:148:LEU:HD11	1.97	0.47
1:A:340:LEU:HD21	1:B:340:LEU:HD22	1.98	0.45
1:A:102:GLN:CB	1:B:103:LYS:HB3	2.47	0.44
1:A:377:HIS:HE1	1:C:377:HIS:NE2	2.16	0.44
1:A:377:HIS:HE1	1:C:377:HIS:CD2	2.37	0.42
1:B:340:LEU:HD21	1:C:340:LEU:HD22	2.03	0.41
1:A:179:MET:CE	1:B:169:ILE:CD1	2.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/293 (94%)	269 (98%)	6 (2%)	0	100	100
1	B	273/293 (93%)	266 (97%)	7 (3%)	0	100	100
1	C	272/293 (93%)	267 (98%)	5 (2%)	0	100	100
All	All	820/879 (93%)	802 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/240 (94%)	226 (100%)	0	100	100
1	B	225/240 (94%)	225 (100%)	0	100	100
1	C	224/240 (93%)	224 (100%)	0	100	100
All	All	675/720 (94%)	675 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

25 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	D	1	2	15,15,15	0.16	0	21,21,21	0.40	0
2	BDP	D	2	2	12,12,13	0.29	0	14,17,19	0.62	0
2	NAG	D	3	2	14,14,15	0.28	0	17,19,21	0.58	0
2	BDP	D	4	2	12,12,13	1.09	2 (16%)	14,17,19	0.90	1 (7%)
2	NAG	D	5	2	14,14,15	0.31	0	17,19,21	0.45	0
2	GCD	D	6	2	10,11,12	0.31	0	13,15,17	0.74	0
3	NAG	E	1	3	15,15,15	0.23	0	21,21,21	0.54	0
3	BDP	E	2	3	12,12,13	0.32	0	14,17,19	0.50	0
3	NAG	E	3	3	14,14,15	0.24	0	17,19,21	0.49	0
3	BDP	E	4	3	12,12,13	0.42	0	14,17,19	1.00	0
3	NAG	E	5	3	14,14,15	0.30	0	17,19,21	0.54	0
4	NAG	F	1	4	15,15,15	0.14	0	21,21,21	0.37	0
4	BDP	F	2	4	12,12,13	0.32	0	14,17,19	0.45	0
4	NAG	F	3	4	14,14,15	0.38	0	17,19,21	0.65	0
4	GCD	F	4	4	10,11,12	2.46	4 (40%)	13,15,17	1.55	2 (15%)
2	NAG	G	1	2	15,15,15	0.32	0	21,21,21	1.14	3 (14%)
2	BDP	G	2	2	12,12,13	0.40	0	14,17,19	0.78	0
2	NAG	G	3	2	14,14,15	0.29	0	17,19,21	0.74	0
2	BDP	G	4	2	12,12,13	0.39	0	14,17,19	0.83	1 (7%)
2	NAG	G	5	2	14,14,15	0.32	0	17,19,21	0.52	0
2	GCD	G	6	2	10,11,12	0.29	0	13,15,17	1.04	0
4	NAG	H	1	4	15,15,15	0.22	0	21,21,21	0.48	0
4	BDP	H	2	4	12,12,13	0.31	0	14,17,19	0.52	0
4	NAG	H	3	4	14,14,15	0.26	0	17,19,21	0.69	0
4	GCD	H	4	4	10,11,12	0.20	0	13,15,17	2.85	5 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	BDP	D	2	2	-	0/4/21/24	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	3	2	-	0/6/23/26	0/1/1/1
2	BDP	D	4	2	-	3/4/21/24	0/1/1/1
2	NAG	D	5	2	-	0/6/23/26	0/1/1/1
2	GCD	D	6	2	-	0/4/17/20	0/1/1/1
3	NAG	E	1	3	-	0/6/26/26	0/1/1/1
3	BDP	E	2	3	-	0/4/21/24	0/1/1/1
3	NAG	E	3	3	-	0/6/23/26	0/1/1/1
3	BDP	E	4	3	-	0/4/21/24	0/1/1/1
3	NAG	E	5	3	-	0/6/23/26	0/1/1/1
4	NAG	F	1	4	-	0/6/26/26	0/1/1/1
4	BDP	F	2	4	-	0/4/21/24	0/1/1/1
4	NAG	F	3	4	-	0/6/23/26	0/1/1/1
4	GCD	F	4	4	-	2/4/17/20	0/1/1/1
2	NAG	G	1	2	-	2/6/26/26	0/1/1/1
2	BDP	G	2	2	-	2/4/21/24	0/1/1/1
2	NAG	G	3	2	-	0/6/23/26	0/1/1/1
2	BDP	G	4	2	-	1/4/21/24	0/1/1/1
2	NAG	G	5	2	-	2/6/23/26	0/1/1/1
2	GCD	G	6	2	-	0/4/17/20	0/1/1/1
4	NAG	H	1	4	-	0/6/26/26	0/1/1/1
4	BDP	H	2	4	-	0/4/21/24	0/1/1/1
4	NAG	H	3	4	-	0/6/23/26	0/1/1/1
4	GCD	H	4	4	-	2/4/17/20	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	4	GCD	O5-C1	-5.40	1.37	1.45
4	F	4	GCD	O5-C5	4.20	1.43	1.37
2	D	4	BDP	O6B-C6	-3.00	1.20	1.30
4	F	4	GCD	O6B-C6	-2.35	1.23	1.30
4	F	4	GCD	C5-C6	-2.17	1.43	1.48
2	D	4	BDP	O6A-C6	2.06	1.28	1.22

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	4	GCD	C2-C3-C4	-8.31	100.95	112.32
4	H	4	GCD	C3-C4-C5	-3.34	115.95	121.60
4	F	4	GCD	C2-C3-C4	-2.80	108.49	112.32
2	G	1	NAG	O5-C5-C4	2.79	114.77	109.69

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	NAG	C3-C4-C5	2.78	115.19	110.24
4	H	4	GCD	C1-C2-C3	-2.54	106.54	109.67
4	H	4	GCD	O5-C5-C4	-2.30	122.87	124.81
2	G	4	BDP	C1-C2-C3	2.29	112.48	109.67
2	G	1	NAG	C1-O5-C5	2.25	117.90	113.66
4	H	4	GCD	O3-C3-C2	2.17	113.18	109.42
4	F	4	GCD	C3-C4-C5	-2.12	118.01	121.60
2	D	4	BDP	O6B-C6-C5	2.10	121.33	113.65

There are no chirality outliers.


All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	5	NAG	C8-C7-N2-C2
2	G	5	NAG	O7-C7-N2-C2
4	F	4	GCD	O5-C5-C6-O6A
2	G	1	NAG	C4-C5-C6-O6
4	F	4	GCD	O5-C5-C6-O6B
4	H	4	GCD	O5-C5-C6-O6A
2	G	1	NAG	O5-C5-C6-O6
4	H	4	GCD	C4-C5-C6-O6A
2	D	4	BDP	O5-C5-C6-O6A
2	D	4	BDP	O5-C5-C6-O6B
2	G	2	BDP	O5-C5-C6-O6B
2	G	4	BDP	O5-C5-C6-O6A
2	D	4	BDP	C4-C5-C6-O6B
2	G	2	BDP	C4-C5-C6-O6B



There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	276/293 (94%)	0.16	15 (5%) 25 32	32, 47, 87, 128	0
1	B	275/293 (93%)	0.15	11 (4%) 38 45	31, 45, 87, 115	0
1	C	274/293 (93%)	0.15	10 (3%) 42 49	30, 45, 95, 117	0
All	All	825/879 (93%)	0.16	36 (4%) 34 41	30, 46, 90, 128	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	376	HIS	5.6
1	C	113	LEU	4.6
1	A	110	ILE	4.2
1	C	111	THR	3.7
1	C	377	HIS	3.6
1	B	123	VAL	3.3
1	A	115	SER	3.1
1	C	112	LYS	3.0
1	C	124	TYR	2.9
1	B	376	HIS	2.9
1	C	115	SER	2.9
1	A	116	SER	2.8
1	A	122	ALA	2.8
1	B	112	LYS	2.8
1	A	377	HIS	2.8
1	B	111	THR	2.7
1	B	110	ILE	2.7
1	B	368	ILE	2.6
1	A	372	LEU	2.6
1	C	373	GLU	2.6
1	B	122	ALA	2.6
1	C	376	HIS	2.5
1	B	115	SER	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	112	LYS	2.5
1	A	124	TYR	2.5
1	A	108	SER	2.4
1	B	127	ALA	2.4
1	B	374	HIS	2.4
1	A	113	LEU	2.4
1	A	121	SER	2.3
1	C	116	SER	2.2
1	A	119	ASP	2.2
1	A	114	GLU	2.2
1	C	123	VAL	2.1
1	B	106	THR	2.1
1	A	127	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GCD	D	6	11/12	0.80	0.23	74,91,99,102	0
2	GCD	G	6	11/12	0.80	0.26	84,90,95,96	0
4	GCD	H	4	11/12	0.84	0.26	66,78,86,86	0
4	NAG	H	1	15/15	0.85	0.12	46,66,72,73	0
4	GCD	F	4	11/12	0.85	0.19	66,71,83,84	0
3	NAG	E	5	14/15	0.86	0.14	46,55,64,64	0
4	NAG	F	1	15/15	0.89	0.14	52,63,69,69	0
2	NAG	G	1	15/15	0.89	0.21	50,57,64,66	0
2	BDP	D	2	12/13	0.91	0.13	40,44,48,48	0
4	BDP	F	2	12/13	0.92	0.11	47,49,59,61	0
4	NAG	H	3	14/15	0.92	0.11	39,51,61,61	0
3	BDP	E	4	12/13	0.92	0.10	49,53,56,61	0
2	NAG	D	5	14/15	0.93	0.12	47,53,64,68	0
4	NAG	F	3	14/15	0.93	0.12	41,48,54,56	0
2	NAG	G	5	14/15	0.93	0.14	46,54,64,65	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BDP	G	4	12/13	0.94	0.17	44,53,61,65	0
3	NAG	E	1	15/15	0.94	0.14	44,53,63,63	0
4	BDP	H	2	12/13	0.94	0.11	54,61,62,64	0
3	BDP	E	2	12/13	0.94	0.11	41,47,49,50	0
2	BDP	G	2	12/13	0.94	0.12	47,54,59,64	0
2	NAG	G	3	14/15	0.95	0.14	42,47,49,50	0
2	NAG	D	1	15/15	0.95	0.12	42,51,59,59	0
3	NAG	E	3	14/15	0.96	0.11	37,44,47,49	0
2	BDP	D	4	12/13	0.97	0.13	35,40,57,58	0
2	NAG	D	3	14/15	0.97	0.15	32,38,40,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NI	B	406	1/1	0.90	0.04	80,80,80,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.