



wwPDB X-ray Structure Validation Summary Report

Jun 12, 2024 – 06:35 PM EDT

PDB ID : 3X2R
Title : Structure of the nonameric bacterial amyloid secretion channel CsgG
Authors : Huang, Y.; Cao, B.; Zhao, Y.; Kou, Y.; Ni, D.; Zhang, X.C.
Deposited on : 2014-12-29
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	
1	B	277	
1	C	277	
1	D	277	
1	E	277	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	277	
1	G	277	
1	H	277	
1	I	277	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 13042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CsgG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	189	1424	904	236	279	5	0	0	0
1	A	196	1467	932	243	287	5	0	0	0
1	B	190	1427	904	236	282	5	0	0	0
1	C	198	1498	954	248	291	5	0	0	0
1	E	198	1498	954	248	291	5	0	0	0
1	F	189	1424	904	236	279	5	0	0	0
1	G	190	1440	918	237	280	5	0	0	0
1	H	191	1441	915	238	283	5	0	0	0
1	I	190	1423	902	236	280	5	0	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CsgG




- Molecule 1: CsgG




- Molecule 1: CsgG






• Molecule 1: CsgG



• Molecule 1: CsgG



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	171.88Å 176.72Å 103.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.07 – 2.90 48.07 – 2.80	Depositor EDS
% Data completeness (in resolution range)	86.8 (48.07-2.90) 75.9 (48.07-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.244 , 0.292 0.247 , 0.294	Depositor DCC
R_{free} test set	3428 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	58.7	Xtrriage
Anisotropy	0.943	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 49.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for k,h,-l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13042	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1488	0.63	0/2019
1	B	0.33	0/1444	0.61	0/1956
1	C	0.39	0/1522	0.67	1/2067 (0.0%)
1	D	0.34	0/1443	0.62	0/1956
1	E	0.38	0/1522	0.67	1/2067 (0.0%)
1	F	0.33	0/1443	0.59	0/1956
1	G	0.34	0/1461	0.61	0/1983
1	H	0.32	0/1460	0.61	0/1979
1	I	0.33	0/1440	0.61	0/1951
All	All	0.35	0/13223	0.63	2/17934 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	251	LEU	CA-CB-CG	7.25	131.99	115.30
1	C	251	LEU	CA-CB-CG	6.38	129.98	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	55	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1467	0	1451	55	0
1	B	1427	0	1415	43	0
1	C	1498	0	1480	50	1
1	D	1424	0	1408	43	0
1	E	1498	0	1480	52	1
1	F	1424	0	1408	44	0
1	G	1440	0	1425	55	0
1	H	1441	0	1425	55	0
1	I	1423	0	1411	53	0
All	All	13042	0	12903	373	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 373 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:92:TRP:HE3	1:E:92:TRP:H	1.22	0.83
1:C:150:LYS:HB3	1:C:169:LEU:HD12	1.61	0.82
1:E:196:ILE:HD13	1:E:227:VAL:HG22	1.64	0.80
1:G:49:ILE:HG21	1:G:92:TRP:HB2	1.65	0.79
1:E:48:LYS:HG2	1:E:92:TRP:HB2	1.66	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:244:ASN:OD1	1:E:244:ASN:ND2[4_555]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/277 (69%)	176 (93%)	12 (6%)	2 (1%)	14	42
1	B	184/277 (66%)	175 (95%)	8 (4%)	1 (0%)	29	61
1	C	192/277 (69%)	176 (92%)	15 (8%)	1 (0%)	29	61
1	D	183/277 (66%)	173 (94%)	9 (5%)	1 (0%)	29	61
1	E	192/277 (69%)	176 (92%)	15 (8%)	1 (0%)	29	61
1	F	183/277 (66%)	173 (94%)	9 (5%)	1 (0%)	29	61
1	G	184/277 (66%)	173 (94%)	10 (5%)	1 (0%)	29	61
1	H	185/277 (67%)	174 (94%)	10 (5%)	1 (0%)	29	61
1	I	184/277 (66%)	175 (95%)	8 (4%)	1 (0%)	29	61
All	All	1677/2493 (67%)	1571 (94%)	96 (6%)	10 (1%)	25	58

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	149	VAL
1	A	149	VAL
1	B	149	VAL
1	C	149	VAL
1	E	149	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	157/236 (66%)	133 (85%)	24 (15%)	2	8
1	B	154/236 (65%)	132 (86%)	22 (14%)	3	10
1	C	161/236 (68%)	133 (83%)	28 (17%)	2	6
1	D	153/236 (65%)	128 (84%)	25 (16%)	2	7
1	E	161/236 (68%)	137 (85%)	24 (15%)	3	9
1	F	153/236 (65%)	131 (86%)	22 (14%)	3	9
1	G	155/236 (66%)	132 (85%)	23 (15%)	3	9
1	H	155/236 (66%)	134 (86%)	21 (14%)	4	11
1	I	153/236 (65%)	131 (86%)	22 (14%)	3	9
All	All	1402/2124 (66%)	1191 (85%)	211 (15%)	3	9

5 of 211 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	151	SER
1	F	235	ILE
1	I	101	LEU
1	E	231	LEU
1	F	80	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	224	ASN
1	I	77	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/277 (70%)	0.44	19 (9%) 7 6	49, 76, 153, 185	0
1	B	190/277 (68%)	0.27	10 (5%) 26 22	46, 76, 158, 191	0
1	C	198/277 (71%)	0.74	20 (10%) 7 5	47, 70, 156, 199	0
1	D	189/277 (68%)	0.45	17 (8%) 9 7	48, 74, 149, 188	0
1	E	198/277 (71%)	0.56	20 (10%) 7 5	47, 73, 156, 191	0
1	F	189/277 (68%)	0.51	13 (6%) 16 13	50, 77, 145, 192	0
1	G	190/277 (68%)	0.47	17 (8%) 9 7	50, 78, 168, 187	0
1	H	191/277 (68%)	0.45	13 (6%) 17 13	47, 79, 157, 188	0
1	I	190/277 (68%)	0.47	12 (6%) 20 16	45, 77, 157, 186	0
All	All	1731/2493 (69%)	0.49	141 (8%) 12 9	45, 76, 157, 199	0

The worst 5 of 141 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	158	TYR	6.9
1	B	251	LEU	6.1
1	C	206	PHE	6.0
1	G	215	LEU	5.5
1	A	208	PHE	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.