



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 13, 2023 – 12:00 PM JST

PDB ID : 5X6C
Title : Crystal structure of SepRS-SepCysE from Methanocaldococcus jannaschii
Authors : Chen, M.; Kato, K.; Yao, M.
Deposited on : 2017-02-21
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	 80% 16%
1	B	553	 81% 16%
2	E	216	 7% 88%
2	F	216	 8% 88%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9363 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-phosphoserine--tRNA(Cys) ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	539	4379	2826	727	807	19	0	0	0
1	B	539	4379	2826	727	807	19	0	0	0


- Molecule 2 is a protein called Uncharacterized protein MJ1481.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	26	212	133	41	37	1	0	0	0
2	F	26	212	133	41	37	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	-2	MET	-	initiating methionine	UNP Q58876
E	-1	ASN	-	expression tag	UNP Q58876
E	0	HIS	-	expression tag	UNP Q58876
F	-2	MET	-	initiating methionine	UNP Q58876
F	-1	ASN	-	expression tag	UNP Q58876
F	0	HIS	-	expression tag	UNP Q58876

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	5	13	3	0	0
3	B	1	31	10	5	13	3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
4	A	1	5	4 1	0	0
4	B	1	5	4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	53	Total 53	O 53	0	0
5	B	56	Total 56	O 56	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: O-phosphoserine--tRNA(Cys) ligase



- Molecule 1: O-phosphoserine--tRNA(Cys) ligase



- Molecule 2: Uncharacterized protein MJ1481



LYS ASN
GLU LYS
ASN ASP
ASN LYS
PRO PRO
LYS ASP
ASN LYS
LEU VAL
LYS VAL
VAL CYS
LEU THR
LEU ARG
GLU THR
GLY LYS
MET LEU
LEU ASP
LEU ASP
ILE ASP
GLU ASP
GLU MET
LYS ARG
LEU ARG
ALA ASP
ALA ASP
ILE ASP
GLY ASP
VAL VAL
PRO VAL
GLU LYS
LYS LYS
LYS VAL
LEU VAL
GLU GLY
GLY LYS
LYS VAL
VAL VAL
GLU ARG
ALA ASP
LYS ASN
ASN VAL
VAL VAL
ARG ARG
PHE ARG
ILE ILE
GLU GLU
LYS LYS
TYR TYR
LYS LYS
PRO PRO
GLU LYS
ILE ILE
PHE PHE
VAL VAL
VAL VAL

VAL
GLU
ASP
ASP
LYS
ASP
GLU
LEU
LEU
TYR
LEU
ARG
ALA
LYS
ASN
LEU
TYR
ASN
ALA
GLU
LYS
LEU
ASP

• Molecule 2: Uncharacterized protein MJ1481

Chain F: 8% . 88%

MET
R-I
R-O
M1
R2
V3
K7
R11
K12
S15
S18
K21
K22
K23
K24
ILE
ARG
VAL
VAL
GLN
LYS
ASP
ASP
LYS
LYS
ILE
ILE
SER
TYR
LYS
ASP
ALA
PHE
LYS
PHE
LYS
PRO
GLY
GLY
LYS
ILE
ASP
VAL
ASN
GLU
PHE
LYS
ALA
TYR
TYR
LEU
LEU
LEU
ASP
ASP
PHE
PHE
TYR

LYS
LYS
ALA
PRO
LYS
HIS
GLU
LEU
ASN
VAL
GLU
GLU
GLU
ALA
LYS
PHE
CYS
LYS
LEU
ILE
ILE
LYS
CYS
GLN
GLY
HIS
LEU
ASN
LYS
ILE
ALA
LEU
ALA
ASN
PHE
GLY
PHE
LYS
PHE
GLY
GLU
LYS
GLU
LYS
ILE
ASP
GLU
GLY
LYS
ALA
LEU
TYR
ILE
VAL
SER
ASN
LYS
LYS
GLY
LEU
LEU
LYS
PHE
LEU
ASP
ASP
PHE
LYS
LEU

LYS
ASN
LYS
ASN
PRO
LYS
ASN
GLU
LYS
VAL
VAL
CYS
THR
GLU
GLY
MET
LEU
LEU
ASP
ILE
GLU
ASP
MET
LYS
ARG
ALA
ILE
ILE
GLY
VAL
PRO
GLU
LYS
ALA
LEU
GLY
LYS
LYS
VAL
GLY
ILE
ALA
ARG
LYS
ASN
VAL
GLU
ARG
PHE
ILE
GLY
LYS
TYR
LYS
PRO
GLY
ILE
PHE
VAL

VAL
GLU
ASP
ASP
LYS
ASP
GLU
LEU
LEU
TYR
LEU
ARG
ALA
LYS
ASN
LEU
TYR
ASN
ALA
GLU
LYS
LEU
LEU
ASP
ILE
LEU
LEU
LEU
ASP

4 Data and refinement statistics i

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	279.82Å 279.82Å 279.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.99 – 3.10 47.99 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.99-3.10) 99.9 (47.99-3.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.209 0.192 , 0.217	Depositor DCC
R_{free} test set	3331 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	68.8	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 23.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.477 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9363	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/4468	0.64	1/6023 (0.0%)
1	B	0.64	0/4468	0.63	0/6023
2	E	0.44	0/213	0.64	0/280
2	F	0.47	0/213	0.62	0/280
All	All	0.64	0/9362	0.64	1/12606 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	ARG	NE-CZ-NH2	5.81	123.20	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	118	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4379	0	4462	67	0
1	B	4379	0	4462	57	0
2	E	212	0	236	11	0
2	F	212	0	236	7	0
3	A	31	0	12	0	0
3	B	31	0	12	0	0
4	A	5	0	0	1	0
4	B	5	0	0	1	0
5	A	53	0	0	1	0
5	B	56	0	0	1	0
All	All	9363	0	9420	127	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:ASN:HD22	1:A:330:LEU:N	1.77	0.83
1:A:386:GLU:OE1	1:A:407:ARG:NH1	2.10	0.83
1:A:363:ARG:HG3	1:A:363:ARG:HH21	1.48	0.77
1:A:118:GLY:HA2	1:A:122:VAL:HG23	1.70	0.72
1:B:394:ASN:HD22	1:B:403:VAL:HG21	1.55	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/553 (97%)	523 (97%)	13 (2%)	1 (0%)	47	79
1	B	537/553 (97%)	519 (97%)	17 (3%)	1 (0%)	47	79
2	E	24/216 (11%)	24 (100%)	0	0	100	100
2	F	24/216 (11%)	24 (100%)	0	0	100	100
All	All	1122/1538 (73%)	1090 (97%)	30 (3%)	2 (0%)	47	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	507	PRO
1	B	507	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	481/494 (97%)	471 (98%)	10 (2%)	53	79
1	B	481/494 (97%)	477 (99%)	4 (1%)	81	92
2	E	24/195 (12%)	23 (96%)	1 (4%)	30	62
2	F	24/195 (12%)	23 (96%)	1 (4%)	30	62
All	All	1010/1378 (73%)	994 (98%)	16 (2%)	62	84

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	12	LYS
1	B	460	GLN
1	A	363	ARG
1	B	212	LYS
1	A	329	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	460	GLN
1	B	329	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	602	-	4,4,4	0.62	0	6,6,6	0.24	0
3	ATP	B	601	-	26,33,33	1.05	2 (7%)	31,52,52	1.44	6 (19%)
3	ATP	A	601	-	26,33,33	1.01	1 (3%)	31,52,52	1.51	6 (19%)
4	SO4	B	602	-	4,4,4	0.58	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	B	601	-	-	5/18/38/38	0/3/3/3
3	ATP	A	601	-	-	6/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	ATP	C2'-C1'	-2.65	1.49	1.53
3	A	601	ATP	C2'-C1'	-2.34	1.50	1.53
3	B	601	ATP	C5-C4	2.02	1.46	1.40

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	ATP	C3'-C2'-C1'	3.55	106.32	100.98
3	B	601	ATP	C3'-C2'-C1'	3.19	105.78	100.98
3	A	601	ATP	N3-C2-N1	-3.12	123.80	128.68
3	B	601	ATP	N3-C2-N1	-2.86	124.20	128.68
3	B	601	ATP	C2'-C3'-C4'	2.69	107.86	102.64

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	ATP	C5'-O5'-PA-O1A
3	B	601	ATP	C5'-O5'-PA-O2A
3	B	601	ATP	PA-O3A-PB-O3B
3	A	601	ATP	C5'-O5'-PA-O3A
3	B	601	ATP	C5'-O5'-PA-O3A


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	602	SO4	1	0
4	B	602	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	539/553 (97%)	0.12	0 100 100	25, 40, 69, 96	0
1	B	539/553 (97%)	0.11	0 100 100	25, 41, 68, 98	0
2	E	26/216 (12%)	0.14	0 100 100	42, 61, 90, 102	0
2	F	26/216 (12%)	0.11	0 100 100	42, 61, 89, 100	0
All	All	1130/1538 (73%)	0.11	0 100 100	25, 42, 72, 102	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ATP	B	601	31/31	0.93	0.23	34,40,71,88	0
3	ATP	A	601	31/31	0.95	0.24	33,41,70,87	0
4	SO4	A	602	5/5	0.98	0.22	33,36,43,59	0
4	SO4	B	602	5/5	0.98	0.24	34,34,41,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ATP B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.