



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2023 – 02:13 PM JST

PDB ID : 7X7H  
Title : Crystal structure of Fructose regulator/Histidine phosphocarrier protein complex from *Vibrio cholerae*  
Authors : Kim, M.-K.; Zhang, J.; Yoon, C.-K.; Seok, Y.-J.  
Deposited on : 2022-03-09  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.32.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	 2% 70% 11% • 18%
1	C	326	 2% 68% 13% • 18%
2	B	91	 9% 81% 12% 7%
2	D	91	 8% 75% 16% • 7%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11372 atoms, of which 5484 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catabolite repressor/activator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	267	4170	1315	2091	359	400	5	0	0	0
1	C	267	4168	1315	2089	359	400	5	0	0	0

- Molecule 2 is a protein called HPr family phosphocarrier protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	85	1293	404	652	111	125	1	0	0	0
2	D	85	1293	404	652	111	125	1	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	MET	-	initiating methionine	UNP A0A085PY40
B	-4	HIS	-	expression tag	UNP A0A085PY40
B	-3	HIS	-	expression tag	UNP A0A085PY40
B	-2	HIS	-	expression tag	UNP A0A085PY40
B	-1	HIS	-	expression tag	UNP A0A085PY40
B	0	HIS	-	expression tag	UNP A0A085PY40
B	1	HIS	-	expression tag	UNP A0A085PY40
D	-5	MET	-	initiating methionine	UNP A0A085PY40
D	-4	HIS	-	expression tag	UNP A0A085PY40
D	-3	HIS	-	expression tag	UNP A0A085PY40
D	-2	HIS	-	expression tag	UNP A0A085PY40
D	-1	HIS	-	expression tag	UNP A0A085PY40
D	0	HIS	-	expression tag	UNP A0A085PY40
D	1	HIS	-	expression tag	UNP A0A085PY40

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	195	Total O 195 195	0	0
4	B	54	Total O 54 54	0	0
4	C	172	Total O 172 172	0	0
4	D	25	Total O 25 25	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Catabolite repressor/activator




- Molecule 1: Catabolite repressor/activator




- Molecule 2: HPr family phosphocarrier protein



- Molecule 2: HPr family phosphocarrier protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.94Å 62.35Å 312.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.13 – 2.00 31.13 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (31.13-2.00) 99.9 (31.13-2.00)	Depositor EDS
$R_{merge}$	1.00	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.24 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, $R_{free}$	0.179 , 0.227 0.180 , 0.183	Depositor DCC
$R_{free}$ test set	2283 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtrriage
Anisotropy	0.333	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 50.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11372	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.54	5/2112 (0.2%)	0.69	4/2864 (0.1%)
1	C	0.31	0/2112	0.58	0/2864
2	B	0.30	0/650	0.51	0/878
2	D	0.47	1/650 (0.2%)	0.65	1/878 (0.1%)
All	All	0.43	6/5524 (0.1%)	0.63	5/7484 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	227	LYS	CD-CE	15.04	1.88	1.51
1	A	228	VAL	CB-CG1	6.80	1.67	1.52
2	D	71	LYS	CE-NZ	6.33	1.64	1.49
1	A	227	LYS	CE-NZ	6.26	1.64	1.49
1	A	236	ASP	CG-OD2	6.12	1.39	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	236	ASP	CB-CG-OD1	-11.77	107.70	118.30
1	A	227	LYS	CB-CG-CD	-9.18	87.73	111.60
2	D	40	LYS	CD-CE-NZ	-8.56	92.01	111.70
1	A	277	LEU	CA-CB-CG	6.88	131.12	115.30
1	A	236	ASP	OD1-CG-OD2	6.34	135.36	123.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2079	2091	2091	29	0
1	C	2079	2089	2091	29	0
2	B	641	652	654	11	0
2	D	641	652	654	11	1
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	195	0	0	5	1
4	B	54	0	0	0	0
4	C	172	0	0	5	2
4	D	25	0	0	0	0
All	All	5888	5484	5490	77	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:LYS:CD	1:A:227:LYS:CE	1.88	1.47
1:C:166:ALA:HB1	1:C:288:LEU:HD21	1.59	0.84
1:A:293:GLU:O	4:A:501:HOH:O	1.98	0.81
1:A:223:GLU:HG2	1:A:227:LYS:HE3	1.62	0.80
1:A:79:LEU:HD21	1:A:147:ILE:HD12	1.64	0.80

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:82:ASP:OD2	4:C:504:HOH:O[4_445]	2.00	0.20
4:A:695:HOH:O	4:C:654:HOH:O[1_655]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/326 (81%)	263 (99%)	2 (1%)	0	100	100
1	C	265/326 (81%)	261 (98%)	4 (2%)	0	100	100
2	B	83/91 (91%)	81 (98%)	2 (2%)	0	100	100
2	D	83/91 (91%)	81 (98%)	2 (2%)	0	100	100
All	All	696/834 (84%)	686 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/270 (83%)	221 (99%)	2 (1%)	78	83
1	C	223/270 (83%)	217 (97%)	6 (3%)	44	46
2	B	68/74 (92%)	67 (98%)	1 (2%)	65	69
2	D	68/74 (92%)	66 (97%)	2 (3%)	42	43
All	All	582/688 (85%)	571 (98%)	11 (2%)	57	61

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	268	ARG
1	C	312	GLN
2	D	81	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	1	HIS
1	C	207	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	267/326 (81%)	0.08	7 (2%) 56 54	12, 23, 51, 65	0
1	C	267/326 (81%)	0.18	7 (2%) 56 54	14, 29, 56, 73	0
2	B	85/91 (93%)	0.47	8 (9%) 8 8	21, 33, 63, 80	0
2	D	85/91 (93%)	0.59	7 (8%) 11 11	26, 44, 67, 77	0
All	All	704/834 (84%)	0.22	29 (4%) 37 36	12, 29, 59, 80	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	326	THR	7.1
2	B	0	HIS	6.6
1	C	312	GLN	4.2
2	B	1	HIS	4.2
2	B	68	PRO	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	400	1/1	1.00	0.07	13,13,13,13	0
3	CA	C	400	1/1	1.00	0.05	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.