



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 21, 2023 – 09:47 PM JST

PDB ID : 7X7O  
Title : SARS-CoV-2 spike RBD in complex with neutralizing antibody UT28K  
Authors : Ozawa, T.; Tani, H.; Anraku, Y.; Kita, S.; Igarashi, E.; Saga, Y.; Inasaki, N.;  
Kawasuji, H.; Yamada, H.; Sasaki, S.; Somekawa, M.; Sasaki, J.; Hayakawa,  
Y.; Yamamoto, Y.; Morinaga, Y.; Kurosawa, N.; Isobe, M.; Fukuhara, H.;  
Maenaka, K.; Hashiguchi, T.; Kishi, H.; Kitajima, I.; Saito, S.; Niimi, H.  
Deposited on : 2022-03-10  
Resolution : 3.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*







The reported resolution of this entry is 3.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1039 (3.94-3.58)
Clashscore	141614	1051 (3.92-3.60)
Ramachandran outliers	138981	1015 (3.92-3.60)
Sidechain outliers	138945	1011 (3.92-3.60)
RSRZ outliers	127900	1050 (3.96-3.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	
1	B	215	
1	C	215	
1	D	215	
2	E	257	
2	G	257	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	257	
2	J	257	
3	F	235	
3	I	235	
3	K	235	
3	L	235	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	167	Total 1334	C 854	N 225	O 249	S 6	0	0	0
1	B	172	Total 1371	C 880	N 231	O 253	S 7	0	0	0
1	C	157	Total 1261	C 809	N 213	O 233	S 6	0	0	0
1	D	163	Total 1309	C 840	N 220	O 243	S 6	0	0	0

- Molecule 2 is a protein called UT28K Fab, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	122	Total 935	C 587	N 159	O 181	S 8	0	0	0
2	G	122	Total 935	C 587	N 159	O 181	S 8	0	0	0
2	H	122	Total 935	C 587	N 159	O 181	S 8	0	0	0
2	J	122	Total 935	C 587	N 159	O 181	S 8	0	0	0


- Molecule 3 is a protein called UT28K Fab, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	F	107	Total 816	C 513	N 138	O 163	S 2	0	0	0
3	I	107	Total 816	C 513	N 138	O 163	S 2	0	0	0
3	K	107	Total 816	C 513	N 138	O 163	S 2	0	0	0
3	L	107	Total 816	C 513	N 138	O 163	S 2	0	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 1: Spike protein S1






- Molecule 1: Spike protein S1











• Molecule 1: Spike protein S1



• Molecule 2: UT28K Fab, heavy chain




• Molecule 2: UT28K Fab, heavy chain



TYR  
ILE  
CYS  
ASN  
VAL  
VAL  
ASN  
HIS  
LYS  
PRO  
SER  
SER  
ASN  
THR  
LYS  
VAL  
ASP  
LYS  
VAL  
GLY  
LYS  
VAL  
GLY  
PRO

- Molecule 2: UT28K Fab, heavy chain




MET  
ASP  
TRP  
ILE  
TRP  
ARG  
ARG  
ILE  
LEU  
PHE  
LEU  
VAL  
VAL  
GLY  
ALA  
ALA  
THR  
GLY  
ALA  
HIS  
HIS  
GLY  
SER  
GLN  
M2  
Q3  
L4  
S7  
V11  
K12  
K19  
V20  
S21  
C22  
A24  
T28  
A33  
V34  
Q35  
W36  
V37  
R38  
Q39  
A40  
Q43  
R44  
L45  
E46  
W47  
I48  
V53  
Y60  
F64  
Q65

E66  
R67  
V68  
T69  
I70  
T71  
R72  
D73  
M74  
M81  
E82  
T86  
H87  
Y94  
D104  
C105  
S106  
D110  
I111  
W112  
V118  
T119  
V120  
A123  
SER  
THR  
LYS  
GLY  
PRO  
SER  
VAL  
PHE  
PRO  
LEU  
ALA  
PRO  
SER  
SER  
LYS  
GLY  
THR  
SER  
GLN  
THR  
TYR  
THR  
GLY  
ALA  
CYS  
ASN  
VAL  
LEU  
GLY  
CYS  
LEU

VAL  
LYS  
ASP  
TYR  
PHE  
PRO  
GLU  
PRO  
VAL  
THR  
VAL  
VAL  
SER  
TPP  
ASN  
SER  
GLY  
SER  
ALA  
HIS  
THR  
SER  
GLY  
VAL  
VAL  
THR  
HIS  
HIS  
PHE  
PRO  
ALA  
VAL  
GLN  
SER  
SER  
SER  
GLY  
THR  
TYR  
SER  
SER  
VAL  
THR  
THR  
TYR  
ILE  
HIS  
ASN  
VAL  
HIS  
LYS

PRO  
SER  
ASN  
THR  
LYS  
VAL  
ASP  
LYS  
PRO  
LYS  
SER  
GLY  
SER  
SER  
HIS  
HIS  
HIS  
HIS  
HIS  
HIS

- Molecule 2: UT28K Fab, heavy chain




MET  
ASP  
TRP  
ILE  
TRP  
ARG  
ARG  
ILE  
LEU  
PHE  
LEU  
VAL  
VAL  
GLY  
ALA  
ALA  
THR  
GLY  
SER  
HIS  
HIS  
GLN  
M2  
Q3  
L4  
V5  
Q6  
E10  
V11  
K12  
V18  
K23  
A24  
F27  
S32  
A33  
V34  
Q35  
W36  
V37  
R38  
Q39  
A40  
R41  
G42  
Q43  
R44  
L45  
G54  
S55  
G56  
M57  
T58  
F64

V68  
T69  
I70  
T71  
R72  
D73  
M74  
M81  
E82  
S85  
L86  
R87  
T91  
C101  
C105  
S106  
D107  
D110  
I111  
W112  
V118  
T119  
V120  
S121  
S122  
A123  
SER  
THR  
LYS  
GLY  
PRO  
SER  
VAL  
PHE  
PRO  
LEU  
ALA  
PRO  
SER  
SER  
LYS  
GLY  
THR  
SER  
GLN  
THR  
TYR  
THR  
GLY  
ALA  
CYS  
ASN  
VAL  
LEU  
GLY  
CYS  
LEU

VAL  
LYS  
ASP  
TYR  
PHE  
PRO  
GLU  
PRO  
VAL  
THR  
VAL  
VAL  
SER  
TPP  
ASN  
SER  
GLY  
SER  
ALA  
HIS  
THR  
SER  
GLY  
VAL  
VAL  
THR  
HIS  
HIS  
PHE  
PRO  
ALA  
VAL  
GLN  
SER  
SER  
SER  
VAL  
THR  
TYR  
SER  
SER  
VAL  
THR  
THR  
TYR  
ILE  
HIS  
ASN  
VAL  
HIS  
LYS

PRO  
SER  
ASN  
THR  
LYS  
VAL  
ASP  
LYS  
PRO  
LYS  
SER  
GLY  
SER  
SER  
HIS  
HIS  
HIS  
HIS  
HIS  
HIS

- Molecule 3: UT28K Fab, light chain



MET  
GLU  
THR  
PRO  
ALA  
GLN  
LEU  
LEU  
PHE  
LEU  
VAL  
LEU  
LEU  
LEU  
TRP  
LEU  
PRO  
ASP  
THR  
THR  
GLY  
E1  
I2  
S26  
Q27  
S28  
V29  
Q38  
Q39  
R39  
P45  
R46  
L47  
L48  
I49  
Y50  
S54  
R55  
A56  
T57  
G58  
I59  
R62  
F63  
I76  
P81  
E82  
D83  
F84  
A85  
Q90  
Q91  
N94

F99  
G100  
Q101  
I107  
LYS  
ARG  
THR  
VAL  
ASP  
ALA  
ALA  
PRO  
PRO  
SER  
VAL  
PHE  
PHE  
THR  
PRO  
SER  
SER  
LEU  
LEU  
SER  
SER  
ASP  
GLU  
GLN  
ALA  
ASP  
TYR  
GLY  
GLY  
HIS  
LYS  
VAL  
THR  
VAL  
VAL  
ALA  
CYS  
GLU  
VAL  
VAL  
HIS  
GLN  
TYR  
LEU  
SER  
ARG  
GLY  
ALA  
VAL  
VAL  
THR  
LYS  
GLN  
TRP  
LYS  
VAL  
ASP  
ASN  
ARG  
GLY  
ALA  
GLN  
SER  
SER  
GLY  
ASN  
SER  
GLN

GLU  
SER  
VAL  
THR  
GLU  
GLN  
ASP  
SER  
LYS  
ASP  
SER  
THR  
THR  
SER  
SER  
PHE  
SER  
PHE  
THR  
THR  
THR  
LEU  
LEU  
SER  
SER  
ASP  
SER  
LYS  
SER  
ASP  
GLU  
GLY  
HIS  
VAL  
TYR  
ALA  
CYS  
GLU  
VAL  
VAL  
HIS  
GLN  
TYR  
LEU  
SER  
ARG  
GLY  
ALA  
VAL  
VAL  
THR  
LYS  
GLN  
TRP  
PHE  
ASN  
ARG  
GLY  
CYS

- Molecule 3: UT28K Fab, light chain





## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	138.25Å 138.41Å 221.11Å 90.00° 90.05° 90.00°	Depositor
Resolution (Å)	48.14 – 3.75 48.91 – 3.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.14-3.75) 99.7 (48.91-3.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 3.77Å)	Xtrriage
Refinement program	REFMAC 5.8.0267, PHENIX 1.18_3845	Depositor
R, $R_{free}$	0.303 , 0.336 0.315 , 0.339	Depositor DCC
$R_{free}$ test set	2193 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	111.2	Xtrriage
Anisotropy	0.307	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 94.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.398 for k,h,-l 0.367 for -k,-h,-l 0.380 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.84	EDS
Total number of atoms	12279	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	133.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1367	0.48	0/1851
1	B	0.35	0/1408	0.51	0/1909
1	C	0.34	0/1295	0.48	0/1755
1	D	0.36	0/1342	0.50	0/1817
2	E	0.34	0/955	0.53	0/1293
2	G	0.33	0/955	0.54	0/1293
2	H	0.39	0/955	0.55	0/1293
2	J	0.36	0/955	0.55	0/1293
3	F	0.28	0/836	0.48	0/1137
3	I	0.29	0/836	0.48	0/1137
3	K	0.35	0/836	0.52	0/1137
3	L	0.31	0/836	0.50	0/1137
All	All	0.34	0/12576	0.51	0/17052

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1334	0	1261	39	0
1	B	1371	0	1290	46	0
1	C	1261	0	1190	16	0
1	D	1309	0	1232	26	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	935	0	899	30	0
2	G	935	0	899	40	0
2	H	935	0	899	24	0
2	J	935	0	899	28	0
3	F	816	0	786	14	0
3	I	816	0	786	14	0
3	K	816	0	786	13	0
3	L	816	0	786	14	0
All	All	12279	0	11713	282	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 282 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:ALA:CB	2:G:31:ILE:HD11	1.71	1.20
1:A:475:ALA:HB1	2:G:31:ILE:HD11	1.10	1.09
1:A:473:TYR:HE2	2:G:31:ILE:HD12	1.20	1.05
2:G:31:ILE:HG13	2:G:31:ILE:O	1.61	0.97
2:G:31:ILE:HG12	2:G:101:CYS:SG	2.08	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	157/215 (73%)	147 (94%)	10 (6%)	0	100 100
1	B	164/215 (76%)	154 (94%)	10 (6%)	0	100 100
1	C	151/215 (70%)	141 (93%)	10 (7%)	0	100 100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	153/215 (71%)	143 (94%)	10 (6%)	0	100	100
2	E	120/257 (47%)	115 (96%)	5 (4%)	0	100	100
2	G	120/257 (47%)	109 (91%)	10 (8%)	1 (1%)	19	56
2	H	120/257 (47%)	114 (95%)	6 (5%)	0	100	100
2	J	120/257 (47%)	113 (94%)	7 (6%)	0	100	100
3	F	105/235 (45%)	99 (94%)	6 (6%)	0	100	100
3	I	105/235 (45%)	99 (94%)	6 (6%)	0	100	100
3	K	105/235 (45%)	98 (93%)	7 (7%)	0	100	100
3	L	105/235 (45%)	99 (94%)	6 (6%)	0	100	100
All	All	1525/2828 (54%)	1431 (94%)	93 (6%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	G	100	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/188 (77%)	139 (96%)	5 (4%)	36	63
1	B	147/188 (78%)	136 (92%)	11 (8%)	13	44
1	C	136/188 (72%)	124 (91%)	12 (9%)	10	38
1	D	141/188 (75%)	124 (88%)	17 (12%)	5	25
2	E	101/216 (47%)	89 (88%)	12 (12%)	5	26
2	G	101/216 (47%)	92 (91%)	9 (9%)	9	38
2	H	101/216 (47%)	87 (86%)	14 (14%)	3	21
2	J	101/216 (47%)	87 (86%)	14 (14%)	3	21
3	F	89/204 (44%)	87 (98%)	2 (2%)	52	73

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	I	89/204 (44%)	86 (97%)	3 (3%)	37 64
3	K	89/204 (44%)	78 (88%)	11 (12%)	4 24
3	L	89/204 (44%)	84 (94%)	5 (6%)	21 53
All	All	1328/2432 (55%)	1213 (91%)	115 (9%)	10 39

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	26	SER
3	L	31	SER
2	H	19	LYS
3	K	107	ILE
3	K	21	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	501	ASN
1	C	493	GLN
1	D	498	GLN
2	E	39	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	167/215 (77%)	0.48	14 (8%) 11 10	85, 136, 214, 248	0
1	B	172/215 (80%)	0.47	13 (7%) 13 11	83, 140, 207, 267	0
1	C	157/215 (73%)	0.46	11 (7%) 16 13	81, 131, 200, 287	0
1	D	163/215 (75%)	0.54	14 (8%) 10 9	64, 137, 202, 283	0
2	E	122/257 (47%)	0.36	3 (2%) 57 51	72, 120, 172, 250	0
2	G	122/257 (47%)	0.55	7 (5%) 23 20	76, 122, 182, 202	0
2	H	122/257 (47%)	0.46	7 (5%) 23 20	62, 117, 172, 198	0
2	J	122/257 (47%)	0.45	6 (4%) 29 26	75, 120, 179, 209	0
3	F	107/235 (45%)	0.58	9 (8%) 11 10	80, 127, 178, 200	0
3	I	107/235 (45%)	0.49	6 (5%) 24 21	88, 125, 171, 198	0
3	K	107/235 (45%)	0.43	5 (4%) 31 28	77, 134, 176, 234	0
3	L	107/235 (45%)	0.60	7 (6%) 18 15	87, 128, 179, 203	0
All	All	1575/2828 (55%)	0.49	102 (6%) 18 15	62, 130, 193, 287	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	373	SER	5.5
2	G	48	ILE	4.4
1	A	341	VAL	4.4
1	A	342	PHE	4.3
1	D	432	CYS	4.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.