



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 11:20 pm BST

PDB ID : 4XC3
Title : Crystal structure of human 4E10 Fab in complex with its peptide epitope on HIV-1 gp41; crystals cryoprotected with rac-glycerol 1-phosphate
Authors : Irimia, A.; Stanfield, R.L.; Wilson, I.A.
Deposited on : 2014-12-17
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	215	2% (Poor fit) 90% (0 outliers) 9% (1 outlier) 0% (2 outliers) 0% (3+ outliers)
2	H	230	6% (Poor fit) 88% (0 outliers) 8% (1 outlier) 0% (2 outliers) 0% (3+ outliers)
3	P	16	6% (Poor fit) 69% (0 outliers) 13% (1 outlier) 6% (2 outliers) 13% (3+ outliers)

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4E10 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	214	1746	1089	302	349	6	0	21	0


- Molecule 2 is a protein called 4E10 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	220	1725	1091	296	330	8	0	20	0

- Molecule 3 is a protein called MODIFIED FRAGMENT OF HIV-1 GLYCOPROTEIN (GP41) INCLUDING THE MPER REGION 671-683.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	P	14	157	110	24	23	0	2	0

- Molecule 4 is SN-GLYCEROL-1-PHOSPHATE (three-letter code: 1GP) (formula: C₃H₉O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	L	1	Total	C	O	P	0	0
			10	3	6	1		
4	H	1	Total	C	O	P	0	0
			6	1	4	1		

- Molecule 5 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	H	1	Total	C	O	0	0
			14	9	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	401	Total	O	0	0
			401	401		
6	H	352	Total	O	0	0
			352	352		
6	P	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4E10 Fab light chain



- Molecule 2: 4E10 Fab heavy chain



- Molecule 3: MODIFIED FRAGMENT OF HIV-1 GLYCOPROTEIN (GP41) INCLUDING THE MPER REGION 671-683



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.54Å 44.70Å 85.94Å 90.00° 113.22° 90.00°	Depositor
Resolution (Å)	32.80 – 1.63 32.80 – 1.63	Depositor EDS
% Data completeness (in resolution range)	97.1 (32.80-1.63) 97.1 (32.80-1.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.63Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.137 , 0.180 0.139 , 0.180	Depositor DCC
R_{free} test set	3360 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtrriage
Anisotropy	0.320	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4419	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 41H, UNL, 1GP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.35	0/1836	0.53	0/2486
2	H	0.35	0/1808	0.52	0/2464
3	P	0.26	0/152	0.42	0/209
All	All	0.35	0/3796	0.52	0/5159

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1746	0	1752	14	0
2	H	1725	0	1746	15	0
3	P	157	0	132	4	0
4	H	6	0	0	0	0
4	L	10	0	7	2	0
5	H	14	0	0	0	0
6	H	352	0	0	3	0
6	L	401	0	0	3	0
6	P	8	0	0	1	0
All	All	4419	0	3637	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:73[B]:ARG:NH2	6:H:401:HOH:O	2.13	0.81
2:H:73[B]:ARG:NH1	6:H:402:HOH:O	2.22	0.72
2:H:67[B]:ILE:HD12	2:H:80:LEU:HD11	1.70	0.72
2:H:83[B]:ARG:HG3	2:H:85:GLU:HG2	1.74	0.70
3:P:671[B]:ASN:ND2	6:P:701:HOH:O	2.25	0.70
2:H:34:LEU:HD21	2:H:92[B]:CYS:SG	2.37	0.64
3:P:671[A]:ASN:HD21	3:P:673:41H:H7	1.60	0.63
2:H:83[A]:ARG:NH1	6:H:404:HOH:O	2.29	0.63
2:H:73[B]:ARG:HH21	2:H:73[B]:ARG:HB2	1.69	0.57
1:L:146[B]:VAL:HG12	1:L:196:VAL:HG22	1.85	0.57
1:L:143:GLU:HG2	4:L:301:1GP:H31	1.87	0.57
2:H:67[B]:ILE:HD11	2:H:82:LEU:HD13	1.87	0.56
2:H:83[B]:ARG:HG2	2:H:86:ASP:OD2	2.06	0.56
1:L:198:HIS:HA	4:L:301:1GP:H32	1.87	0.56
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.88	0.56
2:H:63:PHE:HB3	2:H:67[B]:ILE:HG13	1.90	0.54
2:H:194:PRO:HG2	2:H:197:SER:HB3	1.92	0.51
1:L:32[B]:LYS:HD2	1:L:92:GLY:HA2	1.94	0.50
1:L:142[B]:ARG:HD3	1:L:163:VAL:HG11	1.95	0.48
2:H:205[A]:THR:HG23	2:H:222:LYS:HE3	1.94	0.48
3:P:671[A]:ASN:HB3	3:P:674:ASP:OD2	2.14	0.47
1:L:142[B]:ARG:HD2	1:L:142[B]:ARG:O	2.17	0.45
1:L:45[A]:ARG:NH1	6:L:412:HOH:O	2.49	0.45
2:H:34:LEU:HD11	2:H:92[B]:CYS:SG	2.57	0.45
1:L:142[B]:ARG:NH1	6:L:413:HOH:O	2.49	0.44
1:L:39:ARG:NH1	6:L:407:HOH:O	2.39	0.44
1:L:54[A]:ARG:HG3	1:L:58:VAL:HB	1.99	0.44
1:L:30:ASN:HB2	1:L:32[B]:LYS:HE3	1.99	0.43
2:H:83[B]:ARG:HG3	2:H:85:GLU:CG	2.46	0.43
1:L:134[B]:CYS:SG	1:L:177:SER:HB3	2.59	0.42
2:H:119:PRO:HB3	2:H:147:TYR:HB3	2.03	0.41
1:L:93:GLN:HA	3:P:671[B]:ASN:OD1	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	233/215 (108%)	227 (97%)	6 (3%)	0	100	100
2	H	236/230 (103%)	234 (99%)	1 (0%)	1 (0%)	34	15
3	P	12/16 (75%)	12 (100%)	0	0	100	100
All	All	481/461 (104%)	473 (98%)	7 (2%)	1 (0%)	47	26

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	100	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	205/185 (111%)	202 (98%)	3 (2%)	65	42
2	H	197/189 (104%)	195 (99%)	2 (1%)	76	59
3	P	14/15 (93%)	12 (86%)	2 (14%)	3	0
All	All	416/389 (107%)	409 (98%)	7 (2%)	73	36

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	70[A]	ASP
1	L	70[B]	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	105	GLU
2	H	43[A]	ARG
2	H	43[B]	ARG
3	P	671[A]	ASN
3	P	671[B]	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	41H	P	673	3	11,12,13	2.21	3 (27%)	12,15,17	1.53	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	41H	P	673	3	-	3/9/10/12	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	673	41H	CG-CB	4.95	1.58	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	673	41H	CW-CB	-4.58	1.44	1.53
3	P	673	41H	CE1-CD1	-2.23	1.34	1.38

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	673	41H	CW-CB-CA	4.11	117.33	111.34

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	673	41H	O-C-CA-CB
3	P	673	41H	C-CA-CB-CW
3	P	673	41H	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	P	673	41H	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is unknown - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	1GP	H	301	-	5,5,9	0.55	0	7,7,12	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	1GP	L	301	-	9,9,9	0.60	0	11,12,12	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1GP	H	301	-	-	0/1/3/8	-
4	1GP	L	301	-	-	2/8/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	301	1GP	C1-O1P-P-O2P
4	L	301	1GP	C1-O1P-P-O4P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	1GP	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	214/215 (99%)	-0.42	4 (1%) 66 67	12, 19, 33, 58	0
2	H	220/230 (95%)	-0.02	13 (5%) 22 20	13, 18, 47, 72	0
3	P	13/16 (81%)	0.58	1 (7%) 13 11	17, 26, 55, 58	0
All	All	447/461 (96%)	-0.19	18 (4%) 38 36	12, 18, 42, 72	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	100	TRP	13.0
2	H	100(B)	TRP	9.0
2	H	99	GLY	7.8
2	H	100(A)	GLY	7.6
2	H	100(C)	LEU	5.9
2	H	199	GLY	4.9
3	P	681	TYR	4.9
1	L	126	LYS	3.2
2	H	200	THR	3.1
1	L	212	GLY	2.9
2	H	43[A]	ARG	2.8
2	H	196	SER	2.8
2	H	195	SER	2.3
1	L	213	GLU	2.2
2	H	197	SER	2.2
2	H	143	LEU	2.1
1	L	203[A]	SER	2.0
2	H	26	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	41H	P	673	12/13	0.96	0.08	15,18,21,21	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	1GP	H	301	6/10	0.80	0.33	43,52,72,89	0
5	UNL	H	302	14/-	0.85	0.20	43,50,61,62	0
4	1GP	L	301	10/10	0.87	0.18	28,45,59,64	10

6.5 Other polymers [i](#)

There are no such residues in this entry.