



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 15, 2023 – 06:01 PM EDT

PDB ID : 1XDL  
Title : Structure of human aldolase B associated with hereditary fructose intolerance (A149P), at 277K  
Authors : Malay, A.D.; Allen, K.N.; Tolan, D.R.  
Deposited on : 2004-09-07  
Resolution : 3.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	
1	B	365	
1	C	365	
1	D	365	
1	W	365	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	X	365	
1	Y	365	
1	Z	365	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-bisphosphate aldolase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	303	2331	1469	411	439	12	0	0	0
1	B	299	2298	1448	405	433	12	0	0	0
1	C	329	2527	1589	446	479	13	0	0	0
1	D	297	2282	1436	403	431	12	0	0	0
1	W	296	2274	1431	400	431	12	0	0	0
1	X	329	2523	1584	445	481	13	0	0	0
1	Y	302	2318	1460	408	438	12	0	0	0
1	Z	301	2318	1461	408	437	12	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	cloning artifact	UNP P05062
A	0	SER	-	cloning artifact	UNP P05062
A	149	PRO	ALA	engineered mutation	UNP P05062
B	-1	GLY	-	cloning artifact	UNP P05062
B	0	SER	-	cloning artifact	UNP P05062
B	149	PRO	ALA	engineered mutation	UNP P05062
C	-1	GLY	-	cloning artifact	UNP P05062
C	0	SER	-	cloning artifact	UNP P05062
C	149	PRO	ALA	engineered mutation	UNP P05062
D	-1	GLY	-	cloning artifact	UNP P05062
D	0	SER	-	cloning artifact	UNP P05062
D	149	PRO	ALA	engineered mutation	UNP P05062
W	-1	GLY	-	cloning artifact	UNP P05062

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
W	0	SER	-	cloning artifact	UNP P05062
W	149	PRO	ALA	engineered mutation	UNP P05062
X	-1	GLY	-	cloning artifact	UNP P05062
X	0	SER	-	cloning artifact	UNP P05062
X	149	PRO	ALA	engineered mutation	UNP P05062
Y	-1	GLY	-	cloning artifact	UNP P05062
Y	0	SER	-	cloning artifact	UNP P05062
Y	149	PRO	ALA	engineered mutation	UNP P05062
Z	-1	GLY	-	cloning artifact	UNP P05062
Z	0	SER	-	cloning artifact	UNP P05062
Z	149	PRO	ALA	engineered mutation	UNP P05062

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	W	1	Total O S 5 4 1	0	0
2	X	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	Y	1	Total	O	S	0	0
			5	4	1		
2	Y	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		
3	B	4	Total	O	0	0
			4	4		
3	C	3	Total	O	0	0
			3	3		
3	D	2	Total	O	0	0
			2	2		
3	W	2	Total	O	0	0
			2	2		
3	X	1	Total	O	0	0
			1	1		
3	Y	2	Total	O	0	0
			2	2		
3	Z	3	Total	O	0	0
			3	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Fructose-bisphosphate aldolase B


Chain A: 





#### • Molecule 1: Fructose-bisphosphate aldolase B

Chain B: 








• Molecule 1: Fructose-bisphosphate aldolase B





• Molecule 1: Fructose-bisphosphate aldolase B













- Molecule 1: Fructose-bisphosphate aldolase B




- Molecule 1: Fructose-bisphosphate aldolase B





- Molecule 1: Fructose-bisphosphate aldolase B

Chain Y:  26% 49% 7% 17%



• Molecule 1: Fructose-bisphosphate aldolase B

Chain Z:  29% 46% 7% 18%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.48Å 153.51Å 186.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.15 – 3.00 20.15 – 2.50	Depositor EDS
% Data completeness (in resolution range)	78.1 (20.15-3.00) 69.9 (20.15-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.16 (at 2.50Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.304 , 0.349 0.299 , 0.340	Depositor DCC
$R_{free}$ test set	7001 reflections (5.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.5	Xtrriage
Anisotropy	0.708	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 18.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.458 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	18932	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/2365	0.67	0/3190
1	B	0.48	0/2333	0.71	0/3146
1	C	0.48	0/2566	0.72	1/3465 (0.0%)
1	D	0.42	0/2315	0.69	0/3121
1	W	0.42	0/2306	0.67	0/3108
1	X	0.48	1/2563 (0.0%)	0.74	2/3464 (0.1%)
1	Y	0.46	0/2353	0.71	0/3174
1	Z	0.40	0/2351	0.66	0/3170
All	All	0.44	1/19152 (0.0%)	0.70	3/25838 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
1	W	0	1
1	Y	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	70	ASN	CB-CG	-5.20	1.39	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	70	ASN	CB-CA-C	-10.72	88.96	110.40
1	X	66	ASP	N-CA-C	5.65	126.26	111.00
1	C	66	ASP	N-CA-C	5.43	125.65	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	213	TYR	Sidechain
1	C	213	TYR	Sidechain
1	C	84	TYR	Sidechain
1	W	213	TYR	Sidechain
1	Y	213	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2331	0	2359	214	0
1	B	2298	0	2313	247	0
1	C	2527	0	2555	262	0
1	D	2282	0	2301	220	0
1	W	2274	0	2293	201	0
1	X	2523	0	2538	259	0
1	Y	2318	0	2338	242	0
1	Z	2318	0	2345	220	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	10	0	0	0	0
2	W	5	0	0	0	0
2	X	5	0	0	0	0
2	Y	10	0	0	0	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
3	C	3	0	0	2	0
3	D	2	0	0	0	0
3	W	2	0	0	0	0
3	X	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	Y	2	0	0	0	0
3	Z	3	0	0	1	0
All	All	18932	0	19042	1839	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 48.

The worst 5 of 1839 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:W:303:ARG:H	1:W:303:ARG:HD2	1.05	1.19
1:B:303:ARG:H	1:B:303:ARG:HD2	1.02	1.17
1:Z:228:LEU:HG	1:Z:230:PRO:HD3	1.26	1.17
1:C:303:ARG:H	1:C:303:ARG:HD2	1.03	1.17
1:C:149:PRO:HG3	1:C:170:LEU:HD11	1.20	1.17

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/365 (80%)	223 (76%)	55 (19%)	15 (5%)	2	12
1	B	289/365 (79%)	209 (72%)	59 (20%)	21 (7%)	1	5
1	C	321/365 (88%)	244 (76%)	57 (18%)	20 (6%)	1	8
1	D	287/365 (79%)	218 (76%)	51 (18%)	18 (6%)	1	7
1	W	286/365 (78%)	222 (78%)	47 (16%)	17 (6%)	1	9
1	X	323/365 (88%)	246 (76%)	56 (17%)	21 (6%)	1	7
1	Y	292/365 (80%)	214 (73%)	60 (20%)	18 (6%)	1	8
1	Z	291/365 (80%)	219 (75%)	55 (19%)	17 (6%)	1	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2382/2920 (82%)	1795 (75%)	440 (18%)	147 (6%)	<b>1</b> <b>8</b>

5 of 147 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	SER
1	A	149	PRO
1	A	233	VAL
1	B	67	SER
1	B	88	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	247/292 (85%)	230 (93%)	17 (7%)	<b>15</b> 48
1	B	242/292 (83%)	225 (93%)	17 (7%)	<b>15</b> 47
1	C	269/292 (92%)	247 (92%)	22 (8%)	<b>11</b> 39
1	D	241/292 (82%)	220 (91%)	21 (9%)	<b>10</b> 37
1	W	240/292 (82%)	223 (93%)	17 (7%)	<b>14</b> 46
1	X	268/292 (92%)	243 (91%)	25 (9%)	<b>9</b> 33
1	Y	245/292 (84%)	227 (93%)	18 (7%)	<b>14</b> 44
1	Z	245/292 (84%)	229 (94%)	16 (6%)	<b>17</b> 50
All	All	1997/2336 (86%)	1844 (92%)	153 (8%)	<b>13</b> 42

5 of 153 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	254	THR
1	Z	173	TYR
1	Y	7	LEU
1	Y	203	TYR
1	Z	295	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 115 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	341	GLN
1	Z	200	HIS
1	W	334	ASN
1	Z	179	GLN
1	Y	334	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	X	364	-	4,4,4	0.28	0	6,6,6	0.13	0
2	SO4	W	364	-	4,4,4	0.26	0	6,6,6	0.10	0
2	SO4	B	364	-	4,4,4	0.32	0	6,6,6	0.13	0
2	SO4	Y	365	-	4,4,4	0.34	0	6,6,6	0.08	0
2	SO4	Y	364	-	4,4,4	0.31	0	6,6,6	0.08	0
2	SO4	C	365	-	4,4,4	0.28	0	6,6,6	0.11	0
2	SO4	A	364	-	4,4,4	0.31	0	6,6,6	0.09	0
2	SO4	C	364	-	4,4,4	0.29	0	6,6,6	0.17	0



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	303/365 (83%)	-0.35	0 <a href="#">100</a> <a href="#">100</a>	15, 74, 94, 94	0
1	B	299/365 (81%)	-0.37	0 <a href="#">100</a> <a href="#">100</a>	1, 54, 89, 94	0
1	C	329/365 (90%)	-0.39	1 (0%) <a href="#">94</a> <a href="#">84</a>	9, 65, 94, 94	0
1	D	297/365 (81%)	-0.36	1 (0%) <a href="#">94</a> <a href="#">84</a>	5, 69, 94, 94	0
1	W	296/365 (81%)	-0.38	0 <a href="#">100</a> <a href="#">100</a>	7, 68, 94, 94	0
1	X	329/365 (90%)	-0.45	0 <a href="#">100</a> <a href="#">100</a>	7, 68, 94, 94	0
1	Y	302/365 (82%)	-0.42	1 (0%) <a href="#">94</a> <a href="#">84</a>	1, 55, 89, 94	0
1	Z	301/365 (82%)	-0.35	1 (0%) <a href="#">94</a> <a href="#">84</a>	16, 74, 94, 94	0
All	All	2456/2920 (84%)	-0.38	4 (0%) <a href="#">95</a> <a href="#">87</a>	1, 66, 94, 94	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	192	PRO	3.6
1	Y	150	VAL	3.1
1	D	320	LYS	2.5
1	Z	314	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	B	364	5/5	0.80	0.13	93,93,93,93	0
2	SO4	C	364	5/5	0.88	0.12	93,93,93,93	0
2	SO4	C	365	5/5	0.88	0.15	93,93,93,93	0
2	SO4	X	364	5/5	0.88	0.13	93,93,93,93	0
2	SO4	W	364	5/5	0.89	0.14	93,93,93,93	0
2	SO4	Y	365	5/5	0.91	0.17	93,93,93,93	0
2	SO4	Y	364	5/5	0.93	0.12	93,93,93,93	0
2	SO4	A	364	5/5	0.94	0.13	93,93,93,93	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.