wwPDB EM Validation Summary Report (i)

Oct 27, 2026 – 04:19 PM JST

PDB ID	:	9XFP / pdb_00009xfp
EMDB ID	:	EMD-66835
Title	:	Cryo-EM structure of Retron-ATPaseHNH
Authors	:	Hu, C.; Zhang, S.; Wang, Q.
Deposited on	:	2025-10-29
Resolution	:	3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	67% 16% 17%
1	B	543	5% 73% 18% 9%
1	C	543	6% 85% 8% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	D	543	71% 15% 14%
2	E	227	10% 61% 33% 6%
2	F	227	10% 90%
3	G	317	85% 12%
4	H	84	5% 49% 51%
5	I	39	41% 38% 18%

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 22335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AAA family ATPase.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	453	Total	C	N	O	S	0	0
			3651	2348	601	698	4		
1	B	493	Total	C	N	O	S	0	0
			3891	2492	650	744	5		
1	C	510	Total	C	N	O	S	0	0
			4024	2575	670	774	5		
1	D	467	Total	C	N	O	S	0	0
			3784	2429	628	723	4		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	GLY	GLU	conflict	UNP A0AAD2V6K7
B	155	GLY	GLU	conflict	UNP A0AAD2V6K7
C	155	GLY	GLU	conflict	UNP A0AAD2V6K7
D	155	GLY	GLU	conflict	UNP A0AAD2V6K7

- Molecule 2 is a protein called TIGR02646 family protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	214	Total	C	N	O	S	0	0
			1719	1076	307	322	14		
2	F	23	Total	C	N	O		0	0
			114	68	23	23			

- Molecule 3 is a protein called RNA-directed DNA polymerase.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	310	Total	C	N	O	S	0	0
			2511	1624	433	442	12		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	312	GLY	-	expression tag	UNP A0AAD2V6H6
G	313	GLY	-	expression tag	UNP A0AAD2V6H6
G	314	GLY	-	expression tag	UNP A0AAD2V6H6
G	315	SER	-	expression tag	UNP A0AAD2V6H6
G	316	GLY	-	expression tag	UNP A0AAD2V6H6
G	317	GLY	-	expression tag	UNP A0AAD2V6H6


- Molecule 4 is a DNA chain called DNA (97-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	84	Total	C	N	O	P	0	0
			1718	816	318	500	84		

- Molecule 5 is a RNA chain called RNA (69-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	39	Total	C	N	O	P	0	0
			829	370	141	279	39		

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
6	B	1	Total	C	N	O	P	0
			31	10	5	13	3	
6	C	1	Total	C	N	O	P	0
			31	10	5	13	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
6	D	1	Total	C	N	O	P	0
			31	10	5	13	3	

- Molecule 7 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
7	E	1	Total	Zn	0
			1	1	


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AAA family ATPase

Chain A:






• Molecule 1: AAA family ATPase


Chain B:









• Molecule 1: AAA family ATPase




• Molecule 1: AAA family ATPase



• Molecule 2: TIGR02646 family protein






• Molecule 2: TIGR02646 family protein

Chain F:






• Molecule 3: RNA-directed DNA polymerase

Chain G:






• Molecule 4: DNA (97-MER)


Chain H:






• Molecule 5: RNA (69-MER)

Chain I:





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	243379	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2800	Depositor
Magnification	Not provided	
Image detector	TFS FALCON 4i (4k x 4k)	Depositor
Maximum map value	1.872	Depositor
Minimum map value	-0.877	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.029	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	376.32, 376.32, 376.32	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.98, 0.98, 0.98	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.13	0/3721	0.32	0/5045
1	B	0.12	0/3963	0.32	0/5378
1	C	0.12	0/4095	0.28	0/5552
1	D	0.12	0/3856	0.31	0/5223
2	E	0.13	0/1750	0.41	0/2345
2	F	0.06	0/113	0.15	0/156
3	G	0.15	0/2564	0.34	0/3448
4	H	0.73	0/1927	0.91	1/2970 (0.0%)
5	I	0.39	0/925	0.89	1/1439 (0.1%)
All	All	0.25	0/22914	0.45	2/31556 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	I	32	G	P-O3'-C3'	5.56	126.38	119.70
4	H	35	DT	N3-C4-O4	5.44	123.16	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3651	0	3634	68	0
1	B	3891	0	3784	63	0
1	C	4024	0	3944	33	0
1	D	3784	0	3795	57	0
2	E	1719	0	1659	58	0
2	F	114	0	55	0	0
3	G	2511	0	2595	25	0
4	H	1718	0	943	37	0
5	I	829	0	416	18	0
6	B	31	0	12	2	0
6	C	31	0	12	1	0
6	D	31	0	12	2	0
7	E	1	0	0	0	0
All	All	22335	0	20861	337	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 337 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:LEU:CD2	1:A:413:PHE:HB2	1.67	1.23
2:E:92:CYS:HB2	2:E:98:CYS:SG	2.04	0.96
4:H:99:DC:H42	5:I:64:G:H1	1.10	0.96
1:A:68:LEU:HD23	1:A:413:PHE:HB2	1.49	0.91
1:A:68:LEU:HD22	1:A:413:PHE:HB2	1.53	0.90

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/543 (82%)	431 (96%)	16 (4%)	0	100	100
1	B	487/543 (90%)	465 (96%)	22 (4%)	0	100	100
1	C	504/543 (93%)	484 (96%)	20 (4%)	0	100	100
1	D	463/543 (85%)	441 (95%)	22 (5%)	0	100	100
2	E	210/227 (92%)	199 (95%)	11 (5%)	0	100	100
2	F	21/227 (9%)	21 (100%)	0	0	100	100
3	G	308/317 (97%)	294 (96%)	14 (4%)	0	100	100
All	All	2440/2943 (83%)	2335 (96%)	105 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/500 (82%)	410 (100%)	0	100	100
1	B	421/500 (84%)	418 (99%)	3 (1%)	76	78
1	C	438/500 (88%)	436 (100%)	2 (0%)	81	80
1	D	428/500 (86%)	425 (99%)	3 (1%)	76	78
2	E	185/207 (89%)	183 (99%)	2 (1%)	65	74
3	G	279/284 (98%)	276 (99%)	3 (1%)	65	74
All	All	2161/2491 (87%)	2148 (99%)	13 (1%)	76	79

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	520	ASP
2	E	155	LEU
3	G	291	LEU
3	G	245	LEU
3	G	250	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	7	GLN
3	G	46	GLN
3	G	310	ASN
2	E	208	ASN
1	D	75	ASN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	I	39/39 (100%)	12 (30%)	2 (5%)

5 of 12 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	I	32	G
5	I	33	G
5	I	37	A
5	I	38	U
5	I	41	G

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
5	I	31	C
5	I	32	G

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	ATP	B	600	-	29,33,33	0.28	0	44,52,52	0.45	1 (2%)
6	ATP	D	601	-	29,33,33	0.32	0	44,52,52	0.48	0
6	ATP	C	600	-	29,33,33	0.30	0	44,52,52	0.48	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ATP	B	600	-	-	5/22/38/38	0/3/3/3
6	ATP	D	601	-	-	6/22/38/38	0/3/3/3
6	ATP	C	600	-	-	5/22/38/38	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
6	C	600	ATP	PB-O3B-PG	2.07	139.92	132.83
6	B	600	ATP	PB-O3B-PG	2.04	139.82	132.83

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	600	ATP	C3'-C4'-C5'-O5'
6	D	601	ATP	C5'-O5'-PA-O3A
6	D	601	ATP	O4'-C4'-C5'-O5'
6	D	601	ATP	C3'-C4'-C5'-O5'
6	B	600	ATP	O4'-C4'-C5'-O5'


There are no ring outliers.

3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	600	ATP	2	0
6	D	601	ATP	2	0
6	C	600	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand ATP B 600




Ligand ATP D 601




Bond lengths



Bond angles




Torsions



Rings

Ligand ATP C 600



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.


6 Map visualisation

This section contains visualisations of the EMDB entry EMD-66835. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192



Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



X Index: 186




Y Index: 177




Z Index: 240


6.3.2 Raw map



X Index: 176



Y Index: 192




Z Index: 244


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)


6.5.1 Primary map



X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate (i)



The volume at the recommended contour level is 300 nm^3; this corresponds to an approximate mass of 271 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.286 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.286 Å^{-1}

8.2 Resolution estimates (i)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.13	4.64	4.19

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.13 differs from the reported value 3.5 by more than 10 %


9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-66835 and PDB model 9XFP. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.


9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion (i)



At the recommended contour level, 92% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8990	0.4510
A	0.9390	0.4620
B	0.8910	0.4340
C	0.9030	0.4850
D	0.8950	0.4620
E	0.6970	0.3310
F	0.7110	0.3520
G	0.9770	0.5240
H	0.8890	0.3740
I	0.9880	0.4710

