



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 06:54 AM EDT

PDB ID : 4XI6  
Title : Crystal structure of the MZM-REP domains of Mind bomb 1  
Authors : McMillan, B.J.; Blacklow, S.C.  
Deposited on : 2015-01-06  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5748 atoms, of which 2714 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase MIB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	363	5550	1768	2714	517	534	17	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	SER	-	expression tag	UNP Q86YT6

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	186	Total	O	0	0
			186	186		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 ubiquitin-protein ligase MIB1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.08Å 73.08Å 170.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.94 – 2.04 44.94 – 1.99	Depositor EDS
% Data completeness (in resolution range)	93.6 (44.94-2.04) 93.6 (44.94-1.99)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.208 , 0.248 0.211 , 0.250	Depositor DCC
$R_{free}$ test set	2000 reflections (6.58%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 59.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5748	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	1/2901 (0.0%)	0.51	0/3931

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	141	ARG	CZ-NH2	-5.49	1.25	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2836	2714	2723	16	8
2	A	2	0	0	0	0
3	A	10	0	0	0	0
4	A	186	0	0	0	0
All	All	3034	2714	2723	16	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:CYS:SG	1:A:295:VAL:HG12	2.06	0.95
1:A:63:SER:O	1:A:65:ALA:N	2.03	0.90
1:A:141:ARG:HD2	1:A:355:HIS:CD2	2.45	0.52
1:A:388:THR:HG22	1:A:389:SER:H	1.76	0.49
1:A:141:ARG:HD3	1:A:141:ARG:O	2.13	0.48
1:A:118:HIS:CG	1:A:119:HIS:H	2.32	0.47
1:A:294:ILE:HD12	1:A:294:ILE:N	2.29	0.47
1:A:286:CYS:SG	1:A:303:ARG:HD2	2.56	0.45
1:A:169:ASP:OD1	1:A:169:ASP:N	2.50	0.45
1:A:55:GLY:HA3	1:A:216:LYS:HD2	1.99	0.44
1:A:390:TRP:HB2	1:A:392:TYR:CE2	2.53	0.43
1:A:275:MET:HG2	1:A:304:TRP:CD2	2.53	0.43
1:A:74:ALA:N	1:A:75:PRO:CD	2.82	0.42
1:A:63:SER:C	1:A:65:ALA:N	2.71	0.42
1:A:89:ARG:HG3	1:A:258:LEU:HD23	2.02	0.42
1:A:278:THR:HG22	1:A:298:TYR:CZ	2.55	0.41

All (8) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:LEU:HD21	1:A:264:LEU:HD22[7_644]	0.00	1.60
1:A:264:LEU:HD22	1:A:264:LEU:HD23[7_644]	0.72	0.88
1:A:264:LEU:HD23	1:A:264:LEU:HD23[7_644]	0.81	0.79
1:A:264:LEU:CG	1:A:264:LEU:CD2[7_644]	1.54	0.66
1:A:264:LEU:CD2	1:A:264:LEU:HD22[7_644]	0.97	0.63
1:A:264:LEU:CD2	1:A:264:LEU:HD23[7_644]	0.97	0.63
1:A:264:LEU:CD2	1:A:264:LEU:HD21[7_644]	0.97	0.63
1:A:264:LEU:HD21	1:A:264:LEU:HD23[7_644]	1.33	0.27

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/380 (94%)	341 (95%)	15 (4%)	3 (1%)	19 10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	GLY
1	A	331	SER
1	A	314	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	301/314 (96%)	299 (99%)	2 (1%)	84 84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	A	203	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	504	-	4,4,4	0.17	0	6,6,6	0.14	0
3	SO4	A	503	-	4,4,4	0.14	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/380 (95%)	1.07	65 (17%) <b>1</b> <b>1</b>	10, 30, 90, 113	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	310	VAL	9.7
1	A	260	ILE	9.3
1	A	283	GLY	8.7
1	A	309	ALA	8.6
1	A	286	CYS	8.2
1	A	271	TRP	7.8
1	A	252	VAL	7.8
1	A	264	LEU	7.7
1	A	297	GLN	7.1
1	A	292	HIS	6.6
1	A	291	ASP	6.2
1	A	306	PHE	6.2
1	A	256	LEU	6.0
1	A	287	GLY	5.9
1	A	308	PRO	5.9
1	A	298	TYR	5.8
1	A	263	SER	5.8
1	A	288	ILE	5.5
1	A	257	ASP	5.2
1	A	296	VAL	5.1
1	A	267	GLY	5.0
1	A	301	GLY	4.6
1	A	284	THR	4.6
1	A	313	LYS	4.5
1	A	303	ARG	4.4
1	A	270	GLY	4.4
1	A	276	PHE	4.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	278	THR	4.2
1	A	299	PRO	4.2
1	A	294	ILE	4.2
1	A	266	HIS	4.1
1	A	262	GLN	4.1
1	A	295	VAL	3.9
1	A	254	ILE	3.9
1	A	282	THR	3.9
1	A	272	THR	3.8
1	A	331	SER	3.8
1	A	258	LEU	3.8
1	A	307	ASN	3.6
1	A	259	GLU	3.6
1	A	300	SER	3.5
1	A	265	GLN	3.4
1	A	275	MET	3.4
1	A	311	LEU	3.4
1	A	304	TRP	3.2
1	A	290	GLU	3.2
1	A	281	THR	3.1
1	A	305	THR	3.1
1	A	268	HIS	3.1
1	A	314	ALA	3.0
1	A	293	ASP	3.0
1	A	186	ALA	3.0
1	A	285	VAL	2.9
1	A	289	ASP	2.9
1	A	269	GLY	2.8
1	A	332	GLN	2.7
1	A	273	ASP	2.6
1	A	312	THR	2.5
1	A	167	ASP	2.4
1	A	251	LEU	2.2
1	A	170	GLY	2.2
1	A	400	VAL	2.1
1	A	261	VAL	2.1
1	A	166	GLU	2.0
1	A	255	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	504	5/5	0.98	0.16	23,27,28,29	0
3	SO4	A	503	5/5	0.99	0.17	15,15,16,17	0
2	ZN	A	501	1/1	0.99	0.14	22,22,22,22	0
2	ZN	A	502	1/1	1.00	0.16	18,18,18,18	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.