



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 11, 2023 – 12:29 AM EDT

PDB ID : 4XLR
Title : Crystal structure of *T.aquaticus* transcription initiation complex with CarD containing bubble promoter and RNA
Authors : Bae, B.; Darst, S.A.
Deposited on : 2015-01-13
Resolution : 4.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1014 (4.80-3.80)
Clashscore	141614	1077 (4.80-3.80)
Ramachandran outliers	138981	1029 (4.80-3.80)
Sidechain outliers	138945	1012 (4.80-3.80)
RSRZ outliers	127900	1075 (4.90-3.70)
RNA backbone	3102	1058 (5.60-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	 37% 31% • 28%
1	B	314	 42% 26% • 28%
1	G	314	 37% 31% • 28%
1	H	314	 42% 26% • 28%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	1119	 51% 43% 6%
2	I	1119	 % 51% 43% 6%
3	D	1524	 52% 41% 5%
3	J	1524	 49% 37% 10%
4	E	99	 55% 35% 6%
4	K	99	 56% 34% 6%
5	F	347	 % 59% 35% 6%
5	L	347	 % 57% 36% 6%
6	M	164	 3% 65% 29% ..
6	N	164	 3% 66% 27% ..
7	O	48	 8% 48% 52%
7	R	48	 2% 42% 58%
8	P	48	 8% 38% 63%
8	S	48	 2% 42% 54%
9	Q	4	 25% 25% 75%
9	T	4	 50% 50%

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 60854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	B	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	G	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			
1	H	227	Total	C	N	O	S	0	0	0
			1770	1130	303	334	3			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	1117	Total	C	N	O	S	0	0	0
			8762	5544	1558	1637	23			
2	I	1117	Total	C	N	O	S	0	0	0
			8762	5544	1558	1637	23			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	1490	Total	C	N	O	S	0	0	0
			11761	7439	2088	2196	38			
3	J	1367	Total	C	N	O	S	0	0	0
			10779	6810	1923	2010	36			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	E	93	Total	C	N	O	S	0	0	0
			768	490	136	138	4			
4	K	93	Total	C	N	O	S	0	0	0
			768	490	136	138	4			

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	345	Total	C	N	O	S	0	0	0
			2787	1758	502	523	4			
5	L	345	Total	C	N	O	S	0	0	0
			2787	1758	502	523	4			

- Molecule 6 is a protein called CarD-like transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	M	158	Total	C	N	O	S	0	0	0
			1239	787	229	221	2			
6	N	158	Total	C	N	O	S	0	0	0
			1239	787	229	221	2			

- Molecule 7 is a DNA chain called DNA (48-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	O	48	Total	C	N	O	P	0	0	0
			988	472	182	287	47			
7	R	48	Total	C	N	O	P	0	0	0
			988	472	182	287	47			

- Molecule 8 is a DNA chain called DNA (48-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	P	48	Total	C	N	O	P	0	0	0
			985	471	183	284	47			
8	S	48	Total	C	N	O	P	0	0	0
			985	471	183	284	47			

- Molecule 9 is a RNA chain called RNA (5'-R(P*UP*CP*GP*A)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	Q	4	Total	C	N	O	P	0	0	0
			85	38	15	28	4			
9	T	4	Total	C	N	O	P	0	0	0
			85	38	15	28	4			

- Molecule 10 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	2	Total	Zn	0	0
			2	2		
10	J	2	Total	Zn	0	0
			2	2		

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	D	1	Total	Mg	0	0
			1	1		
11	J	1	Total	Mg	0	0
			1	1		



• Molecule 3: DNA-directed RNA polymerase subunit beta'




R75	T154	VAL	LEU	V355	M527	L619	K688	D847	V936	R1042	R1136	V1221
C76	D155	GLU	VAL	P356	T537	G620	V699	L850	Y937	M1045	K1136	G1222
V78	E156	LYS	VAL	E357	S538	K621	V700	L859	L941	M1048	S1138	V1223
S83	E150	ALA	GLY	Q362	D539	V623	L701	D859	I947	P1048	D1139	A1225
T84	L161	TYR	GLY	E365	L543	R628	L702	O861	T948	V1055	I1140	A1226
R86	K165	ARG	ILE	K366	Y544	S629	N703	D862	I949	P1056	L1144	T1234
R88	Q166	PRO	VAL	I367	I548	W630	A705	H863	G950	V1057	Y1145	L1144
Y88	E167	GLY	GLU	V368	I631	V632	T707	V864	I951	S1060	G1146	M1238
H82	T168	VAL	VAL	R87	N849	V633	L708	T865	D952	F1061	R1147	ARG
A96	Y169	LEU	GLN	I371	R550	G634	L708	Y868	D953	R1062	V1148	THR
A99	L171	ALA	PRU	D372	R553	G635	I713	L879	A954	L1068	L1149	PHE
A100	P172	GLU	LEU	F373	K554	Q636	Q714	I880	E958	L1068	G1157	HIS
H101	V175	GLY	ALA	E374	K554	L637	A715	E874	E959	L1068	R1158	THR
H102	D176	GLY	GLU	E375	K554	L637	A715	E874	E959	L1068	R1158	GLY
W103	A177	TYR	LYS	E376	G561	K638	F716	T875	K960	S1072	R1159	VAL
F104	K180	PHE	LEU	I378	A562	G643	Q717	S876	I964	S1073	L1160	GLY
V105	D181	ARG	LEU	A379	I566	L644	V719	R879	E965	S1074	E1161	ALA
K106	V186	ALA	GLY	V385	I567	K646	L720	I880	E965	H1075	E1162	VAL
D107	L387	PRO	ARG	L387	K571	K647	E722	F882	Q976	A1077	S1167	GLY
I112	K187	GLY	HIS	E389	M572	A649	G723	A883	Q976	R1077	L1168	THR
G113	L191	VAL	THR	P390	M573	L650	Q724	I885	I974	K1079	E1169	ASP
T114	A192	VAL	MET	A391	L574	F653	Q727	V895	E975	L1086	V1171	ILE
L115	G194	GLU	THR	H386	A577	K653	L728	R796	Q976	R1087	H1172	GLY
L116	G194	GLU	THR	H386	A577	K653	L728	R796	Q976	R1087	H1172	GLY
D117	V195	LEU	ALA	V395	D579	P655	F730	K800	T984	A1077	L1174	ASP
L118	S197	ASP	GLU	V395	A580	F656	L731	G801	T984	D1090	I1175	ASP
T121	R198	LEU	ALA	K397	V581	L657	L731	G801	R988	Y1093	E1184	ILE
E122	L204	GLY	GLU	V400	I582	L658	F736	A807	L995	L1094	E1185	VAL
L123	Y205	GLY	GLU	V400	I582	L658	F736	A807	L995	L1094	E1185	VAL
E124	Q125	HIS	GLY	D406	D583	K659	A738	E810	V1003	R1096	V1189	THR
Q125	P208	LEU	ASP	V407	N584	K660	D739	E898	V1007	K1097	S1190	THR
V126	R209	ILE	GLU	E408	G585	M661	F740	L813	F1008	V1099	P1191	GLY
L127	R210	TYR	SER	V409	R587	E662	D743	E817	F1011	D1100	L1182	GLY
Y128	V211	LEU	ASP	T410	P590	F666	Q744	V821	E1012	V1101	C1194	THR
F129	V211	LEU	ASP	T411	T592	V670	M745	A822	E1013	A1102	C1194	THR
M130	R212	ARG	THR	T411	N593	K671	A746	A822	N1014	H1103	Q1195	THR
K131	V213	GLN	THR	T411	P594	A672	V747	A825	N1015	V1106	T1196	THR
Y132	L216	GLU	PHE	G418	P594	R675	H748	A825	P1016	V1107	R1197	THR
P137	ARG	GLU	LEU	D419	P599	M676	P750	W828	F1017	R1108	V1200	THR
A140	LYS	VAL	LEU	V420	L600	L603	L751	G831	F1017	R1108	C1201	THR
V141	GLU	VAL	LEU	L421	L514	L603	S752	R833	N1018	T1114	Q1202	THR
V145	ALA	ALA	ALA	A422	E515	L603	F754	E833	Y1021	T1115	Q1203	THR
P146	ARG	ARG	ARG	G425	W517	I606	A755	T834	Q1025	M1116	C1204	THR
V147	TYR	ARG	TYR	G425	P518	I606	A757	S835	A1028	L1122	D1208	THR
E148	ALA	LEU	PHE	K428	V519	K610	E758	R838	A1028	E1127	L1209	THR
Q151	LEU	LEU	LEU	V519	L520	V687	A759	L839	A1029	E1127	S1210	THR
L152	PRO	ILE	LEU	E430	P521	W688	R760	L839	R929	T1129	M1211	THR
L153	ALA	ALA	ALA	V347	P522	R613	R761	L839	Q1037	T1129	A1212	THR
	LEU	GLY	GLY	P349	D523	F614	I761	L931	L1038	R1130	R1213	THR
	SER	THR	MET	R434	L524	M617	Q762	L931	C1039	T1131	P1214	THR
	ALA	ALA	ALA	W351	M525	E693	M763	L934	G1040	L1132	V1215	THR
	TRP	PRO	TRP	E436	V694	L764		K935	M1041		P1306	THR




- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain E: 55% 35% 6%




- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain K: 56% 34% 6%



- Molecule 5: RNA polymerase sigma factor SigA

Chain F: 59% 35% 6%




- Molecule 5: RNA polymerase sigma factor SigA


Chain L: 57% 36% 6%



- Molecule 8: DNA (48-MER)



- Molecule 8: DNA (48-MER)



- Molecule 9: RNA (5'-R(P*UP*CP*GP*A)-3')



- Molecule 9: RNA (5'-R(P*UP*CP*GP*A)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	289.84Å 289.84Å 536.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.56 – 4.30 39.56 – 4.30	Depositor EDS
% Data completeness (in resolution range)	94.8 (39.56-4.30) 94.8 (39.56-4.30)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.11 (at 4.28Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, R_{free}	0.275 , 0.310 0.275 , 0.310	Depositor DCC
R_{free} test set	7337 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	165.1	Xtrriage
Anisotropy	0.026	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 123.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.19$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	60854	wwPDB-VP
Average B, all atoms (Å ²)	179.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1804	0.52	0/2455
1	B	0.27	0/1804	0.49	0/2455
1	G	0.30	0/1804	0.52	0/2455
1	H	0.27	0/1804	0.49	0/2455
2	C	0.29	0/8929	0.51	1/12074 (0.0%)
2	I	0.29	0/8929	0.51	1/12074 (0.0%)
3	D	0.29	0/11963	0.50	0/16165
3	J	0.28	0/10959	0.49	0/14802
4	E	0.27	0/783	0.53	0/1054
4	K	0.27	0/783	0.53	0/1054
5	F	0.34	0/2829	0.54	0/3804
5	L	0.33	0/2829	0.54	0/3804
6	M	0.35	0/1267	0.55	0/1719
6	N	0.35	0/1267	0.55	0/1719
7	O	0.59	0/1109	0.92	0/1712
7	R	0.56	0/1109	0.92	0/1712
8	P	0.64	0/1106	0.88	0/1706
8	S	0.61	0/1106	0.90	2/1706 (0.1%)
9	Q	0.24	0/94	0.71	0/144
9	T	0.24	0/94	0.76	0/144
All	All	0.33	0/62372	0.55	4/85213 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	2
2	I	0	2
3	D	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
3	J	0	1
6	M	0	2
6	N	0	2
All	All	0	10

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	S	23	DA	O5'-P-OP1	-6.61	99.75	105.70
2	C	242	LEU	CA-CB-CG	5.68	128.36	115.30
2	I	242	LEU	CA-CB-CG	5.57	128.10	115.30
8	S	22	DT	OP1-P-O3'	5.13	116.48	105.20

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	360	VAL	Peptide
2	C	71	TYR	Mainchain
3	D	1270	ALA	Peptide
2	I	360	VAL	Peptide
2	I	71	TYR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1770	0	1799	87	0
1	B	1770	0	1799	66	0
1	G	1770	0	1799	88	0
1	H	1770	0	1799	65	0
2	C	8762	0	8854	435	0
2	I	8762	0	8854	436	0
3	D	11761	0	11976	537	0
3	J	10779	0	10993	490	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	768	0	784	38	0
4	K	768	0	784	36	0
5	F	2787	0	2866	122	0
5	L	2787	0	2866	127	0
6	M	1239	0	1259	38	0
6	N	1239	0	1259	39	0
7	O	988	0	544	30	0
7	R	988	0	544	38	0
8	P	985	0	543	36	0
8	S	985	0	543	30	0
9	Q	85	0	43	1	0
9	T	85	0	43	2	0
10	D	2	0	0	0	0
10	J	2	0	0	0	0
11	D	1	0	0	0	0
11	J	1	0	0	0	0
All	All	60854	0	59951	2363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 2363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:73:CYS:HB3	3:D:76:CYS:SG	1.97	1.04
3:J:73:CYS:HB3	3:J:76:CYS:SG	1.97	1.04
3:D:105:VAL:HA	3:D:112:ILE:HD11	1.55	0.89
3:D:412:GLY:HA2	3:D:434:ARG:HD3	1.55	0.89
3:J:105:VAL:HA	3:J:112:ILE:HD11	1.55	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	225/314 (72%)	191 (85%)	32 (14%)	2 (1%)	17	56
1	B	225/314 (72%)	196 (87%)	27 (12%)	2 (1%)	17	56
1	G	225/314 (72%)	190 (84%)	33 (15%)	2 (1%)	17	56
1	H	225/314 (72%)	196 (87%)	27 (12%)	2 (1%)	17	56
2	C	1115/1119 (100%)	974 (87%)	137 (12%)	4 (0%)	34	72
2	I	1115/1119 (100%)	974 (87%)	137 (12%)	4 (0%)	34	72
3	D	1486/1524 (98%)	1306 (88%)	171 (12%)	9 (1%)	25	65
3	J	1361/1524 (89%)	1200 (88%)	156 (12%)	5 (0%)	34	72
4	E	91/99 (92%)	82 (90%)	9 (10%)	0	100	100
4	K	91/99 (92%)	82 (90%)	9 (10%)	0	100	100
5	F	343/347 (99%)	301 (88%)	41 (12%)	1 (0%)	41	76
5	L	343/347 (99%)	300 (88%)	42 (12%)	1 (0%)	41	76
6	M	156/164 (95%)	143 (92%)	11 (7%)	2 (1%)	12	48
6	N	156/164 (95%)	142 (91%)	12 (8%)	2 (1%)	12	48
All	All	7157/7762 (92%)	6277 (88%)	844 (12%)	36 (0%)	29	68

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	53	VAL
3	D	681	ARG
3	D	683	ILE
3	D	1128	VAL
1	G	53	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/270 (72%)	171 (88%)	23 (12%)	5	23
1	B	194/270 (72%)	167 (86%)	27 (14%)	3	19
1	G	194/270 (72%)	171 (88%)	23 (12%)	5	23
1	H	194/270 (72%)	167 (86%)	27 (14%)	3	19
2	C	931/936 (100%)	820 (88%)	111 (12%)	5	23
2	I	931/936 (100%)	820 (88%)	111 (12%)	5	23
3	D	1252/1281 (98%)	1115 (89%)	137 (11%)	6	26
3	J	1150/1281 (90%)	1033 (90%)	117 (10%)	7	28
4	E	83/88 (94%)	77 (93%)	6 (7%)	14	41
4	K	83/88 (94%)	77 (93%)	6 (7%)	14	41
5	F	296/299 (99%)	267 (90%)	29 (10%)	8	29
5	L	296/299 (99%)	267 (90%)	29 (10%)	8	29
6	M	127/133 (96%)	122 (96%)	5 (4%)	32	57
6	N	127/133 (96%)	122 (96%)	5 (4%)	32	57
All	All	6052/6554 (92%)	5396 (89%)	656 (11%)	6	26

5 of 656 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	848	VAL
3	J	1078	ARG
2	I	934	PHE
2	I	834	GLN
3	J	387	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 109 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	80	GLN
2	I	1050	GLN
5	L	263	ASN
2	I	187	ASN
2	I	683	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
9	Q	3/4 (75%)	2 (66%)	0
9	T	3/4 (75%)	0	0
All	All	6/8 (75%)	2 (33%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
9	Q	2	C
9	Q	3	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/314 (72%)	-0.23	1 (0%) 92 87	144, 176, 208, 241	0
1	B	227/314 (72%)	-0.41	1 (0%) 92 87	143, 164, 196, 228	0
1	G	227/314 (72%)	-0.03	2 (0%) 84 77	148, 188, 218, 252	0
1	H	227/314 (72%)	-0.34	0 100 100	145, 175, 207, 241	0
2	C	1117/1119 (99%)	-0.19	4 (0%) 92 87	144, 172, 211, 253	0
2	I	1117/1119 (99%)	-0.20	14 (1%) 77 68	144, 182, 221, 270	0
3	D	1490/1524 (97%)	-0.23	3 (0%) 95 93	117, 162, 195, 251	0
3	J	1367/1524 (89%)	-0.20	6 (0%) 92 87	120, 171, 204, 250	0
4	E	93/99 (93%)	-0.23	0 100 100	144, 165, 194, 217	0
4	K	93/99 (93%)	-0.23	0 100 100	144, 179, 206, 228	0
5	F	345/347 (99%)	-0.17	2 (0%) 89 84	144, 179, 222, 245	0
5	L	345/347 (99%)	-0.17	2 (0%) 89 84	145, 186, 225, 258	0
6	M	158/164 (96%)	0.17	5 (3%) 47 37	159, 207, 235, 243	0
6	N	158/164 (96%)	0.42	5 (3%) 47 37	171, 215, 240, 267	0
7	O	48/48 (100%)	0.33	4 (8%) 11 10	157, 217, 256, 270	0
7	R	48/48 (100%)	-0.22	1 (2%) 63 54	163, 207, 251, 276	0
8	P	48/48 (100%)	0.38	4 (8%) 11 10	161, 219, 260, 270	0
8	S	48/48 (100%)	-0.09	1 (2%) 63 54	167, 212, 250, 261	0
9	Q	4/4 (100%)	0.94	1 (25%) 0 1	175, 177, 186, 189	0
9	T	4/4 (100%)	-0.17	0 100 100	165, 183, 184, 196	0
All	All	7391/7962 (92%)	-0.18	56 (0%) 86 79	117, 174, 219, 276	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	187	GLY	4.7
2	I	246	ASP	4.4
3	J	216	LEU	3.4
3	D	1253	THR	3.4
7	O	1	DC	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
10	ZN	J	2001	1/1	0.89	0.14	277,277,277,277	0
10	ZN	D	2002	1/1	0.92	0.16	237,237,237,237	0
10	ZN	J	2002	1/1	0.95	0.07	157,157,157,157	0
11	MG	J	2003	1/1	0.96	0.19	270,270,270,270	0
11	MG	D	2003	1/1	0.97	0.09	283,283,283,283	0
10	ZN	D	2001	1/1	0.98	0.10	116,116,116,116	0

6.5 Other polymers [i](#)

There are no such residues in this entry.