



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 28, 2024 – 12:30 pm BST

PDB ID : 2XM3  
Title : Deinococcus radiodurans ISDr2 Transposase Left end DNA complex  
Authors : Hickman, A.B.; James, J.A.; Barabas, O.; Pasternak, C.; Ton-Hoang, B.;  
Chandler, M.; Sommer, S.; Dyda, F.  
Deposited on : 2010-07-22  
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	 4% 68% 19% • 10%
1	B	140	 4% 74% 16% • 8%
1	C	140	 5% 69% 19% • 9%
1	D	140	 8% 72% 17% • 8%
1	E	140	 10% 67% 20% • 10%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	140	 6% 66% 20% 10%
2	G	27	 56% 41%
2	I	27	 44% 44% 11%
2	K	27	 37% 56% 7%
2	M	27	 33% 59% 7%
2	O	27	 15% 11% 81% 7%
2	Q	27	 7% 26% 70%
3	H	5	 80% 20%
3	J	5	 80% 20%
3	L	5	 20% 80%
3	N	5	 80% 20%
3	P	5	 20% 80%
3	R	5	 80% 20%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSPOSASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	126	Total 1017	C 656	N 176	O 179	S 6	0	0	0
1	B	129	Total 1040	C 673	N 179	O 182	S 6	0	0	0
1	C	128	Total 1032	C 667	N 178	O 181	S 6	0	0	0
1	D	129	Total 1042	C 670	N 182	O 184	S 6	0	0	0
1	E	126	Total 1017	C 656	N 176	O 179	S 6	0	0	0
1	F	126	Total 1017	C 656	N 176	O 179	S 6	0	0	0


- Molecule 2 is a DNA chain called DRA2 TRANSPOSASE BINDING ELEMENT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	G	27	Total 547	C 262	N 98	O 161	P 26	0	0	0
2	I	27	Total 547	C 262	N 98	O 161	P 26	0	0	0
2	K	27	Total 547	C 262	N 98	O 161	P 26	0	0	0
2	M	27	Total 547	C 262	N 98	O 161	P 26	0	0	0
2	O	27	Total 547	C 262	N 98	O 161	P 26	0	0	0
2	Q	27	Total 547	C 262	N 98	O 161	P 26	0	0	0

- Molecule 3 is a DNA chain called 5'-D(\*TP\*TP\*AP\*GP\*T)-3'.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
3	H	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			
3	J	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			
3	L	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			
3	N	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			
3	P	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			
3	R	5	Total	C	N	O	P	0	0	0
			100	50	16	30	4			

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	C O	0	0
			4	2 2		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	E	1	Total Mg	0	0
			1 1		
5	F	1	Total Mg	0	0
			1 1		
5	G	2	Total Mg	0	0
			2 2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	I	2	Total Mg 2 2	0	0
5	K	2	Total Mg 2 2	0	0
5	M	2	Total Mg 2 2	0	0
5	O	1	Total Mg 1 1	0	0
5	Q	1	Total Mg 1 1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	G	1	Total C O 6 3 3	0	0
6	J	1	Total C O 6 3 3	0	0
6	L	1	Total C O 6 3 3	0	0
6	P	1	Total C O 6 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	58	Total 58	O 58	0	0
7	B	53	Total 53	O 53	0	0
7	C	58	Total 58	O 58	0	0
7	D	43	Total 43	O 43	0	0
7	E	39	Total 39	O 39	0	0
7	F	30	Total 30	O 30	0	0
7	G	35	Total 35	O 35	0	0
7	H	3	Total 3	O 3	0	0
7	I	42	Total 42	O 42	0	0
7	J	8	Total 8	O 8	0	0
7	K	45	Total 45	O 45	0	0
7	L	4	Total 4	O 4	0	0
7	M	29	Total 29	O 29	0	0
7	N	8	Total 8	O 8	0	0
7	O	23	Total 23	O 23	0	0
7	Q	25	Total 25	O 25	0	0
7	R	6	Total 6	O 6	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TRANSPOSASE




- Molecule 1: TRANSPOSASE



- Molecule 1: TRANSPOSASE




- Molecule 1: TRANSPOSASE




- Molecule 1: TRANSPOSASE






- Molecule 1: TRANSPOSASE



- Molecule 2: DRA2 TRANSPOSASE BINDING ELEMENT



- Molecule 2: DRA2 TRANSPOSASE BINDING ELEMENT



- Molecule 2: DRA2 TRANSPOSASE BINDING ELEMENT



- Molecule 2: DRA2 TRANSPOSASE BINDING ELEMENT



- Molecule 2: DRA2 TRANSPOSASE BINDING ELEMENT





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.58Å 128.38Å 140.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 29.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.1 (30.00-2.30) 93.1 (29.97-2.30)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 2.31Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.218 , 0.249 0.204 , 0.237	Depositor DCC
$R_{free}$ test set	1575 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.9	Xtrriage
Anisotropy	0.286	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10596	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1040	0.76	0/1408
1	B	0.55	0/1064	0.75	0/1442
1	C	0.55	0/1056	0.77	0/1430
1	D	0.53	0/1066	0.79	1/1443 (0.1%)
1	E	0.55	0/1040	0.79	0/1408
1	F	0.51	0/1040	0.76	1/1408 (0.1%)
2	G	0.63	0/612	0.83	0/942
2	I	0.70	0/612	0.83	2/942 (0.2%)
2	K	0.70	0/612	0.83	0/942
2	M	0.62	0/612	0.84	1/942 (0.1%)
2	O	0.50	0/612	0.74	0/942
2	Q	0.60	0/612	0.81	0/942
3	H	0.60	0/111	0.76	0/170
3	J	0.75	0/111	0.63	0/170
3	L	0.68	0/111	0.85	0/170
3	N	0.57	0/111	0.73	0/170
3	P	0.59	0/111	0.75	0/170
3	R	0.67	0/111	0.80	0/170
All	All	0.58	0/10644	0.78	5/15211 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	G	0	1
2	I	0	2
2	K	0	5
2	M	0	1
2	O	0	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	#Chirality outliers	#Planarity outliers
2	Q	0	1
3	P	0	1
All	All	0	13

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	30	DT	O5'-P-OP2	-7.52	98.94	105.70
2	I	13	DC	C1'-O4'-C4'	-5.57	104.53	110.10
1	D	14	ARG	NE-CZ-NH2	-5.30	117.65	120.30
2	I	31	DC	O4'-C4'-C3'	-5.28	102.39	104.50
1	F	107	TRP	N-CA-C	-5.01	97.47	111.00

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	G	15	DC	Sidechain
2	I	15	DC	Sidechain
2	I	28	DA	Sidechain
2	K	15	DC	Sidechain
2	K	23	DA	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1017	0	1033	23	0
1	B	1040	0	1062	20	0
1	C	1032	0	1052	27	0
1	D	1042	0	1053	25	0
1	E	1017	0	1033	25	0
1	F	1017	0	1033	33	0
2	G	547	0	306	13	0
2	I	547	0	306	14	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	K	547	0	306	13	0
2	M	547	0	306	26	0
2	O	547	0	306	55	0
2	Q	547	0	306	37	0
3	H	100	0	60	1	0
3	J	100	0	60	1	0
3	L	100	0	60	5	0
3	N	100	0	60	1	0
3	P	100	0	60	6	0
3	R	100	0	60	1	0
4	B	4	0	3	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	2	0	0	0	0
5	I	2	0	0	0	0
5	K	2	0	0	0	0
5	M	2	0	0	0	0
5	O	1	0	0	0	0
5	Q	1	0	0	0	0
6	G	6	0	8	1	0
6	J	6	0	8	2	0
6	L	6	0	8	2	0
6	P	6	0	8	2	0
7	A	58	0	0	2	0
7	B	53	0	0	1	0
7	C	58	0	0	1	0
7	D	43	0	0	0	0
7	E	39	0	0	0	0
7	F	30	0	0	0	0
7	G	35	0	0	0	0
7	H	3	0	0	0	0
7	I	42	0	0	1	0
7	J	8	0	0	0	0
7	K	45	0	0	0	0
7	L	4	0	0	0	0
7	M	29	0	0	0	0
7	N	8	0	0	0	0
7	O	23	0	0	0	0
7	Q	25	0	0	0	0
7	R	6	0	0	0	0
All	All	10596	0	8497	294	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 294 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:O:18:DT:H5'	2:O:18:DT:C6	1.79	1.18
2:O:18:DT:H5'	2:O:18:DT:H6	0.98	1.14
2:M:36:DG:C2'	2:M:37:DT:H5''	1.76	1.13
2:O:17:DC:C2'	2:O:18:DT:H5''	1.78	1.13
2:M:12:DG:H2''	2:M:13:DC:H5''	1.34	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/140 (89%)	119 (96%)	4 (3%)	1 (1%)	19	23
1	B	127/140 (91%)	125 (98%)	2 (2%)	0	100	100
1	C	126/140 (90%)	125 (99%)	1 (1%)	0	100	100
1	D	127/140 (91%)	123 (97%)	3 (2%)	1 (1%)	19	23
1	E	124/140 (89%)	123 (99%)	1 (1%)	0	100	100
1	F	124/140 (89%)	121 (98%)	3 (2%)	0	100	100
All	All	752/840 (90%)	736 (98%)	14 (2%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	GLU
1	D	124	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/122 (88%)	102 (94%)	6 (6%)	21	29
1	B	111/122 (91%)	104 (94%)	7 (6%)	18	24
1	C	110/122 (90%)	105 (96%)	5 (4%)	27	39
1	D	111/122 (91%)	105 (95%)	6 (5%)	22	30
1	E	108/122 (88%)	100 (93%)	8 (7%)	13	17
1	F	108/122 (88%)	100 (93%)	8 (7%)	13	17
All	All	656/732 (90%)	616 (94%)	40 (6%)	18	25

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	111	LEU
1	F	88	LYS
1	E	118	ILE
1	F	68	VAL
1	F	118	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	113	ASN
1	D	69	HIS
1	F	113	ASN
1	F	23	HIS
1	F	69	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 12 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	G	1038	-	5,5,5	0.68	0	5,5,5	0.35	0
6	GOL	P	1007	-	5,5,5	0.72	0	5,5,5	0.29	0
4	ACT	B	1135	-	3,3,3	0.71	0	3,3,3	0.79	0
6	GOL	L	1006	-	5,5,5	0.46	0	5,5,5	0.25	0
6	GOL	J	1006	-	5,5,5	0.65	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	P	1007	-	-	0/4/4/4	-
6	GOL	J	1006	-	-	0/4/4/4	-
6	GOL	L	1006	-	-	0/4/4/4	-
6	GOL	G	1038	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	1038	GOL	1	0
6	P	1007	GOL	2	0
6	L	1006	GOL	2	0
6	J	1006	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	126/140 (90%)	0.18	5 (3%) 38 45	17, 30, 65, 98	0
1	B	129/140 (92%)	0.32	6 (4%) 31 38	19, 32, 63, 90	0
1	C	128/140 (91%)	0.20	7 (5%) 25 31	19, 30, 66, 97	0
1	D	129/140 (92%)	0.44	11 (8%) 10 14	19, 33, 85, 106	0
1	E	126/140 (90%)	0.51	14 (11%) 5 7	22, 39, 81, 107	0
1	F	126/140 (90%)	0.56	9 (7%) 16 21	23, 44, 71, 79	0
2	G	27/27 (100%)	-0.25	0 100 100	24, 34, 49, 59	0
2	I	27/27 (100%)	-0.24	0 100 100	22, 35, 43, 52	0
2	K	27/27 (100%)	-0.40	0 100 100	18, 28, 38, 49	0
2	M	27/27 (100%)	-0.14	0 100 100	26, 36, 52, 62	0
2	O	27/27 (100%)	0.68	4 (14%) 2 3	31, 53, 68, 77	0
2	Q	27/27 (100%)	0.58	2 (7%) 14 19	28, 46, 75, 79	0
3	H	5/5 (100%)	-0.60	0 100 100	25, 30, 35, 37	0
3	J	5/5 (100%)	-0.75	0 100 100	26, 28, 29, 29	0
3	L	5/5 (100%)	-0.37	0 100 100	29, 29, 31, 35	0
3	N	5/5 (100%)	-0.83	0 100 100	30, 31, 33, 36	0
3	P	5/5 (100%)	-0.09	0 100 100	33, 35, 43, 46	0
3	R	5/5 (100%)	-0.91	0 100 100	30, 32, 34, 43	0
All	All	956/1032 (92%)	0.28	58 (6%) 21 27	17, 35, 72, 107	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	131	LYS	7.0
1	D	137	HIS	5.8
1	B	130	GLN	5.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	6	LEU	5.3
1	E	133	ILE	4.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q < 0.9
4	ACT	B	1135	4/4	0.79	0.27	62,63,63,63	0
6	GOL	L	1006	6/6	0.79	0.38	44,48,48,50	0
6	GOL	J	1006	6/6	0.81	0.28	44,45,46,48	0
5	MG	G	1114	1/1	0.84	0.07	36,36,36,36	0
6	GOL	G	1038	6/6	0.86	0.37	46,46,47,48	0
6	GOL	P	1007	6/6	0.91	0.25	40,41,43,43	0
5	MG	O	1112	1/1	0.92	0.06	52,52,52,52	0
5	MG	Q	1111	1/1	0.92	0.11	40,40,40,40	0
5	MG	M	1121	1/1	0.92	0.05	43,43,43,43	0
5	MG	I	1038	1/1	0.94	0.20	26,26,26,26	0
5	MG	E	1135	1/1	0.94	0.21	35,35,35,35	0
5	MG	K	1038	1/1	0.95	0.16	21,21,21,21	0
5	MG	K	1039	1/1	0.96	0.04	34,34,34,34	0
5	MG	G	1039	1/1	0.98	0.22	23,23,23,23	0
5	MG	F	1135	1/1	0.98	0.09	39,39,39,39	0
5	MG	M	1038	1/1	0.99	0.19	24,24,24,24	0
5	MG	I	1116	1/1	0.99	0.04	34,34,34,34	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.