



## wwPDB EM Validation Summary Report ⓘ

Nov 7, 2022 – 04:44 PM JST

PDB ID : 5XXX  
EMDB ID : EMD-6783  
Title : GMPCPP-microtubule complexed with nucleotide-free KIF5C  
Authors : Morikawa, M.; Shigematsu, H.; Nitta, R.; Hirokawa, N.  
Deposited on : 2017-07-05  
Resolution : 6.43 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 6.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	<div style="display: flex; justify-content: space-between;"> <span>34%</span> <span>74%</span> <span>22%</span> </div>
1	C	438	<div style="display: flex; justify-content: space-between;"> <span>33%</span> <span>73%</span> <span>22%</span> </div>
1	E	438	<div style="display: flex; justify-content: space-between;"> <span>34%</span> <span>74%</span> <span>22%</span> </div>
1	G	438	<div style="display: flex; justify-content: space-between;"> <span>36%</span> <span>74%</span> <span>22%</span> </div>
1	I	438	<div style="display: flex; justify-content: space-between;"> <span>37%</span> <span>73%</span> <span>22%</span> </div>
1	K	438	<div style="display: flex; justify-content: space-between;"> <span>37%</span> <span>74%</span> <span>21%</span> </div>
1	M	438	<div style="display: flex; justify-content: space-between;"> <span>47%</span> <span>75%</span> <span>21%</span> </div>
1	O	438	<div style="display: flex; justify-content: space-between;"> <span>47%</span> <span>75%</span> <span>20%</span> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	Q	438	47%	76% 20% ..
2	B	426	34%	76% 20% .
2	D	426	34%	77% 19% .
2	F	426	34%	77% 20% .
2	H	426	44%	79% 18% .
2	J	426	44%	78% 19% .
2	L	426	43%	79% 18% .
2	N	426	38%	78% 19% .
2	P	426	37%	76% 20% .
2	R	426	38%	77% 20% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	G2P	B	501	-	-	X	-
5	G2P	D	501	-	-	X	-
5	G2P	F	501	-	-	X	-
5	G2P	H	501	-	-	X	-
5	G2P	J	501	-	-	X	-
5	G2P	L	501	-	-	X	-
5	G2P	N	501	-	-	X	-
5	G2P	P	501	-	-	X	-
5	G2P	R	501	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 60948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1A chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	428	3350	2121	570	638	21	0	0
1	C	428	3350	2121	570	638	21	0	0
1	E	428	3350	2121	570	638	21	0	0
1	G	428	3350	2121	570	638	21	0	0
1	I	428	3350	2121	570	638	21	0	0
1	K	428	3350	2121	570	638	21	0	0
1	M	428	3350	2121	570	638	21	0	0
1	O	428	3350	2121	570	638	21	0	0
1	Q	428	3350	2121	570	638	21	0	0

- Molecule 2 is a protein called Tubulin beta chain.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	426	3352	2105	575	647	25	0	0
2	D	426	3352	2105	575	647	25	0	0
2	F	426	3352	2105	575	647	25	0	0
2	H	426	3352	2105	575	647	25	0	0
2	J	426	3352	2105	575	647	25	0	0
2	L	426	3352	2105	575	647	25	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	N	426	Total	C	N	O	S	0	0
			3352	2105	575	647	25		
2	P	426	Total	C	N	O	S	0	0
			3352	2105	575	647	25		
2	R	426	Total	C	N	O	S	0	0
			3352	2105	575	647	25		

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
3	A	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	C	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	E	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	G	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	I	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	K	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	M	1	Total	C	N	O	P	0
			32	10	5	14	3	
3	O	1	Total	C	N	O	P	0
			32	10	5	14	3	

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
3	Q	1	32	10	5	14	3	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
4	A	1	Total	Mg	0
			1	1	
4	B	1	Total	Mg	0
			1	1	
4	C	1	Total	Mg	0
			1	1	
4	D	1	Total	Mg	0
			1	1	
4	E	1	Total	Mg	0
			1	1	
4	F	1	Total	Mg	0
			1	1	
4	G	1	Total	Mg	0
			1	1	
4	H	1	Total	Mg	0
			1	1	
4	I	1	Total	Mg	0
			1	1	
4	J	1	Total	Mg	0
			1	1	
4	K	1	Total	Mg	0
			1	1	
4	L	1	Total	Mg	0
			1	1	
4	M	1	Total	Mg	0
			1	1	
4	N	1	Total	Mg	0
			1	1	
4	O	1	Total	Mg	0
			1	1	
4	P	1	Total	Mg	0
			1	1	
4	Q	1	Total	Mg	0
			1	1	
4	R	1	Total	Mg	0
			1	1	

- Molecule 5 is PHOSPHOMETHYLPHOSPHONIC ACID GUANYLATE ESTER (three-letter code: G2P) (formula:  $C_{11}H_{18}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
5	B	1	Total 32	C 11	N 5	O 13	P 3	0
5	D	1	Total 32	C 11	N 5	O 13	P 3	0
5	F	1	Total 32	C 11	N 5	O 13	P 3	0
5	H	1	Total 32	C 11	N 5	O 13	P 3	0
5	J	1	Total 32	C 11	N 5	O 13	P 3	0
5	L	1	Total 32	C 11	N 5	O 13	P 3	0
5	N	1	Total 32	C 11	N 5	O 13	P 3	0
5	P	1	Total 32	C 11	N 5	O 13	P 3	0
5	R	1	Total 32	C 11	N 5	O 13	P 3	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		AltConf
			Total	O	
6	A	4	Total 4	O 4	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		AltConf
6	C	4	Total 4	O 4	0
6	E	4	Total 4	O 4	0
6	G	4	Total 4	O 4	0
6	I	4	Total 4	O 4	0
6	K	4	Total 4	O 4	0
6	M	4	Total 4	O 4	0
6	O	4	Total 4	O 4	0
6	Q	4	Total 4	O 4	0




### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tubulin alpha-1A chain





- Molecule 1: Tubulin alpha-1A chain








• Molecule 1: Tubulin alpha-1A chain





• Molecule 1: Tubulin alpha-1A chain







• Molecule 1: Tubulin alpha-1A chain





• Molecule 1: Tubulin alpha-1A chain







• Molecule 1: Tubulin alpha-1A chain





• Molecule 1: Tubulin alpha-1A chain





• Molecule 1: Tubulin alpha-1A chain




• Molecule 2: Tubulin beta chain




• Molecule 2: Tubulin beta chain





• Molecule 2: Tubulin beta chain








• Molecule 2: Tubulin beta chain





• Molecule 2: Tubulin beta chain








• Molecule 2: Tubulin beta chain





• Molecule 2: Tubulin beta chain










• Molecule 2: Tubulin beta chain



• Molecule 2: Tubulin beta chain





## 4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-25.7541°, rise=8.8632 Å, axial sym=C1	Depositor
Number of segments used	5822	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{Å}^2$ )	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	13.831	Depositor
Minimum map value	-4.366	Depositor
Average map value	1.018	Depositor
Map value standard deviation	2.451	Depositor
Recommended contour level	7.4	Depositor
Map size (Å)	246.52802, 246.52802, 282.48	wwPDB
Map dimensions	192, 192, 220	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.284, 1.284, 1.284	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: G2P, MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
1	C	1.14	10/3427 (0.3%)	1.44	45/4651 (1.0%)
1	E	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
1	G	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
1	I	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
1	K	1.14	10/3427 (0.3%)	1.44	45/4651 (1.0%)
1	M	1.14	10/3427 (0.3%)	1.44	45/4651 (1.0%)
1	O	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
1	Q	1.14	10/3427 (0.3%)	1.44	44/4651 (0.9%)
2	B	1.05	5/3427 (0.1%)	1.56	49/4642 (1.1%)
2	D	1.05	5/3427 (0.1%)	1.56	47/4642 (1.0%)
2	F	1.05	5/3427 (0.1%)	1.56	46/4642 (1.0%)
2	H	1.05	5/3427 (0.1%)	1.56	46/4642 (1.0%)
2	J	1.05	5/3427 (0.1%)	1.56	47/4642 (1.0%)
2	L	1.05	5/3427 (0.1%)	1.56	49/4642 (1.1%)
2	N	1.05	5/3427 (0.1%)	1.56	47/4642 (1.0%)
2	P	1.05	5/3427 (0.1%)	1.56	48/4642 (1.0%)
2	R	1.05	5/3427 (0.1%)	1.56	47/4642 (1.0%)
All	All	1.09	135/61686 (0.2%)	1.50	825/83637 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
2	F	0	1
2	H	0	1
2	J	0	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	1
2	N	0	1
2	P	0	1
2	R	0	1
All	All	0	9

The worst 5 of 135 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	M	265	GLY	CA-C	-15.12	1.27	1.51
1	G	265	GLY	CA-C	-15.03	1.27	1.51
1	A	265	GLY	CA-C	-15.03	1.27	1.51
1	C	265	GLY	CA-C	-15.02	1.27	1.51
1	Q	265	GLY	CA-C	-14.99	1.27	1.51

The worst 5 of 825 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	48	ARG	NE-CZ-NH1	20.58	130.59	120.30
2	P	48	ARG	NE-CZ-NH1	20.45	130.53	120.30
2	B	48	ARG	NE-CZ-NH1	20.42	130.51	120.30
2	J	48	ARG	NE-CZ-NH1	20.36	130.48	120.30
2	L	48	ARG	NE-CZ-NH1	20.31	130.46	120.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	248	LEU	Peptide
2	D	248	LEU	Peptide
2	F	248	LEU	Peptide
2	H	248	LEU	Peptide
2	J	248	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3350	0	3254	332	0
1	C	3350	0	3254	314	0
1	E	3350	0	3254	331	0
1	G	3350	0	3254	322	0
1	I	3350	0	3254	343	0
1	K	3350	0	3254	317	0
1	M	3350	0	3254	242	0
1	O	3350	0	3254	273	0
1	Q	3350	0	3254	268	0
2	B	3352	0	3227	348	0
2	D	3352	0	3227	324	0
2	F	3352	0	3227	336	0
2	H	3352	0	3227	279	0
2	J	3352	0	3227	302	0
2	L	3352	0	3227	275	0
2	N	3352	0	3227	313	0
2	P	3352	0	3227	357	0
2	R	3352	0	3227	346	0
3	A	32	0	12	3	0
3	C	32	0	12	2	0
3	E	32	0	12	2	0
3	G	32	0	12	2	0
3	I	32	0	12	3	0
3	K	32	0	12	2	0
3	M	32	0	12	3	0
3	O	32	0	12	3	0
3	Q	32	0	12	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
4	L	1	0	0	0	0
4	M	1	0	0	0	0
4	N	1	0	0	0	0
4	O	1	0	0	0	0
4	P	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Q	1	0	0	0	0
4	R	1	0	0	0	0
5	B	32	0	13	12	0
5	D	32	0	13	11	0
5	F	32	0	13	12	0
5	H	32	0	13	11	0
5	J	32	0	13	12	0
5	L	32	0	13	11	0
5	N	32	0	13	11	0
5	P	32	0	13	12	0
5	R	32	0	13	12	0
6	A	4	0	0	0	0
6	C	4	0	0	0	0
6	E	4	0	0	0	0
6	G	4	0	0	0	0
6	I	4	0	0	0	0
6	K	4	0	0	0	0
6	M	4	0	0	0	0
6	O	4	0	0	0	0
6	Q	4	0	0	0	0
All	All	60948	0	58554	3168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 3168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:401:LYS:HD3	2:H:346:TRP:CD2	1.32	1.64
1:K:101:ASN:HB2	2:L:254:LYS:CE	1.24	1.63
1:I:101:ASN:HB2	2:J:254:LYS:CE	1.20	1.62
1:G:181:VAL:CA	2:H:352:LYS:HE3	1.30	1.61
1:Q:181:VAL:CA	2:R:352:LYS:HE3	1.27	1.60

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	C	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	E	424/438 (97%)	408 (96%)	14 (3%)	2 (0%)	29	69
1	G	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	I	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	K	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	M	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	O	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
1	Q	424/438 (97%)	409 (96%)	13 (3%)	2 (0%)	29	69
2	B	424/426 (100%)	409 (96%)	12 (3%)	3 (1%)	22	63
2	D	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	F	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	H	424/426 (100%)	409 (96%)	12 (3%)	3 (1%)	22	63
2	J	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	L	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	N	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	P	424/426 (100%)	408 (96%)	13 (3%)	3 (1%)	22	63
2	R	424/426 (100%)	409 (96%)	12 (3%)	3 (1%)	22	63
All	All	7632/7776 (98%)	7355 (96%)	232 (3%)	45 (1%)	29	66

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	98	GLY
2	D	98	GLY
2	F	98	GLY
2	H	98	GLY

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
2	J	98	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	C	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	E	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	G	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	I	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	K	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	M	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	O	360/367 (98%)	359 (100%)	1 (0%)	92	95
1	Q	360/367 (98%)	359 (100%)	1 (0%)	92	95
2	B	367/367 (100%)	367 (100%)	0	100	100
2	D	367/367 (100%)	367 (100%)	0	100	100
2	F	367/367 (100%)	367 (100%)	0	100	100
2	H	367/367 (100%)	367 (100%)	0	100	100
2	J	367/367 (100%)	367 (100%)	0	100	100
2	L	367/367 (100%)	367 (100%)	0	100	100
2	N	367/367 (100%)	367 (100%)	0	100	100
2	P	367/367 (100%)	367 (100%)	0	100	100
2	R	367/367 (100%)	367 (100%)	0	100	100
All	All	6543/6606 (99%)	6534 (100%)	9 (0%)	93	97

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	255	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	Q	255	PHE
1	G	255	PHE
1	I	255	PHE
1	K	255	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 97 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	301	GLN
2	N	96	GLN
2	L	8	GLN
1	M	61	HIS
2	N	406	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 18 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GTP	I	501	4	26,34,34	1.29	4 (15%)	32,54,54	1.10	3 (9%)
5	G2P	D	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.25	18 (54%)
5	G2P	P	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.24	18 (54%)
5	G2P	J	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.25	18 (54%)
3	GTP	K	501	4	26,34,34	1.29	4 (15%)	32,54,54	1.10	3 (9%)
5	G2P	R	501	4	27,34,34	4.63	23 (85%)	33,54,54	4.23	18 (54%)
5	G2P	L	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.26	18 (54%)
3	GTP	E	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
3	GTP	Q	501	4	26,34,34	1.31	4 (15%)	32,54,54	1.10	3 (9%)
3	GTP	A	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
3	GTP	M	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
3	GTP	C	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
5	G2P	N	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.24	18 (54%)
3	GTP	O	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
5	G2P	F	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.25	18 (54%)
3	GTP	G	501	4	26,34,34	1.30	4 (15%)	32,54,54	1.10	3 (9%)
5	G2P	H	501	4	27,34,34	4.65	23 (85%)	33,54,54	4.24	18 (54%)
5	G2P	B	501	4	27,34,34	4.64	23 (85%)	33,54,54	4.25	18 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	I	501	4	-	3/18/38/38	0/3/3/3
5	G2P	D	501	4	-	4/15/38/38	0/3/3/3
5	G2P	P	501	4	-	4/15/38/38	0/3/3/3
5	G2P	J	501	4	-	4/15/38/38	0/3/3/3
3	GTP	K	501	4	-	3/18/38/38	0/3/3/3
5	G2P	R	501	4	-	4/15/38/38	0/3/3/3
5	G2P	L	501	4	-	4/15/38/38	0/3/3/3
3	GTP	E	501	4	-	3/18/38/38	0/3/3/3
3	GTP	Q	501	4	-	3/18/38/38	0/3/3/3
3	GTP	A	501	4	-	3/18/38/38	0/3/3/3
3	GTP	M	501	4	-	3/18/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	C	501	4	-	3/18/38/38	0/3/3/3
5	G2P	N	501	4	-	4/15/38/38	0/3/3/3
3	GTP	O	501	4	-	3/18/38/38	0/3/3/3
5	G2P	F	501	4	-	4/15/38/38	0/3/3/3
3	GTP	G	501	4	-	3/18/38/38	0/3/3/3
5	G2P	H	501	4	-	4/15/38/38	0/3/3/3
5	G2P	B	501	4	-	4/15/38/38	0/3/3/3

The worst 5 of 243 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	R	501	G2P	C2-N2	8.53	1.51	1.33
5	B	501	G2P	C2-N2	8.51	1.50	1.33
5	F	501	G2P	C2-N2	8.50	1.50	1.33
5	H	501	G2P	C2-N2	8.49	1.50	1.33
5	P	501	G2P	C2-N2	8.49	1.50	1.33

The worst 5 of 189 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	501	G2P	N3-C2-N1	-11.10	112.41	127.22
5	J	501	G2P	N3-C2-N1	-11.09	112.43	127.22
5	D	501	G2P	N3-C2-N1	-11.07	112.46	127.22
5	L	501	G2P	N3-C2-N1	-11.07	112.46	127.22
5	B	501	G2P	N3-C2-N1	-11.05	112.48	127.22

There are no chirality outliers.

5 of 63 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	501	GTP	C3'-C4'-C5'-O5'
3	C	501	GTP	C3'-C4'-C5'-O5'
3	E	501	GTP	C3'-C4'-C5'-O5'
3	I	501	GTP	C3'-C4'-C5'-O5'
3	K	501	GTP	C3'-C4'-C5'-O5'



There are no ring outliers.


18 monomers are involved in 127 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	501	GTP	3	0
5	D	501	G2P	11	0
5	P	501	G2P	12	0
5	J	501	G2P	12	0
3	K	501	GTP	2	0
5	R	501	G2P	12	0
5	L	501	G2P	11	0
3	E	501	GTP	2	0
3	Q	501	GTP	3	0
3	A	501	GTP	3	0
3	M	501	GTP	3	0
3	C	501	GTP	2	0
5	N	501	G2P	11	0
3	O	501	GTP	3	0
5	F	501	G2P	12	0
3	G	501	GTP	2	0
5	H	501	G2P	11	0
5	B	501	G2P	12	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-6783. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 96



Y Index: 96




Z Index: 110


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 66



Y Index: 74



Z Index: 105


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 7.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.




## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 583 nm<sup>3</sup>; this corresponds to an approximate mass of 527 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

## 7.3 Rotationally averaged power spectrum [i](#)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-6783 and PDB model 5XXX. Per-residue inclusion information can be found in section [3](#) on page [9](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 7.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (7.4).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 47% of all backbone atoms, 46% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (7.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4643	0.1540
A	0.5005	0.1600
B	0.5062	0.1510
C	0.5035	0.1600
D	0.5086	0.1540
E	0.4974	0.1590
F	0.5044	0.1520
G	0.4860	0.1520
H	0.4472	0.1510
I	0.4800	0.1490
J	0.4466	0.1490
K	0.4800	0.1520
L	0.4475	0.1490
M	0.4161	0.1580
N	0.4809	0.1540
O	0.4143	0.1580
P	0.4794	0.1530
Q	0.4119	0.1560
R	0.4767	0.1500

