



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2022 – 12:07 AM JST

PDB ID : 7Y3I  
Title : Structure of DNA bound SALL4  
Authors : Ru, W.; Xu, C.  
Deposited on : 2022-06-10  
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	12	
1	L	12	
2	H	12	
2	K	12	
3	A	78	
3	B	78	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	78	 33% 65%
3	D	78	 28% 5% 67%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2357 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (12-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	L	12	247	118	47	70	12	0	0	0
1	G	12	247	118	47	70	12	0	0	0

- Molecule 2 is a DNA chain called DNA (12-mer).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	K	12	245	118	41	74	12	0	0	0
2	H	12	245	118	41	74	12	0	0	0

- Molecule 3 is a protein called Sal-like protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	54	413	253	82	73	5	0	0	0
3	B	53	398	245	77	71	5	0	0	0
3	C	27	207	132	39	33	3	0	0	0
3	D	26	203	130	38	32	3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	853	GLY	-	expression tag	UNP Q9UJQ4
A	854	HIS	-	expression tag	UNP Q9UJQ4
B	853	GLY	-	expression tag	UNP Q9UJQ4
B	854	HIS	-	expression tag	UNP Q9UJQ4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	853	GLY	-	expression tag	UNP Q9UJQ4
C	854	HIS	-	expression tag	UNP Q9UJQ4
D	853	GLY	-	expression tag	UNP Q9UJQ4
D	854	HIS	-	expression tag	UNP Q9UJQ4

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total 2	Zn 2	0	0
4	B	2	Total 2	Zn 2	0	0
4	C	1	Total 1	Zn 1	0	0
4	D	1	Total 1	Zn 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	20	Total 20	O 20	0	0
5	K	21	Total 21	O 21	0	0
5	G	20	Total 20	O 20	0	0
5	H	15	Total 15	O 15	0	0
5	A	18	Total 18	O 18	0	0
5	B	30	Total 30	O 30	0	0
5	C	13	Total 13	O 13	0	0
5	D	9	Total 9	O 9	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (12-mer)

Chain L:  67% 33%



- Molecule 1: DNA (12-mer)

Chain G:  75% 25%



- Molecule 2: DNA (12-mer)

Chain K:  75% 17% 8%




- Molecule 2: DNA (12-mer)

Chain H:  92% 8%



- Molecule 3: Sal-like protein 4

Chain A:  % 65% 31%



- Molecule 3: Sal-like protein 4

Chain B:  62% 6% 32%

GLY HIS MET THR PRO LEU LEU ALA ALA PRO ARG ARG GLN ALA LYS Q869 N878 E895 K896 P897 K914 M918 G921 ALA ASN ASN ASN SER ALA ARG ARG GLY

● Molecule 3: Sal-like protein 4

Chain C:  33% 0% 65%

GLY HIS MET THR PRO LEU LEU ALA ALA GLN PRO ARG ARG GLN ALA LYS GLN HIS GLY CYS THR ARG CYS GLY LYS ASN SER PHE SER SER ALA ALA ALA LEU GLN ILE HIS GLU ARG THR HIS THR GLY E895 G921 ALA ASN ASN ASN SER ALA ARG GLY

● Molecule 3: Sal-like protein 4

Chain D:  28% 5% 67%

GLY HIS MET THR PRO LEU LEU ALA ALA GLN PRO ARG ARG GLN ALA LYS GLN HIS GLY CYS THR ARG CYS GLY LYS ASN PHE SER SER SER ALA ALA LEU GLN ILE HIS GLU ARG THR HIS THR GLY K896 T908 H916 M918 T919 H920 G921 ALA ASN ASN ASN SER ALA ARG

ARG  
GLY

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.98Å 69.20Å 110.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.13 – 2.45 43.13 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.13-2.45) 99.7 (43.13-2.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.194 , 0.242 0.194 , 0.239	Depositor DCC
$R_{free}$ test set	678 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.8	Xtrriage
Anisotropy	0.218	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 27.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2357	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5822e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	G	1.06	0/277	1.01	0/425
1	L	0.96	0/277	1.07	0/425
2	H	1.00	0/273	1.09	1/419 (0.2%)
2	K	1.01	1/273 (0.4%)	1.04	1/419 (0.2%)
3	A	0.37	0/422	0.54	0/565
3	B	0.43	0/407	0.57	0/546
3	C	0.42	0/212	0.55	0/283
3	D	0.43	0/208	0.49	0/278
All	All	0.75	1/2349 (0.0%)	0.84	2/3360 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	2	DT	C3'-O3'	-5.70	1.36	1.44

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	K	3	DA	P-O3'-C3'	6.17	127.10	119.70
2	H	4	DA	P-O3'-C3'	5.13	125.86	119.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	247	0	136	4	0
1	L	247	0	136	3	0
2	H	245	0	138	0	0
2	K	245	0	138	1	0
3	A	413	0	393	2	0
3	B	398	0	374	2	0
3	C	207	0	204	1	0
3	D	203	0	203	2	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	18	0	0	1	0
5	B	30	0	0	0	0
5	C	13	0	0	1	0
5	D	9	0	0	0	0
5	G	20	0	0	1	0
5	H	15	0	0	0	0
5	K	21	0	0	0	0
5	L	20	0	0	1	0
All	All	2357	0	1722	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:10:DA:H2''	1:L:11:DC:H5''	1.68	0.74
3:A:910:LYS:NZ	5:A:1101:HOH:O	2.34	0.59
1:G:2:DG:OP1	3:B:914:LYS:NZ	2.39	0.56
3:B:895:GLU:HG2	3:B:897:PRO:HD3	1.92	0.52
3:C:895:GLU:N	5:C:1102:HOH:O	2.42	0.52

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	52/78 (67%)	52 (100%)	0	0	100	100
3	B	51/78 (65%)	51 (100%)	0	0	100	100
3	C	25/78 (32%)	25 (100%)	0	0	100	100
3	D	24/78 (31%)	24 (100%)	0	0	100	100
All	All	152/312 (49%)	152 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	44/62 (71%)	44 (100%)	0	100	100
3	B	42/62 (68%)	40 (95%)	2 (5%)	25	33
3	C	22/62 (36%)	22 (100%)	0	100	100
3	D	22/62 (36%)	22 (100%)	0	100	100
All	All	130/248 (52%)	128 (98%)	2 (2%)	65	76

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	878	ASN
3	B	918	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	G	12/12 (100%)	-0.24	0 <a href="#">100</a> <a href="#">100</a>	27, 29, 44, 50	0
1	L	12/12 (100%)	-0.24	0 <a href="#">100</a> <a href="#">100</a>	28, 33, 40, 51	0
2	H	12/12 (100%)	-0.22	0 <a href="#">100</a> <a href="#">100</a>	27, 32, 44, 50	0
2	K	12/12 (100%)	0.07	0 <a href="#">100</a> <a href="#">100</a>	26, 37, 50, 61	0
3	A	54/78 (69%)	0.22	1 (1%) <a href="#">66</a> <a href="#">64</a>	25, 35, 53, 71	0
3	B	53/78 (67%)	0.25	0 <a href="#">100</a> <a href="#">100</a>	21, 31, 56, 72	0
3	C	27/78 (34%)	-0.10	0 <a href="#">100</a> <a href="#">100</a>	28, 32, 43, 56	0
3	D	26/78 (33%)	0.39	1 (3%) <a href="#">40</a> <a href="#">37</a>	30, 35, 63, 76	0
All	All	208/360 (57%)	0.12	2 (0%) <a href="#">82</a> <a href="#">83</a>	21, 33, 56, 76	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	868	LYS	4.8
3	D	917	TYR	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ZN	B	1001	1/1	0.98	0.08	36,36,36,36	0
4	ZN	A	1002	1/1	0.99	0.11	32,32,32,32	0
4	ZN	A	1001	1/1	0.99	0.09	30,30,30,30	0
4	ZN	C	1001	1/1	0.99	0.08	29,29,29,29	0
4	ZN	D	1001	1/1	0.99	0.10	41,41,41,41	0
4	ZN	B	1002	1/1	1.00	0.12	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ZN A 1002:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)












**Electron density around ZN C 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.