

Full wwPDB X-ray Structure Validation Report (i)

Nov 21, 2023 – 10:57 PM JST

PDB ID	:	7Y7Q
Title	:	QDE-1 in complex with RNA template, RNA primer and 3'-dGTP
Authors	:	Cui, R.X.; Gan, J.H.; Ma, J.B.
Deposited on	:	2022-06-22
Resolution	:	2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

Overall quality at a glance (i) 1

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution	on
---	----

Motria	Whole archive	Similar resolution
Metric	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R_{free}	130704	1692(2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672(2.04-2.04)
RNA backbone	3102	$1018 \ (2.50-1.58)$

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length		Quality of chain					
1	А	1026	4%	83%		10% 7%			
1	В	1026	3%	80%		8% 12%			
2	С	14		64%	7%	29%			
2	Е	14	7%	29%	57%				



Mol	Chain	Length	Quality of chain
	_		14%
3	D	7	100%



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 16072 atoms, of which 18 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	951	Total 7600	C 4864	N 1306	O 1394	S 36	0	13	0
1	В	906	Total 7213	C 4618	N 1254	0 1307	S 34	0	1	0

• Molecule 2 is a RNA chain called RNA (5'-R(*GP*AP*AP*CP*UP*AP*CP*CP*GP*UP* CP*GP*A)-3').

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
9	C	10	Total	С	Ν	Ο	Р	0	0	0
	U	10	213	95	38	70	10	0	0	
2	F	6	Total	С	Ν	Ο	Р	0	2	0
	0	125	56	21	42	6	0		0	

• Molecule 3 is a RNA chain called RNA (5'-R(*UP*CP*CP*GP*AP*CP*G)-3').

Mol	Chain	Residues		At	\mathbf{oms}			ZeroOcc	AltConf	Trace
3	D	7	Total 144	C 66	N 26	O 46	Р 6	0	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	2	Total Ca 2 2	0	0
4	В	2	Total Ca 2 2	0	0

• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	Total Mg 1 1	0	0
5	В	2	Total Mg 2 2	0	0

• Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total C H O 14 3 8 3	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 7 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
7	Λ	1	Total	С	Ν	Ο	Р	0	0
1	Л	1	31	10	5	13	3	0	0
7	В	1	Total	С	Ν	Ο	Р	0	0
1	D	1	31	10	5	13	3	0	0
7	Р	1	Total	С	Ν	Ο	Р	0	0
'	D	1	31	10	5	13	3	0	0

• Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	В	1	Total 17	$\begin{array}{c} \mathrm{C} \\ 4 \end{array}$	H 10	0 3	0	0



• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	351	Total O 351 351	0	0
9	В	245	Total O 245 245	0	0
9	С	22	TotalO2222	0	0
9	D	12	Total O 12 12	0	0
9	Е	4	Total O 4 4	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: RNA-dependent RNA polymerase

• Molecule 1: RNA-dependent RNA polymerase





P624 P624 VAL VAL THR ARG VAL VAL VAL VAL VAL CLU CLU CLU CLU	R642 T661 4662 4663 4673 4673 8677 8677 759	D696 T703 1722	V750 V747 V750	G755 D774 K775 R783	S787 E803 M812 R813	
0818 1821 1821 1821 1821 1821 1832 1832 1	Y919 M922 M927 L959 F967 D1011	A1013 V1022 P1031 F1047 F1075	K1089 E1090 R1091 L1107 V1111	V1115 S1118 K1119 Q1120	61139 61139 91146 91148	
H1183 M1185 M1185 M1186 M1188 A1188 A1188 A1188 A1188 L1X8 E1193 E1193 E1193 E1193 E1193 E1193 E1193 E1193 E1193	R1240 H1244 LN244 LN244 LN244 ARC ARC ARC ARC ARC ARC ARC ARC ARC ARC	71264 71265 71266 71266 71270 812 812 812 812 812 812	SER SER GLY ALA ASN TYR ASN ASP ASP	11285 E1289 W1301	11310 11314 01343 M1348 M1348	
Molecule 2: BNV (2		*CD*IID* V D	9*CD*CD*	·OD*11D*0	₽)
• Molecule 2. RIVA (3	-n GIAIAI	OI UI AI	UI UI	GIUIC	A = G = G = A - 3)
Chain C:	64%		7%	29%	_	
A A A A A A A A A A A A A A A A A A A						
• Molecule 2: RNA (5	·-R(*GP*AP*AP	*CP*UP*AF	*CP*CP*	GP*UP*C	CP*GP*GP*A)-3')
Chain E: 14%	29%		57%		-	
● ● 0 0 0 0 0 0 0 0 0 0 0 0 0						
• Molecule 3: RNA (5	·-R(*UP*CP*CP*	*GP*AP*CP	*G)-3')			
Chain D					_	
Chain D:		100%				
● 5 8						



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	81.21Å 85.81Å 101.58Å	Depositor
a, b, c, α , β , γ	102.56° 104.05° 94.06°	Depositor
Bosolution (Å)	29.31 - 2.05	Depositor
Resolution (A)	29.31 - 2.05	EDS
% Data completeness	94.1 (29.31-2.05)	Depositor
(in resolution range)	94.1 (29.31-2.05)	EDS
R_{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.29 (at 2.06 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
P. P.	0.196 , 0.226	Depositor
n, n_{free}	0.196 , 0.226	DCC
R_{free} test set	7584 reflections (5.01%)	wwPDB-VP
Wilson B-factor $(Å^2)$	33.3	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36 , 49.7	EDS
L-test for $twinning^2$	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16072	wwPDB-VP
Average B, all atoms $(Å^2)$	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, PEG, MG, CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
	Unam	RMSZ # Z > 5		RMSZ	# Z > 5
1	А	0.43	0/7791	0.59	0/10535
1	В	0.42	0/7389	0.59	0/9985
2	С	0.44	0/237	0.82	0/367
2	Е	0.43	0/138	0.72	0/212
3	D	0.48	0/160	0.72	0/247
All	All	0.43	0/15715	0.60	0/21346

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	784[B]	SER	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	7600	0	7428	53	0
1	В	7213	0	7096	52	0
2	С	213	0	109	1	0
2	Ε	125	0	64	3	0
3	D	144	0	76	0	0
4	А	2	0	0	0	0
4	В	2	0	0	0	0
5	А	1	0	0	0	0
5	В	2	0	0	0	0
6	А	6	8	8	1	0
6	В	12	0	16	0	0
7	А	31	0	10	0	0
7	В	62	0	20	1	0
8	В	7	10	10	0	0
9	А	351	0	0	2	0
9	В	245	0	0	0	0
9	С	22	0	0	0	0
9	D	12	0	0	0	0
9	Е	4	0	0	0	0
All	All	16054	18	14837	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:450:VAL:HG21	1:B:472:PRO:HG2	1.55	0.86
1:A:1400:GLY:H	1:B:783:ARG:HH12	1.41	0.66
1:B:1075:PHE:CE1	1:B:1118:SER:HA	2.31	0.65
1:B:575:LEU:HG	1:B:576:VAL:HG23	1.77	0.65
1:A:1095:ILE:HA	6:A:2004:GOL:H31	1.80	0.64
1:A:528:ARG:NH1	9:A:2105:HOH:O	2.31	0.63
1:A:1402:TYR:HB2	1:B:677:SER:HA	1.81	0.63
1:A:940:SER:HA	1:A:986:LYS:HD2	1.84	0.59
1:B:1270:PRO:HG2	1:B:1289:GLU:HG3	1.85	0.58
1:A:1400:GLY:C	1:B:783:ARG:HH22	2.07	0.57
1:B:450:VAL:HG21	1:B:472:PRO:CG	2.31	0.56
1:B:584:PHE:CZ	1:B:673:GLN:HG3	2.41	0.56
1:A:1117:GLN:HG3	1:A:1122:ILE:HB	1.88	0.56
1:B:1111:VAL:O	1:B:1115:VAL:HG23	2.06	0.55
1:A:526:LEU:HD13	1:A:650[B]:MET:HE3	1.88	0.55



Atom-1	Atom-2	Interatomic	Clash
		distance (A)	overlap (Å)
1:B:1089:LYS:HG3	1:B:1107:LEU:HB3	1.89	0.53
1:B:896:TRP:HB3	1:B:1182:PHE:HE1	1.72	0.53
1:B:803:GLU:HG3	1:B:812:MET:HE1	1.90	0.53
1:B:1253:PRO:HG2	1:B:1256:VAL:HG23	1.89	0.53
1:A:1089:LYS:HA	1:A:1107:LEU:HD13	1.90	0.52
1:B:922:MET:HE2	1:B:1011:ASP:HB3	1.92	0.52
1:B:967:PHE:CD1	1:B:1031:PRO:HG3	2.44	0.52
2:E:7:C:H2'	2:E:8:C:C6	2.45	0.52
1:A:423:LEU:HD11	1:A:458:TYR:HE1	1.74	0.51
1:A:922[A]:MET:HE3	1:A:1011:ASP:HB3	1.92	0.50
1:A:450:VAL:HG11	1:A:471:LYS:HB2	1.93	0.50
1:B:541:PRO:HG3	1:B:609:LYS:HE2	1.92	0.50
1:B:1115:VAL:HG11	7:B:1509:GTP:C3'	2.42	0.50
1:A:803:GLU:N	1:A:812[B]:MET:HE3	2.26	0.49
1:A:413:PRO:HG2	1:A:481:MET:HE1	1.94	0.49
1:A:694:LYS:HG2	1:A:695:THR:HG23	1.94	0.49
1:B:1134:ARG:HA	1:B:1139:GLY:HA2	1.94	0.49
1:A:741:SER:HB3	1:A:770:CYS:HB2	1.94	0.49
1:B:812:MET:HE3	1:B:813:ARG:N	2.28	0.49
1:A:534:ARG:HD2	1:A:642:ARG:HD2	1.95	0.48
1:B:959:LEU:HD13	1:B:1022:VAL:HG22	1.94	0.48
1:A:406:PRO:HB2	1:A:408:TRP:CD1	2.48	0.48
1:A:450:VAL:HG11	1:A:472:PRO:HD2	1.95	0.48
1:A:1379:ASP:HB3	1:A:1382:VAL:HB	1.95	0.47
1:A:709:ASP:OD1	1:A:764:SER:HB2	2.15	0.47
1:B:922:MET:SD	1:B:1013:ALA:HB2	2.53	0.47
1:A:543:PRO:HB2	1:A:561:VAL:HG13	1.97	0.47
1:B:750:VAL:HG22	1:B:927:TRP:CE2	2.49	0.47
1:B:774:ASP:OD1	1:B:775:LYS:N	2.47	0.47
1:A:522[B]:GLN:HB3	1:A:678:LYS:HB3	1.97	0.46
1:A:1348:MET:HE1	1:A:1353:TYR:HA	1.98	0.46
1:B:529:ARG:O	1:B:529:ARG:NH1	2.36	0.46
1:B:407:LYS:HA	1:B:410:HIS:CD2	2.51	0.46
1:B:1254:TYR:HB3	1:B:1255:PRO:HD3	1.96	0.46
1:B:1270:PRO:HB3	1:B:1285:ILE:CD1	2.46	0.45
1:A:526:LEU:HD13	1:A:650[B]:MET:CE	2.45	0.45
1:A:799:LEU:HB2	1:A:800:PRO:HD3	1.98	0.45
1:A:611[A]:ARG:NH2	9:A:2118:HOH:O	2.44	0.45
1:B:1310:PHE:O	1:B:1314:TYR:HB3	2.17	0.45
1:A:1392[A]:ASP:OD2	2:E:9:G:O3'	2.33	0.45
1:A:580:TRP:HB3	1:A:614:PHE:HB3	1.99	0.44



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:B:522:GLN:HE21	1:B:679:THR:C	2.20	0.44
1:B:832:LYS:HB2	1:B:832:LYS:HE2	1.72	0.44
1:A:605:LYS:HE2	2:C:5:U:O4	2.18	0.44
1:B:696:ASP:HB3	1:B:706:VAL:HG13	1.99	0.44
1:A:416:VAL:HG22	1:A:453:ILE:HD13	2.00	0.44
1:A:450:VAL:CG1	1:A:471:LYS:HB2	2.48	0.44
1:A:812[B]:MET:HE1	1:A:816:ILE:CD1	2.47	0.44
1:A:413:PRO:HD2	1:A:416:VAL:HB	1.99	0.44
1:A:495:SER:HA	1:A:537:GLU:O	2.18	0.44
1:B:722:ILE:HD13	1:B:735:VAL:HG11	1.99	0.43
1:B:803:GLU:HG3	1:B:812:MET:CE	2.48	0.43
1:B:534:ARG:HD2	1:B:642:ARG:HD2	2.01	0.43
1:A:710:GLY:HA3	1:A:1009:ASP:HB3	2.01	0.43
1:B:662:TRP:CZ3	1:B:663:GLN:HG2	2.54	0.43
1:B:1089:LYS:HA	1:B:1107:LEU:HD13	2.01	0.43
1:A:679:THR:HB	1:A:781:GLU:HB3	2.01	0.43
1:B:890:TYR:CE2	1:B:894:ILE:HD11	2.54	0.42
1:A:1214:SER:HB2	1:A:1221:ALA:HA	2.01	0.42
1:B:860:ARG:HA	1:B:1354:ALA:HB2	2.00	0.42
1:B:919:TYR:CD2	2:E:10[B]:U:H5'	2.55	0.42
1:B:544:THR:O	1:B:545:SER:C	2.58	0.42
1:A:485:PHE:CD2	1:A:492:VAL:HB	2.54	0.42
1:A:860:ARG:HD2	1:B:1343:GLY:HA3	2.02	0.42
1:A:960:VAL:HA	1:A:1014:TRP:O	2.20	0.42
1:B:580:TRP:HB3	1:B:614:PHE:HB3	2.01	0.41
1:B:1091:ARG:HG2	1:B:1146:PRO:HD3	2.01	0.41
1:A:1159:PRO:HB3	1:A:1164:ASP:HB3	2.02	0.41
1:B:1270:PRO:HG3	1:B:1301:TRP:CE3	2.56	0.41
1:A:796:LEU:O	1:A:800:PRO:HD3	2.20	0.41
1:A:1129:TRP:CE2	1:A:1133:ARG:HD2	2.56	0.41
1:A:1310:PHE:HA	1:A:1321:VAL:HG21	2.01	0.41
1:A:428:LYS:HA	1:A:428:LYS:HD2	1.83	0.41
1:A:1030:MET:HA	1:A:1031:PRO:HD3	1.93	0.41
1:B:1270:PRO:HG3	1:B:1301:TRP:CD2	2.56	0.41
1:A:920:ILE:HG22	1:A:988:VAL:HA	2.03	0.41
1:A:1380:PRO:HG3	1:B:787:SER:HB2	2.02	0.41
1:B:818:ASP:O	1:B:821:ILE:HG22	2.21	0.40
1:A:617:GLU:HA	1:A:645:PHE:O	2.21	0.40
1:A:1402:TYR:HE1	1:B:586:LYS:HD2	1.86	0.40
1:B:407:LYS:HG2	1:B:410:HIS:CE1	2.56	0.40
1:A:394:VAL:HG12	1:A:553:VAL:HG22	2.04	0.40



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
1:A:1402:TYR:CD1	1:B:673:GLN:HG2	2.57	0.40	
1:B:661:THR:HA	1:B:1047:PHE:HB3	2.03	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	935/1026~(91%)	920~(98%)	15 (2%)	0	100	100
1	В	884/1026~(86%)	871 (98%)	13~(2%)	0	100	100
All	All	1819/2052~(89%)	1791 (98%)	28 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	813/894~(91%)	791~(97%)	22 (3%)	44	38	
1	В	770/894~(86%)	758~(98%)	12 (2%)	62	59	
All	All	1583/1788~(88%)	1549~(98%)	34 (2%)	53	48	

All (34) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	394	VAL
1	А	439	LYS
1	А	443	SER
1	А	445	SER
1	А	475	ASP
1	А	505	SER
1	А	520	PHE
1	А	547	SER
1	А	554	VAL
1	А	555	SER
1	А	620	ILE
1	А	716	ARG
1	А	743[A]	LYS
1	А	743[B]	LYS
1	А	783	ARG
1	А	920	ILE
1	А	1152	SER
1	А	1177	LYS
1	А	1213	ILE
1	А	1243	LYS
1	А	1282	SER
1	А	1317	SER
1	В	428	LYS
1	В	475	ASP
1	В	553	VAL
1	В	608	ILE
1	В	812	MET
1	В	821	ILE
1	В	870	SER
1	В	1120	GLN
1	В	1214	SER
1	В	1217	SER
1	В	1240	ARG
1	В	1348	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	484	ASN
1	А	831	GLN
1	А	1335	GLN
1	В	400	ASN
1	В	410	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	В	522	GLN
1	В	1184	ASN
1	В	1323	GLN

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	С	9/14~(64%)	0	0
2	Е	5/14~(35%)	0	0
3	D	5/7~(71%)	0	0
All	All	19/35~(54%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 7 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Link Bond lengths			B	ond ang	les	
	туре	Unam	nes	LIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	В	1507	-	$5,\!5,\!5$	0.08	0	$5,\!5,\!5$	0.30	0
7	GTP	А	2005	4	25,33,34	1.09	2 (8%)	29,52,54	0.81	1 (3%)



Mol Type Chain		Dec	Link	Bond lengths			Bond angles			
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	В	1506	-	$5,\!5,\!5$	0.07	0	$5,\!5,\!5$	0.34	0
7	GTP	В	1509	5,4	$25,\!33,\!34$	1.07	2 (8%)	29,52,54	0.84	0
7	GTP	В	1502	-	25,33,34	1.02	2 (8%)	29,52,54	0.77	0
8	PEG	В	1501	-	$6,\!6,\!6$	0.12	0	5,5,5	0.07	0
6	GOL	А	2004	-	$5,\!5,\!5$	0.08	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	В	1507	-	-	1/4/4/4	-
7	GTP	А	2005	4	-	3/18/34/38	0/3/3/3
6	GOL	В	1506	-	-	4/4/4/4	-
7	GTP	В	1509	5,4	-	4/18/34/38	0/3/3/3
7	GTP	В	1502	-	-	7/18/34/38	0/3/3/3
8	PEG	В	1501	-	-	2/4/4/4	-
6	GOL	А	2004	-	-	4/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	В	1509	GTP	C5-C6	-2.71	1.41	1.47
7	А	2005	GTP	C5-C6	-2.69	1.41	1.47
7	В	1502	GTP	C5-C6	-2.63	1.42	1.47
7	А	2005	GTP	C8-N7	-2.18	1.31	1.35
7	В	1509	GTP	C8-N7	-2.11	1.31	1.35
7	В	1502	GTP	C8-N7	-2.09	1.31	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
7	А	2005	GTP	O6-C6-C5	2.02	128.32	124.37

There are no chirality outliers.

All (25) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	А	2004	GOL	O1-C1-C2-C3
6	В	1506	GOL	C1-C2-C3-O3
7	В	1502	GTP	C5'-O5'-PA-O2A
7	В	1502	GTP	O4'-C4'-C5'-O5'
7	В	1509	GTP	C3'-C4'-C5'-O5'
7	В	1509	GTP	O4'-C4'-C5'-O5'
8	В	1501	PEG	O1-C1-C2-O2
6	А	2004	GOL	C1-C2-C3-O3
6	В	1506	GOL	O1-C1-C2-C3
6	В	1507	GOL	C1-C2-C3-O3
8	В	1501	PEG	O2-C3-C4-O4
6	В	1506	GOL	O1-C1-C2-O2
7	А	2005	GTP	C3'-C4'-C5'-O5'
7	В	1502	GTP	C3'-C4'-C5'-O5'
6	А	2004	GOL	O1-C1-C2-O2
6	В	1506	GOL	O2-C2-C3-O3
7	А	2005	GTP	PB-O3A-PA-O5'
7	В	1502	GTP	PB-O3A-PA-O5'
7	В	1509	GTP	PB-O3A-PA-O5'
7	В	1502	GTP	C5'-O5'-PA-O3A
7	В	1502	GTP	C5'-O5'-PA-O1A
7	А	2005	GTP	O4'-C4'-C5'-O5'
7	В	1509	GTP	PB-O3A-PA-O1A
6	А	2004	GOL	O2-C2-C3-O3
7	В	1502	GTP	PB-O3A-PA-O2A

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	В	1509	GTP	1	0
6	А	2004	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ >	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q < 0.9
1	А	951/1026~(92%)	0.07	37 (3%) 39 42	22, 42, 77, 118	5~(0%)
1	В	906/1026~(88%)	0.13	31 (3%) 45 49	22, 45, 80, 100	0
2	С	10/14~(71%)	-0.62	0 100 100	27, 43, 76, 101	0
2	Е	6/14~(42%)	1.13	1 (16%) 1 1	61, 89, 98, 120	2(33%)
3	D	7/7~(100%)	0.34	1 (14%) 2 2	27, 46, 103, 130	0
All	All	1880/2087~(90%)	0.10	70 (3%) 41 45	22, 43, 80, 130	7~(0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	391	ALA	6.0
1	А	1383	TYR	4.7
1	В	506	PRO	4.6
1	В	548	PRO	4.6
1	В	553	VAL	4.3
1	В	753	THR	4.0
1	В	552	PRO	3.9
1	В	1187	LYS	3.9
1	А	455	LYS	3.7
1	А	1154	LEU	3.7
1	А	1401	ASP	3.6
1	А	1382	VAL	3.5
3	D	1	U	3.5
1	А	392	PRO	3.4
1	А	461	ASP	3.4
1	В	638	PRO	3.4
1	А	638	PRO	3.3
1	А	1250	SER	3.2
1	В	703	THR	3.2
1	А	458	TYR	3.2



Mol	Chain	Res	Type	RSRZ
1	В	1370	LEU	3.1
1	А	1183	1183 HIS	
1	В	1188	ALA	3.0
1	А	753	753 THR	
1	В	504	ASN	2.9
1	В	1253	PRO	2.9
1	А	1381	GLU	2.9
1	В	507	THR	2.9
1	А	620	ILE	2.9
1	В	620	ILE	2.8
1	В	1254	TYR	2.7
1	А	1196	ALA	2.7
1	В	560	ALA	2.6
2	Е	6	А	2.6
1	А	1252	ASP	2.5
1	А	604	PRO	2.5
1	А	1392[A]	ASP	2.5
1	В	538	ILE	2.5
1	В	1182	PHE	2.5
1	В	407	LYS	2.5
1	А	507	THR	2.4
1	А	1378	PRO	2.4
1	А	1241	LEU	2.4
1	А	1242	VAL	2.4
1	В	1186	MET	2.4
1	В	672	ILE	2.3
1	В	467	PRO	2.3
1	В	747	VAL	2.3
1	А	508	ALA	2.3
1	В	755	ASP	2.3
1	А	1244	ASN	2.3
1	А	438	LEU	2.3
1	В	746	TRP	2.3
1	А	1393[A]	GLY	2.2
1	A	1185	ALA	2.2
1	А	597	PHE	2.2
1	А	460	LEU	2.2
1	А	434	GLU	2.2
1	А	1399	ASN	2.2
1	А	1222	LEU	2.2
1	В	494	LEU	2.2
1	В	1189	ALA	2.2



Mol	Chain	Res	Type	RSRZ
1	В	1185	ALA	2.1
1	А	701	SER	2.1
1	В	1216	LYS	2.1
1	В	1184	ASN	2.1
1	А	1400	GLY	2.0
1	В	608	ILE	2.0
1	А	747	VAL	2.0
1	А	1379	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
7	GTP	В	1502	31/32	0.78	0.19	68,94,106,115	6
6	GOL	В	1507	6/6	0.84	0.15	45,53,57,58	0
6	GOL	В	1506	6/6	0.84	0.14	44,52,56,57	0
8	PEG	В	1501	7/7	0.86	0.33	67,80,90,90	0
6	GOL	А	2004	6/6	0.88	0.14	44,58,73,73	0
7	GTP	В	1509	31/32	0.96	0.11	40,80,94,99	0
5	MG	В	1508	1/1	0.97	0.14	$55,\!55,\!55,\!55$	0
5	MG	В	1505	1/1	0.98	0.13	32,32,32,32	0
5	MG	А	2003	1/1	0.98	0.09	22,22,22,22	0
7	GTP	А	2005	31/32	0.99	0.12	23,28,40,46	0
4	CA	А	2001	1/1	0.99	0.09	30,30,30,30	0
4	CA	А	2002	1/1	0.99	0.12	31,31,31,31	0
4	CA	В	1504	1/1	0.99	0.09	39,39,39,39	0
4	CA	В	1503	1/1	1.00	0.11	34,34,34,34	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

