



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 23, 2023 – 01:36 AM JST

PDB ID : 7Y96  
Title : Crystal structure of the carboxy-terminal domain of a coronavirus M protein fused with a split GFP  
Authors : Wang, X.; Sun, Z.; Zhou, X.  
Deposited on : 2022-06-24  
Resolution : 3.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1486 (3.50-3.34)
Clashscore	141614	1572 (3.50-3.34)
Ramachandran outliers	138981	1534 (3.50-3.34)
Sidechain outliers	138945	1535 (3.50-3.34)
RSRZ outliers	127900	1395 (3.50-3.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	
1	B	340	

## 2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 4890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein, Membrane protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	310	2432	1537	419	465	11	0	0	0
1	B	313	2458	1555	423	469	11	0	0	0

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	ARG	SER	engineered mutation	UNP P42212
A	39	ASN	TYR	engineered mutation	UNP P42212
A	64	LEU	PHE	engineered mutation	UNP P42212
A	66	CRO	SER	chromophore	UNP P42212
A	66	CRO	TYR	chromophore	UNP P42212
A	66	CRO	GLY	chromophore	UNP P42212
A	80	ARG	GLN	engineered mutation	UNP P42212
A	99	SER	PHE	engineered mutation	UNP P42212
A	105	THR	ASN	engineered mutation	UNP P42212
A	145	VAL	-	linker	UNP P42212
A	146	ASP	-	linker	UNP P42212
A	147	THR	-	linker	UNP P42212
A	148	MET	-	linker	UNP P42212
A	149	GLU	-	linker	UNP P42212
A	238	ALA	-	linker	UNP A3EXD6
A	239	ALA	-	linker	UNP A3EXD6
A	240	ALA	-	linker	UNP A3EXD6
A	248	THR	MET	engineered mutation	UNP P42212
A	258	ALA	VAL	engineered mutation	UNP P42212
A	266	VAL	ILE	engineered mutation	UNP P42212
A	301	VAL	ALA	engineered mutation	UNP P42212
A	326	GLU	-	expression tag	UNP P42212
A	327	ASN	-	expression tag	UNP P42212
A	328	LEU	-	expression tag	UNP P42212
A	329	TYR	-	expression tag	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	330	PHE	-	expression tag	UNP P42212
A	331	GLN	-	expression tag	UNP P42212
A	332	GLY	-	expression tag	UNP P42212
A	333	HIS	-	expression tag	UNP P42212
A	334	HIS	-	expression tag	UNP P42212
A	335	HIS	-	expression tag	UNP P42212
A	336	HIS	-	expression tag	UNP P42212
A	337	HIS	-	expression tag	UNP P42212
A	338	HIS	-	expression tag	UNP P42212
A	339	HIS	-	expression tag	UNP P42212
A	340	HIS	-	expression tag	UNP P42212
A	341	HIS	-	expression tag	UNP P42212
A	342	HIS	-	expression tag	UNP P42212
B	30	ARG	SER	engineered mutation	UNP P42212
B	39	ASN	TYR	engineered mutation	UNP P42212
B	64	LEU	PHE	engineered mutation	UNP P42212
B	66	CRO	SER	chromophore	UNP P42212
B	66	CRO	TYR	chromophore	UNP P42212
B	66	CRO	GLY	chromophore	UNP P42212
B	80	ARG	GLN	engineered mutation	UNP P42212
B	99	SER	PHE	engineered mutation	UNP P42212
B	105	THR	ASN	engineered mutation	UNP P42212
B	145	VAL	-	linker	UNP P42212
B	146	ASP	-	linker	UNP P42212
B	147	THR	-	linker	UNP P42212
B	148	MET	-	linker	UNP P42212
B	149	GLU	-	linker	UNP P42212
B	238	ALA	-	linker	UNP A3EXD6
B	239	ALA	-	linker	UNP A3EXD6
B	240	ALA	-	linker	UNP A3EXD6
B	248	THR	MET	engineered mutation	UNP P42212
B	258	ALA	VAL	engineered mutation	UNP P42212
B	266	VAL	ILE	engineered mutation	UNP P42212
B	301	VAL	ALA	engineered mutation	UNP P42212
B	326	GLU	-	expression tag	UNP P42212
B	327	ASN	-	expression tag	UNP P42212
B	328	LEU	-	expression tag	UNP P42212
B	329	TYR	-	expression tag	UNP P42212
B	330	PHE	-	expression tag	UNP P42212
B	331	GLN	-	expression tag	UNP P42212
B	332	GLY	-	expression tag	UNP P42212
B	333	HIS	-	expression tag	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	334	HIS	-	expression tag	UNP P42212
B	335	HIS	-	expression tag	UNP P42212
B	336	HIS	-	expression tag	UNP P42212
B	337	HIS	-	expression tag	UNP P42212
B	338	HIS	-	expression tag	UNP P42212
B	339	HIS	-	expression tag	UNP P42212
B	340	HIS	-	expression tag	UNP P42212
B	341	HIS	-	expression tag	UNP P42212
B	342	HIS	-	expression tag	UNP P42212



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.02Å 82.02Å 427.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.29 – 3.42 46.33 – 3.42	Depositor EDS
% Data completeness (in resolution range)	93.9 (38.29-3.42) 93.8 (46.33-3.42)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.01 (at 3.40Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.240 , 0.265 0.240 , 0.265	Depositor DCC
$R_{free}$ test set	555 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	70.8	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 22.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	4890	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/2457	0.49	0/3324
1	B	0.26	0/2484	0.48	0/3360
All	All	0.26	0/4941	0.49	0/6684

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2407	56	0
1	B	2458	0	2435	35	0
All	All	4890	0	4842	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:ASN:ND2	1:A:130:PHE:HB2	1.94	0.83
1:A:23:ASN:HD21	1:A:130:PHE:HB2	1.50	0.77
1:A:112:VAL:HG22	1:A:121:ASN:HD22	1.56	0.69
1:A:305:ASP:HB3	1:A:308:GLU:HB2	1.74	0.68
1:B:305:ASP:HB3	1:B:308:GLU:HB2	1.77	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/340 (89%)	294 (97%)	9 (3%)	0	100	100
1	B	306/340 (90%)	300 (98%)	6 (2%)	0	100	100
All	All	609/680 (90%)	594 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/291 (92%)	257 (96%)	11 (4%)	30	62
1	B	270/291 (93%)	264 (98%)	6 (2%)	52	78
All	All	538/582 (92%)	521 (97%)	17 (3%)	39	69

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	275	ASP
1	B	308	GLU
1	A	171	SER
1	A	210	LEU
1	A	307	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CRO	A	66	1	23,23,24	2.70	8 (34%)	30,32,34	2.85	8 (26%)
1	CRO	B	66	1	23,23,24	2.69	8 (34%)	30,32,34	2.79	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. <sup>1,2</sup> means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	66	1	-	6/12/31/32	0/2/2/2

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	B	66	1	-	2/12/31/32	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRO	C1-N2	7.36	1.43	1.32
1	B	66	CRO	C1-N2	7.33	1.43	1.32
1	B	66	CRO	CA2-C2	5.95	1.54	1.48
1	A	66	CRO	CA2-C2	5.74	1.54	1.48
1	A	66	CRO	C1-N3	5.16	1.45	1.37

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CRO	O2-C2-CA2	-8.95	125.93	130.96
1	B	66	CRO	O2-C2-CA2	-8.76	126.04	130.96
1	B	66	CRO	CA2-C2-N3	8.39	107.34	103.37
1	A	66	CRO	CA2-C2-N3	8.13	107.22	103.37
1	B	66	CRO	C2-N3-C1	-5.39	105.24	107.97

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CRO	N1-CA1-CB1-CG1
1	A	66	CRO	C1-CA1-CB1-CG1
1	A	66	CRO	C1-CA1-CB1-OG1
1	A	66	CRO	C3-CA3-N3-C2
1	B	66	CRO	C3-CA3-N3-C2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRO	5	0
1	B	66	CRO	4	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/340 (90%)	0.43	8 (2%) 56 54	47, 70, 101, 117	0
1	B	312/340 (91%)	0.27	5 (1%) 72 71	39, 56, 78, 100	0
All	All	621/680 (91%)	0.35	13 (2%) 63 63	39, 64, 94, 117	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	235	ALA	4.8
1	A	30	ARG	3.4
1	B	248	THR	3.2
1	B	277	TYR	2.8
1	B	233	TYR	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CRO	A	66	22/23	0.91	0.31	49,62,70,73	0
1	CRO	B	66	22/23	0.95	0.27	40,42,45,49	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.