



wwPDB EM Validation Summary Report ⓘ

May 27, 2026 – 12:55 PM EDT

PDB ID : 9YM8 / pdb_00009ym8
EMDB ID : EMD-73098
Title : State 2 focused on PHD FYR of MLL4FC bound to a nucleosome premodified
with H2BK120ub and H4K16ac
Authors : Sun, J.; Roeder, R.
Deposited on : 2025-10-09
Resolution : 3.43 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13927 (2.93 - 3.93)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	147	96% ..
2	J	147	97% ..
3	O	76	5% 97% .
3	U	76	96% 93% 5% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	A	135	
4	E	135	
5	B	102	
5	F	102	
6	C	129	
6	G	129	
7	D	125	
7	H	125	
8	R	334	
9	N	538	
10	K	1256	
11	T	628	
12	P	99	
12	Q	99	

2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 28174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	I	145	Total	C	N	O	P	0	0
			2957	1404	537	871	145		

- Molecule 2 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	J	145	Total	C	N	O	P	0	0
			2988	1414	560	869	145		

- Molecule 3 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	O	76	Total	C	N	O	S	0	0
			603	379	105	117	2		
3	U	76	Total	C	N	O	S	0	0
			603	379	105	117	2		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	76	CYS	GLY	engineered mutation	UNP P0CG47
U	76	CYS	GLY	engineered mutation	UNP P0CG47

- Molecule 4 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	A	97	Total	C	N	O	S	0	0
			801	505	155	137	4		
4	E	124	Total	C	N	O	S	0	0
			1008	630	199	175	4		

- Molecule 5 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	B	87	Total	C	N	O	S	0	0
			706	444	142	119	1		
5	F	84	Total	C	N	O	S	0	0
			673	424	133	115	1		

- Molecule 6 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	C	110	Total	C	N	O		0	0
			849	535	168	146			
6	G	104	Total	C	N	O		0	0
			805	508	157	140			

- Molecule 7 is a protein called Histone H2B type 2-E.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	D	96	Total	C	N	O	S	0	0
			755	474	138	141	2		
7	H	92	Total	C	N	O	S	0	0
			719	453	129	135	2		

- Molecule 8 is a protein called WD repeat-containing protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	R	300	Total	C	N	O	S	0	0
			2326	1485	388	444	9		

- Molecule 9 is a protein called Retinoblastoma-binding protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	N	414	Total	C	N	O	S	0	0
			3238	2036	550	636	16		

- Molecule 10 is a protein called [histone H3]-lysine(4) N-methyltransferase,Histone-lysine N-methyltransferase 2D.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	K	732	Total	C	N	O	S	0	0
			5825	3656	1032	1065	72		

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	1348	MET	-	expression tag	UNP A0A6P6IDI8
K	1349	ASP	-	expression tag	UNP A0A6P6IDI8
K	1350	TYR	-	expression tag	UNP A0A6P6IDI8
K	1351	LYS	-	expression tag	UNP A0A6P6IDI8
K	1352	ASP	-	expression tag	UNP A0A6P6IDI8
K	1353	ASP	-	expression tag	UNP A0A6P6IDI8
K	1354	ASP	-	expression tag	UNP A0A6P6IDI8
K	1355	ASP	-	expression tag	UNP A0A6P6IDI8
K	1356	LYS	-	expression tag	UNP A0A6P6IDI8
K	1357	SER	-	expression tag	UNP A0A6P6IDI8

- Molecule 11 is a protein called Set1/Ash2 histone methyltransferase complex subunit ASH2.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	T	301	Total	C	N	O	S	0	0
			2421	1566	400	447	8		


- Molecule 12 is a protein called Protein dpy-30 homolog.

Mol	Chain	Residues	Atoms				AltConf	Trace
12	P	54	Total	C	N	O	0	0
			431	279	73	79		
12	Q	54	Total	C	N	O	0	0
			432	279	73	80		

- Molecule 13 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
13	K	8	Total	Zn	0
			8	8	

- Molecule 14 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
14	K	1	Total	C	N	O	S	0
			26	14	6	5	1	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (145-MER)

Chain I:  96% ..




- Molecule 2: DNA (145-MER)

Chain J:  97% ..




- Molecule 3: Ubiquitin

Chain O:  97% .




- Molecule 3: Ubiquitin

Chain U:  96% 93% 5% .





- Molecule 4: Histone H3.1



Chain A:  70% 28%





- Molecule 4: Histone H3.1





- Molecule 5: Histone H4




- Molecule 5: Histone H4




- Molecule 6: Histone H2A type 1-B/E



- Molecule 6: Histone H2A type 1-B/E



- Molecule 7: Histone H2B type 2-E



- Molecule 7: Histone H2B type 2-E



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	46989	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	61.52	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.854	Depositor
Minimum map value	-0.947	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.040	Depositor
Recommended contour level	0.2	Depositor
Map size (\AA)	440.32, 440.32, 440.32	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.86, 0.86, 0.86	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, ZN, ALY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	I	0.26	0/3313	0.53	0/5107
2	J	0.27	0/3355	0.50	0/5180
3	O	0.18	0/609	0.37	0/819
3	U	0.18	0/609	0.34	0/819
4	A	0.15	0/813	0.31	0/1090
4	E	0.18	0/981	0.49	2/1308 (0.2%)
5	B	0.15	0/702	0.34	0/937
5	F	0.19	0/680	0.41	0/908
6	C	0.18	0/859	0.38	0/1157
6	G	0.18	0/815	0.38	0/1100
7	D	0.20	0/766	0.47	0/1026
7	H	0.19	0/730	0.44	0/982
8	R	0.15	0/2382	0.35	0/3231
9	N	0.16	0/3304	0.37	0/4489
10	K	0.16	0/5960	0.39	0/8061
11	T	0.17	0/2500	0.39	0/3395
12	P	0.18	0/439	0.43	0/598
12	Q	0.20	0/440	0.43	0/598
All	All	0.20	0/29257	0.42	2/40805 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	10	SER	CA-C-N	9.49	136.27	121.19
4	E	10	SER	C-N-CA	9.49	136.27	121.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	2957	0	1628	2	0
2	J	2988	0	1627	2	0
3	O	603	0	631	1	0
3	U	603	0	631	4	0
4	A	801	0	839	1	0
4	E	1008	0	1066	7	0
5	B	706	0	758	2	0
5	F	673	0	722	4	0
6	C	849	0	915	2	0
6	G	805	0	861	2	0
7	D	755	0	781	3	0
7	H	719	0	737	1	0
8	R	2326	0	2309	2	0
9	N	3238	0	3163	4	0
10	K	5825	0	5652	10	0
11	T	2421	0	2346	7	0
12	P	431	0	448	0	0
12	Q	432	0	448	3	0
13	K	8	0	0	0	0
14	K	26	0	19	0	0
All	All	28174	0	25581	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:N:201:ILE:HD12	9:N:215:ILE:HD11	1.78	0.65
10:K:5125:PHE:CZ	10:K:5131:MET:HE1	2.33	0.63
2:J:83:DT:OP2	4:A:46:VAL:HG13	2.05	0.57
5:B:73:THR:HG21	5:B:81:VAL:HG12	1.88	0.56
3:U:9:THR:HG23	3:U:11:LYS:H	1.72	0.55

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	O	74/76 (97%)	73 (99%)	1 (1%)	0	100	100
3	U	74/76 (97%)	72 (97%)	2 (3%)	0	100	100
4	A	95/135 (70%)	93 (98%)	2 (2%)	0	100	100
4	E	117/135 (87%)	107 (92%)	9 (8%)	1 (1%)	14	45
5	B	85/102 (83%)	80 (94%)	5 (6%)	0	100	100
5	F	82/102 (80%)	78 (95%)	4 (5%)	0	100	100
6	C	108/129 (84%)	106 (98%)	2 (2%)	0	100	100
6	G	102/129 (79%)	101 (99%)	1 (1%)	0	100	100
7	D	94/125 (75%)	89 (95%)	5 (5%)	0	100	100
7	H	90/125 (72%)	87 (97%)	3 (3%)	0	100	100
8	R	298/334 (89%)	280 (94%)	18 (6%)	0	100	100
9	N	408/538 (76%)	364 (89%)	42 (10%)	2 (0%)	24	56
10	K	726/1256 (58%)	675 (93%)	50 (7%)	1 (0%)	48	78
11	T	299/628 (48%)	277 (93%)	22 (7%)	0	100	100
12	P	52/99 (52%)	48 (92%)	4 (8%)	0	100	100
12	Q	52/99 (52%)	50 (96%)	2 (4%)	0	100	100
All	All	2756/4088 (67%)	2580 (94%)	172 (6%)	4 (0%)	49	78

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	E	24	ALA
9	N	359	LYS
9	N	365	THR
10	K	5514	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	O	69/69 (100%)	68 (99%)	1 (1%)	59	71
3	U	69/69 (100%)	68 (99%)	1 (1%)	59	71
4	A	85/107 (79%)	84 (99%)	1 (1%)	63	73
4	E	100/107 (94%)	99 (99%)	1 (1%)	68	75
5	B	71/77 (92%)	69 (97%)	2 (3%)	38	61
5	F	69/77 (90%)	69 (100%)	0	100	100
6	C	86/99 (87%)	84 (98%)	2 (2%)	44	64
6	G	83/99 (84%)	83 (100%)	0	100	100
7	D	82/105 (78%)	80 (98%)	2 (2%)	43	64
7	H	78/105 (74%)	78 (100%)	0	100	100
8	R	262/291 (90%)	256 (98%)	6 (2%)	44	64
9	N	362/462 (78%)	358 (99%)	4 (1%)	65	74
10	K	642/1091 (59%)	634 (99%)	8 (1%)	63	73
11	T	259/514 (50%)	253 (98%)	6 (2%)	44	64
12	P	48/89 (54%)	48 (100%)	0	100	100
12	Q	48/89 (54%)	48 (100%)	0	100	100
All	All	2413/3450 (70%)	2379 (99%)	34 (1%)	57	71

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	T	344	ILE
11	T	425	ILE
11	T	572	CYS
8	R	155	ILE
8	R	140	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
8	R	298	GLN
9	N	291	HIS
12	P	49	GLN
10	K	5105	ASN
11	T	605	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ALY	E	18	4	10,11,12	0.42	0	7,12,14	0.53	0
4	ALY	E	14	4	10,11,12	0.44	0	7,12,14	0.47	0
4	ALY	E	23	4	10,11,12	0.47	0	7,12,14	0.16	0
5	ALY	B	16	5	10,11,12	0.50	0	7,12,14	0.94	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ALY	E	18	4	-	2/9/10/12	-
4	ALY	E	14	4	-	1/9/10/12	-
4	ALY	E	23	4	-	6/9/10/12	-
5	ALY	B	16	5	-	3/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	16	ALY	CE-NZ-CH	2.25	125.88	122.56

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	23	ALY	N-CA-CB-CG
4	E	23	ALY	C-CA-CB-CG
5	B	16	ALY	CG-CD-CE-NZ
5	B	16	ALY	OH-CH-NZ-CE
5	B	16	ALY	CH3-CH-NZ-CE

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	18	ALY	1	0
4	E	23	ALY	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
14	SAH	K	5609	-	27,28,28	0.68	0	36,40,40	0.75	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	SAH	K	5609	-	-	5/15/31/31	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	K	5609	SAH	C3'-C2'-C1'	2.24	105.69	101.46

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	K	5609	SAH	O4'-C4'-C5'-SD
14	K	5609	SAH	C3'-C4'-C5'-SD
14	K	5609	SAH	CA-CB-CG-SD
14	K	5609	SAH	OXT-C-CA-CB
14	K	5609	SAH	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-73098. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 256




Y Index: 256



Z Index: 256


6.2.2 Raw map



X Index: 256



Y Index: 256



Z Index: 256


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 234



Y Index: 272




Z Index: 265


6.3.2 Raw map



X Index: 262



Y Index: 272




Z Index: 265


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 340 nm^3 ; this corresponds to an approximate mass of 307 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.292 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.292 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.43	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.38	9.35	6.04

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.38 differs from the reported value 3.43 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-73098 and PDB model 9YM8. Per-residue inclusion information can be found in section [3](#) on page [8](#).


9.1 Map-model overlay [i](#)



X



Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).







































9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8980	 0.4160
A	 0.9620	 0.5300
B	 0.9020	 0.5220
C	 0.9550	 0.5410
D	 0.9520	 0.5260
E	 0.9060	 0.4820
F	 0.9430	 0.5280
G	 0.9630	 0.5210
H	 0.9670	 0.5190
I	 0.9770	 0.4020
J	 0.9760	 0.3990
K	 0.8700	 0.3640
N	 0.8970	 0.4600
O	 0.8130	 0.2970
P	 0.6860	 0.1760
Q	 0.6930	 0.2070
R	 0.9720	 0.4880
T	 0.8870	 0.3540
U	 0.0710	 0.1560

