



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2023 – 07:59 AM EDT

PDB ID : 1YQU
Title : Escherichia coli purine nucleoside phosphorylase II, the product of the xapA gene
Authors : Dandanell, G.; Szczepanowski, R.H.; Kierdaszuk, B.; Shugar, D.; Bochtler, M.
Deposited on : 2005-02-02
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	 2% 71% 14% • 12%
1	B	277	 3% 71% 13% • 12%
1	C	277	 2% 73% 13% • 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	500	-	-	-	X
2	PO4	B	501	-	X	-	X
2	PO4	C	502	-	X	-	X
3	GUN	B	401	-	-	-	X
3	GUN	C	402	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 5601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xanthosine phosphorylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	243	Total 1851	C 1190	N 312	O 335	S 14	0	0	0
1	B	243	Total 1851	C 1190	N 312	O 335	S 14	0	0	0
1	C	243	Total 1851	C 1190	N 312	O 335	S 14	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	Total 5	O 4	P 1	0	0
2	B	1	Total 5	O 4	P 1	0	0
2	C	1	Total 5	O 4	P 1	0	0

- Molecule 3 is GUANINE (three-letter code: GUN) (formula: C₅H₅N₅O).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			11	5	5	1		
3	B	1	Total	C	N	O	0	0
			11	5	5	1		
3	C	1	Total	C	N	O	0	0
			11	5	5	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Xanthosine phosphorylase



- Molecule 1: Xanthosine phosphorylase



- Molecule 1: Xanthosine phosphorylase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.34Å 71.34Å 267.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 3.10 24.55 – 3.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.00-3.10) 99.9 (24.55-3.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 3.11Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.245 , 0.267 0.236 , 0.247	Depositor DCC
R_{free} test set	749 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	59.5	Xtrriage
Anisotropy	0.348	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.048 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5601	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GUN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/1888	0.86	0/2553
1	B	0.65	0/1888	0.86	0/2553
1	C	0.65	0/1888	0.86	0/2553
All	All	0.65	0/5664	0.86	0/7659

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1851	0	1866	43	0
1	B	1851	0	1866	51	1
1	C	1851	0	1866	36	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	11	0	5	1	0
3	B	11	0	5	1	0
3	C	11	0	5	1	0
All	All	5601	0	5613	112	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204:MET:HE2	1:B:205:MET:N	1.55	1.21
1:B:204:MET:HE2	1:B:205:MET:CA	1.72	1.17
1:C:175:VAL:HG12	1:C:179:GLU:OE1	1.58	1.03
1:B:175:VAL:HG12	1:B:179:GLU:OE1	1.58	1.02
1:B:204:MET:CE	1:B:205:MET:HA	1.91	1.00
1:A:175:VAL:HG12	1:A:179:GLU:OE1	1.65	0.96
1:C:175:VAL:O	1:C:179:GLU:HG3	1.69	0.90
1:B:175:VAL:O	1:B:179:GLU:HG3	1.72	0.89
1:A:175:VAL:O	1:A:179:GLU:HG3	1.70	0.89
1:B:204:MET:HE2	1:B:205:MET:HA	1.45	0.88
1:A:208:ILE:HD13	1:C:208:ILE:HD11	1.57	0.85
1:B:30:PHE:HB2	1:B:81:CYS:SG	2.16	0.84
1:A:69:LEU:HD12	1:A:81:CYS:O	1.77	0.84
1:A:204:MET:CG	1:C:208:ILE:HG13	2.08	0.83
1:A:175:VAL:CG1	1:A:179:GLU:OE1	2.26	0.83
1:A:204:MET:HG2	1:C:208:ILE:HG13	1.61	0.82
1:B:208:ILE:HG13	1:C:204:MET:CG	2.10	0.82
1:B:208:ILE:HG13	1:C:204:MET:HG3	1.61	0.81
1:B:175:VAL:CG1	1:B:179:GLU:OE1	2.31	0.78
1:C:162:ASN:HD21	1:C:166:ALA:H	1.32	0.78
1:C:175:VAL:CG1	1:C:179:GLU:OE1	2.30	0.78
1:B:162:ASN:HD21	1:B:166:ALA:H	1.33	0.77
1:B:204:MET:HE2	1:B:204:MET:C	2.05	0.77
1:A:162:ASN:HD21	1:A:166:ALA:H	1.33	0.76
1:A:178:GLU:O	1:A:179:GLU:HG2	1.86	0.75
1:C:178:GLU:O	1:C:179:GLU:HG2	1.86	0.75
1:B:178:GLU:O	1:B:179:GLU:HG2	1.86	0.74
1:B:71:LEU:HD13	1:B:80:VAL:HG23	1.70	0.73
1:B:204:MET:CE	1:B:204:MET:C	2.57	0.72
1:B:204:MET:CE	1:B:205:MET:N	2.45	0.72
1:A:71:LEU:HD13	1:A:80:VAL:HG23	1.69	0.72
1:B:178:GLU:C	1:B:179:GLU:CG	2.59	0.71
1:C:178:GLU:C	1:C:179:GLU:CG	2.59	0.71
1:B:104:PHE:O	1:B:109:CYS:SG	2.45	0.71
1:A:178:GLU:C	1:A:179:GLU:CG	2.59	0.71
1:B:204:MET:CE	1:B:205:MET:CA	2.51	0.69
1:C:71:LEU:HD13	1:C:80:VAL:HG23	1.75	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:ILE:CD1	1:C:208:ILE:HD11	2.24	0.67
1:B:139:THR:HB	1:B:190:SER:O	1.96	0.66
1:A:139:THR:HB	1:A:190:SER:O	1.97	0.65
1:C:139:THR:HB	1:C:190:SER:O	1.96	0.65
1:A:208:ILE:HD11	1:B:208:ILE:HD13	1.79	0.63
1:A:204:MET:HG3	1:C:208:ILE:HG13	1.79	0.62
1:B:208:ILE:HD11	1:C:208:ILE:HD13	1.82	0.62
1:B:208:ILE:HG13	1:C:204:MET:HG2	1.83	0.60
1:B:136:HIS:CD2	1:C:198:THR:HG21	2.38	0.59
1:A:94:MET:CE	1:A:140:MET:HB2	2.33	0.59
1:A:178:GLU:C	1:A:179:GLU:HG2	2.24	0.58
1:C:178:GLU:C	1:C:179:GLU:HG2	2.24	0.57
1:B:178:GLU:C	1:B:179:GLU:HG2	2.24	0.56
1:B:204:MET:HE2	1:B:205:MET:CG	2.36	0.56
1:B:204:MET:CE	1:B:205:MET:HG3	2.36	0.55
1:B:204:MET:CE	1:B:205:MET:CG	2.85	0.54
1:B:204:MET:HE2	1:B:205:MET:HG3	1.89	0.54
1:A:208:ILE:HD11	1:B:208:ILE:CD1	2.38	0.52
1:A:178:GLU:C	1:A:179:GLU:HG3	2.31	0.51
1:C:178:GLU:C	1:C:179:GLU:HG3	2.31	0.51
1:A:71:LEU:N	1:A:71:LEU:HD22	2.26	0.51
1:B:178:GLU:C	1:B:179:GLU:HG3	2.31	0.51
1:B:30:PHE:CB	1:B:81:CYS:SG	2.95	0.51
1:B:215:MET:HG3	3:B:401:GUN:HN22	1.76	0.50
1:A:181:PHE:HZ	1:A:268:LEU:HB2	1.76	0.50
1:A:198:THR:HG21	1:C:136:HIS:CD2	2.46	0.50
1:B:181:PHE:HZ	1:B:268:LEU:HB2	1.76	0.50
1:C:215:MET:HG3	3:C:402:GUN:HN22	1.76	0.50
1:A:204:MET:HE2	1:A:205:MET:HA	1.94	0.50
1:B:71:LEU:HD22	1:B:71:LEU:N	2.27	0.50
1:C:71:LEU:HD22	1:C:71:LEU:N	2.26	0.49
1:C:181:PHE:HZ	1:C:268:LEU:HB2	1.76	0.49
1:A:82:MET:HG3	1:A:104:PHE:HZ	1.78	0.48
1:B:178:GLU:O	1:B:179:GLU:CG	2.58	0.48
1:C:82:MET:HG3	1:C:104:PHE:HZ	1.77	0.48
1:A:215:MET:HG3	3:A:400:GUN:HN22	1.77	0.48
1:C:178:GLU:O	1:C:179:GLU:CG	2.58	0.47
1:B:208:ILE:CG1	1:C:204:MET:HG3	2.39	0.47
1:A:94:MET:HE2	1:A:140:MET:HB2	1.96	0.46
1:B:204:MET:O	1:B:204:MET:HE3	2.14	0.46
1:C:136:HIS:CD2	1:C:136:HIS:C	2.89	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:HIS:CD2	1:A:136:HIS:C	2.90	0.46
1:A:36:LEU:HA	1:A:36:LEU:HD23	1.66	0.45
1:B:36:LEU:HD23	1:B:36:LEU:HA	1.67	0.45
1:C:36:LEU:HA	1:C:36:LEU:HD23	1.66	0.45
1:B:136:HIS:CD2	1:B:136:HIS:C	2.89	0.45
1:A:30:PHE:HB2	1:A:81:CYS:SG	2.56	0.45
1:B:74:LEU:HD12	1:B:74:LEU:HA	1.85	0.45
1:A:136:HIS:HA	1:A:188:PHE:O	2.17	0.44
1:C:208:ILE:HD12	1:C:208:ILE:HA	1.76	0.44
1:C:201:GLU:O	1:C:204:MET:HB3	2.16	0.44
1:A:208:ILE:HG13	1:B:204:MET:CG	2.48	0.44
1:C:276:ILE:HG12	1:C:276:ILE:O	2.18	0.43
1:A:175:VAL:O	1:A:179:GLU:CG	2.56	0.43
1:A:276:ILE:O	1:A:276:ILE:HG12	2.18	0.43
1:A:94:MET:HE1	1:A:140:MET:HB2	2.00	0.43
1:B:276:ILE:O	1:B:276:ILE:HG12	2.18	0.43
1:B:37:GLY:O	1:B:39:LEU:N	2.52	0.43
1:C:122:ARG:HA	1:C:123:PRO:HD3	1.86	0.43
1:B:208:ILE:HD12	1:B:208:ILE:HA	1.76	0.42
1:A:178:GLU:O	1:A:179:GLU:CG	2.59	0.42
1:B:82:MET:HG3	1:B:104:PHE:HZ	1.83	0.42
1:C:37:GLY:O	1:C:39:LEU:N	2.52	0.42
1:A:162:ASN:HD21	1:A:166:ALA:N	2.11	0.42
1:A:201:GLU:O	1:A:204:MET:HB3	2.20	0.42
1:A:37:GLY:O	1:A:39:LEU:N	2.52	0.42
1:B:139:THR:O	1:B:139:THR:CG2	2.69	0.41
1:A:71:LEU:N	1:A:71:LEU:CD2	2.84	0.41
1:A:136:HIS:CD2	1:B:198:THR:HG21	2.56	0.41
1:B:162:ASN:HD21	1:B:166:ALA:N	2.10	0.41
1:A:195:ASN:HB3	1:C:158:PHE:CZ	2.56	0.40
1:C:175:VAL:O	1:C:179:GLU:CG	2.55	0.40
1:A:139:THR:O	1:A:139:THR:CG2	2.70	0.40
1:B:69:LEU:HD11	1:B:80:VAL:HG22	2.03	0.40
1:A:208:ILE:HG13	1:B:204:MET:HG2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ARG:NH2	1:B:180:GLY:O[4_466]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	237/277 (86%)	225 (95%)	7 (3%)	5 (2%)	7	30
1	B	237/277 (86%)	225 (95%)	7 (3%)	5 (2%)	7	30
1	C	237/277 (86%)	226 (95%)	6 (2%)	5 (2%)	7	30
All	All	711/831 (86%)	676 (95%)	20 (3%)	15 (2%)	7	30

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	34	SER
1	B	34	SER
1	C	34	SER
1	A	38	ALA
1	A	178	GLU
1	B	38	ALA
1	B	178	GLU
1	C	38	ALA
1	C	178	GLU
1	A	33	GLY
1	B	33	GLY
1	C	33	GLY
1	B	58	PRO
1	C	58	PRO
1	A	58	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/223 (89%)	180 (91%)	18 (9%)	9	33
1	B	198/223 (89%)	178 (90%)	20 (10%)	7	28
1	C	198/223 (89%)	180 (91%)	18 (9%)	9	33
All	All	594/669 (89%)	538 (91%)	56 (9%)	8	32

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	71	LEU
1	A	79	VAL
1	A	111	LEU
1	A	124	GLU
1	A	136	HIS
1	A	139	THR
1	A	145	MET
1	A	172	LEU
1	A	184	THR
1	A	189	VAL
1	A	204	MET
1	A	208	ILE
1	A	229	LEU
1	A	265	PHE
1	A	273	LEU
1	A	274	ARG
1	A	276	ILE
1	B	5	GLN
1	B	71	LEU
1	B	79	VAL
1	B	81	CYS
1	B	109	CYS
1	B	111	LEU
1	B	124	GLU
1	B	136	HIS
1	B	139	THR
1	B	145	MET
1	B	172	LEU
1	B	184	THR
1	B	189	VAL
1	B	204	MET
1	B	208	ILE
1	B	229	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	265	PHE
1	B	273	LEU
1	B	274	ARG
1	B	276	ILE
1	C	5	GLN
1	C	71	LEU
1	C	79	VAL
1	C	111	LEU
1	C	124	GLU
1	C	136	HIS
1	C	139	THR
1	C	145	MET
1	C	172	LEU
1	C	184	THR
1	C	189	VAL
1	C	204	MET
1	C	208	ILE
1	C	229	LEU
1	C	265	PHE
1	C	273	LEU
1	C	274	ARG
1	C	276	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	162	ASN
1	A	267	ASN
1	B	75	GLN
1	B	162	ASN
1	B	267	ASN
1	C	75	GLN
1	C	162	ASN
1	C	267	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	C	502	-	4,4,4	2.45	1 (25%)	6,6,6	1.86	3 (50%)
2	PO4	A	500	-	4,4,4	2.46	1 (25%)	6,6,6	1.91	2 (33%)
3	GUN	A	400	-	7,12,12	2.53	4 (57%)	8,17,17	1.47	2 (25%)
3	GUN	C	402	-	7,12,12	2.49	4 (57%)	8,17,17	1.54	2 (25%)
3	GUN	B	401	-	7,12,12	2.55	4 (57%)	8,17,17	1.52	2 (25%)
2	PO4	B	501	-	4,4,4	2.66	1 (25%)	6,6,6	1.94	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GUN	A	400	-	-	-	0/2/2/2
3	GUN	C	402	-	-	-	0/2/2/2
3	GUN	B	401	-	-	-	0/2/2/2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	PO4	P-O1	5.23	1.63	1.50
2	A	500	PO4	P-O1	4.83	1.62	1.50
2	C	502	PO4	P-O1	4.78	1.62	1.50
3	B	401	GUN	C6-N1	4.29	1.44	1.37
3	A	400	GUN	C6-N1	4.28	1.44	1.37
3	C	402	GUN	C6-N1	4.00	1.43	1.37
3	C	402	GUN	O6-C6	3.92	1.31	1.23
3	A	400	GUN	O6-C6	3.55	1.30	1.23
3	B	401	GUN	O6-C6	3.53	1.30	1.23
3	A	400	GUN	C2-N1	2.69	1.44	1.37
3	B	401	GUN	C2-N1	2.63	1.44	1.37
3	C	402	GUN	C2-N1	2.41	1.43	1.37
3	B	401	GUN	C2-N3	2.40	1.39	1.33
3	C	402	GUN	C2-N3	2.13	1.38	1.33
3	A	400	GUN	C2-N3	2.10	1.38	1.33

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	402	GUN	O6-C6-C5	3.11	130.44	124.37
3	A	400	GUN	O6-C6-C5	3.06	130.34	124.37
3	B	401	GUN	O6-C6-C5	3.05	130.33	124.37
2	B	501	PO4	O4-P-O1	-2.74	100.86	110.89
2	A	500	PO4	O4-P-O1	-2.67	101.13	110.89
2	C	502	PO4	O4-P-O1	-2.63	101.29	110.89
2	B	501	PO4	O3-P-O1	-2.45	101.94	110.89
2	A	500	PO4	O3-P-O1	-2.44	101.95	110.89
2	C	502	PO4	O3-P-O1	-2.29	102.53	110.89
3	C	402	GUN	C5-C6-N1	-2.27	109.95	113.95
3	A	400	GUN	C5-C6-N1	-2.22	110.04	113.95
2	C	502	PO4	O2-P-O1	2.06	118.42	110.89
3	B	401	GUN	C5-C6-N1	-2.03	110.37	113.95
2	B	501	PO4	O2-P-O1	2.02	118.27	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	GUN	1	0
3	C	402	GUN	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	GUN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/277 (87%)	-0.18	6 (2%) 57 34	16, 39, 61, 79	0
1	B	243/277 (87%)	-0.21	9 (3%) 41 21	16, 39, 62, 79	0
1	C	243/277 (87%)	-0.19	6 (2%) 57 34	16, 40, 62, 79	0
All	All	729/831 (87%)	-0.20	21 (2%) 51 28	16, 39, 63, 79	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	5	GLN	5.5
1	A	5	GLN	4.5
1	C	5	GLN	4.0
1	B	60	SER	3.5
1	B	151	ASP	3.5
1	C	34	SER	3.3
1	C	151	ASP	3.3
1	B	34	SER	3.2
1	A	60	SER	3.2
1	C	239	ASN	2.9
1	A	6	PHE	2.7
1	A	34	SER	2.6
1	B	179	GLU	2.5
1	C	179	GLU	2.5
1	B	124	GLU	2.4
1	B	239	ASN	2.2
1	B	148	LEU	2.2
1	A	151	ASP	2.2
1	B	23	ASP	2.2
1	A	179	GLU	2.1
1	C	60	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	500	5/5	0.43	0.99	115,115,116,116	0
2	PO4	B	501	5/5	0.49	0.79	115,115,115,116	0
2	PO4	C	502	5/5	0.58	0.67	114,114,115,115	0
3	GUN	B	401	11/11	0.69	0.71	84,85,86,86	0
3	GUN	C	402	11/11	0.74	0.63	84,85,86,86	0
3	GUN	A	400	11/11	0.81	0.45	84,84,84,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.