



Full wwPDB X-ray Structure Validation Report i

Oct 18, 2023 – 06:19 AM EDT

PDB ID : 1YXM
Title : Crystal structure of peroxisomal trans 2-enoyl CoA reductase
Authors : Jansson, A.; Ng, S.; Arrowsmith, C.; Sharma, S.; Edwards, A.M.; Von Delft, F.; Sundstrom, M.; Oppermann, U.; Structural Genomics Consortium (SGC)
Deposited on : 2005-02-22
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)


There are 5 unique types of molecules in this entry. The entry contains 8697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called peroxisomal trans 2-enoyl CoA reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	297	Total	C 2153	N 1376	O 366	S 403	8	0	1	0
1	B	283	Total	C 2102	N 1345	O 358	S 391	8	0	2	0
1	C	264	Total	C 1892	N 1210	O 322	S 352	8	0	0	0
1	D	291	Total	C 2141	N 1369	O 363	S 401	8	0	1	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	O 5	S 4	1	0	0
2	B	1	Total	O 5	S 4	1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is ADENINE (three-letter code: ADE) (formula: C₅H₅N₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C N 10 5 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	101	Total O 101 101	0	0
5	B	96	Total O 96 96	0	0
5	C	70	Total O 70 70	0	0
5	D	107	Total O 107 107	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: peroxisomal trans 2-enoyl CoA reductase




- Molecule 1: peroxisomal trans 2-enoyl CoA reductase



- Molecule 1: peroxisomal trans 2-enoyl CoA reductase



- Molecule 1: peroxisomal trans 2-enoyl CoA reductase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.15 Å 119.01 Å 119.91 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.00 – 1.90 47.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (47.00-1.90) 97.5 (47.34-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle^1$	2.72 (at 1.90 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R , R_{free}	0.204 , 0.245 0.212 , 0.247	Depositor DCC
R_{free} test set	4438 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.003 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8697	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PO4, ADE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	1/2200 (0.0%)	0.79	5/2996 (0.2%)
1	B	0.82	0/2152	0.76	1/2926 (0.0%)
1	C	0.75	2/1926 (0.1%)	0.83	5/2617 (0.2%)
1	D	0.89	4/2185 (0.2%)	0.93	6/2972 (0.2%)
All	All	0.82	7/8463 (0.1%)	0.83	17/11511 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	3
1	D	3	5
All	All	3	13

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	194	ILE	C-N	-15.25	0.98	1.34
1	D	195	ARG	C-N	9.89	1.56	1.34
1	A	111	GLN	C-N	-8.52	1.14	1.34
1	D	198	CYS	CB-SG	-6.13	1.71	1.82
1	D	195	ARG	CD-NE	-5.84	1.36	1.46
1	C	103	PHE	C-N	-5.58	1.21	1.34
1	C	285	GLY	C-N	-5.33	1.21	1.34

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	195	ARG	NE-CZ-NH2	-18.18	111.21	120.30
1	D	195	ARG	O-C-N	-17.14	95.28	122.70
1	D	195	ARG	NE-CZ-NH1	14.56	127.58	120.30
1	C	285	GLY	O-C-N	-10.11	106.53	122.70
1	C	285	GLY	C-N-CA	10.07	146.88	121.70
1	A	110	GLY	O-C-N	-8.39	109.28	122.70
1	D	195	ARG	C-N-CA	7.09	139.42	121.70
1	D	195	ARG	CA-C-N	7.01	132.63	117.20
1	A	110	GLY	C-N-CA	6.52	138.00	121.70
1	C	104	LEU	CB-CG-CD2	6.31	121.73	111.00
1	B	111	GLN	O-C-N	-6.23	112.74	122.70
1	C	285	GLY	CA-C-N	6.13	130.68	117.20
1	C	276	ASP	CB-CG-OD1	5.69	123.42	118.30
1	A	74	VAL	CB-CA-C	-5.45	101.04	111.40
1	D	260	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	60	ASP	CB-CG-OD2	5.28	123.06	118.30
1	A	110	GLY	CA-C-N	5.08	128.39	117.20

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	52	LEU	CA
1	D	53	GLU	CA
1	D	54	ARG	CA

All (13) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	110	GLY	Mainchain
1	A	97	THR	Peptide
1	A	98	PHE	Peptide
1	B	111	GLN	Mainchain
1	B	207	GLN	Peptide
1	C	51	LYS	Peptide
1	C	53	GLU	Peptide
1	C	6	LYS	Peptide
1	D	194	ILE	Mainchain
1	D	195	ARG	Mainchain
1	D	214	GLY	Peptide
1	D	51	LYS	Peptide
1	D	53	GLU	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2077	16	0
1	B	2102	0	2089	21	0
1	C	1892	0	1838	42	0
1	D	2141	0	2090	49	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	D	10	0	0	0	0
3	B	5	0	0	0	0
4	B	10	0	4	0	0
5	A	101	0	0	1	0
5	B	96	0	0	1	0
5	C	70	0	0	2	0
5	D	107	0	0	4	0
All	All	8697	0	8098	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (122) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:52:LEU:CA	1:D:53:GLU:HG3	1.65	1.26
1:C:52:LEU:HB3	1:C:53:GLU:HA	1.30	1.12
1:D:52:LEU:N	1:D:53:GLU:HA	1.49	1.09
1:B:208:THR:HG23	5:B:4085:HOH:O	1.51	1.08
1:C:52:LEU:HD23	1:C:55:LEU:H	1.15	1.08
1:D:52:LEU:H	1:D:53:GLU:CA	1.65	1.08
1:D:52:LEU:H	1:D:53:GLU:HA	0.89	1.01
1:C:52:LEU:HD21	1:C:56:LYS:H	1.22	1.01
1:C:52:LEU:CD2	1:C:55:LEU:N	2.24	1.00
1:C:124:TRP:HE1	1:D:133:THR:HG22	1.26	0.99
1:D:52:LEU:HA	1:D:53:GLU:HG3	1.00	0.99
1:C:52:LEU:HD23	1:C:55:LEU:N	1.81	0.95
1:D:52:LEU:HA	1:D:53:GLU:CG	1.95	0.95
1:C:52:LEU:HD21	1:C:56:LYS:N	1.82	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:52:LEU:CD2	1:C:55:LEU:H	1.88	0.85
1:D:154:ILE:HB	1:D:196:ILE:HG12	1.65	0.78
1:C:52:LEU:HD22	1:C:53:GLU:O	1.83	0.77
1:D:50:ARG:O	1:D:78:GLN:HG3	1.84	0.77
1:B:29:GLY:H	1:B:209:ALA:HA	1.49	0.77
1:C:52:LEU:CB	1:C:53:GLU:HA	2.13	0.76
1:D:52:LEU:CA	1:D:53:GLU:CG	2.58	0.74
1:C:60:ASP:O	1:C:64:ALA:HB2	1.88	0.73
1:B:125[B]:HIS:NE2	1:B:129:GLU:HG3	2.02	0.73
1:D:277:HIS:HD2	1:D:279:ASN:H	1.38	0.70
1:D:50:ARG:O	1:D:78:GLN:CG	2.40	0.69
1:D:52:LEU:H	1:D:53:GLU:CB	2.07	0.67
1:D:52:LEU:N	1:D:53:GLU:HG3	2.10	0.66
1:C:52:LEU:CD2	1:C:55:LEU:CA	2.74	0.66
1:C:55:LEU:O	1:C:56:LYS:C	2.35	0.64
1:D:52:LEU:N	1:D:53:GLU:CA	2.29	0.64
1:B:63:GLN:HE22	1:B:73:ARG:HA	1.62	0.64
1:D:109:GLY:HA3	1:D:131:ASN:OD1	1.98	0.64
1:D:289:VAL:HG12	1:D:293:MET:HE3	1.77	0.64
1:A:39:LEU:HD11	1:A:74:VAL:HG22	1.80	0.63
1:B:205:TYR:O	1:B:236:PRO:HD3	1.98	0.63
1:D:51:LYS:HA	1:D:52:LEU:HB3	1.81	0.63
1:D:49:SER:HB3	1:D:55:LEU:HG	1.82	0.61
1:C:52:LEU:CD2	1:C:56:LYS:N	2.61	0.61
1:C:52:LEU:HB3	1:C:53:GLU:CA	2.17	0.61
1:C:169:VAL:CG1	1:D:181:LEU:HD13	2.32	0.60
1:C:17:GLN:NE2	5:C:356:HOH:O	2.25	0.60
1:C:26:GLY:HA3	1:C:47:ILE:HG23	1.85	0.59
1:D:51:LYS:CA	1:D:52:LEU:HB3	2.34	0.58
1:D:182:THR:HG23	1:D:196:ILE:HG22	1.85	0.58
1:B:111:GLN:HE22	1:B:165:PHE:HE2	1.51	0.57
1:C:52:LEU:HD22	1:C:55:LEU:N	2.17	0.57
1:C:88:ASN:O	1:C:92:LYS:HG3	2.05	0.57
1:B:92:LYS:HE2	1:B:96:ASP:OD1	2.06	0.56
1:C:52:LEU:HD21	1:C:55:LEU:CA	2.34	0.56
1:D:289:VAL:HG12	1:D:293:MET:CE	2.36	0.55
1:C:169:VAL:CG1	1:D:181:LEU:CD1	2.85	0.55
1:D:19:GLN:NE2	5:D:2085:HOH:O	2.39	0.55
1:D:52:LEU:N	1:D:53:GLU:CG	2.70	0.54
1:B:125[B]:HIS:CD2	1:B:129:GLU:HG3	2.42	0.54
1:B:92:LYS:O	1:B:92:LYS:HE3	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:THR:HA	1:D:269:THR:HA	1.89	0.53
1:B:125[B]:HIS:NE2	1:B:129:GLU:CG	2.71	0.53
1:D:46:VAL:HG22	1:D:75:ILE:CG1	2.38	0.53
1:D:277:HIS:CD2	1:D:279:ASN:H	2.23	0.53
1:C:124:TRP:NE1	1:D:133:THR:HG22	2.10	0.53
1:A:210:VAL:HG12	1:A:213:TYR:HB2	1.90	0.53
1:C:290:VAL:HA	1:C:293:MET:HE2	1.91	0.52
1:D:117:GLU:HG3	1:D:118:HIS:CD2	2.45	0.52
1:D:280:TRP:CD2	1:D:281:PRO:HD2	2.45	0.52
1:A:77:ILE:HD13	1:A:93:SER:OG	2.10	0.51
1:D:73:ARG:NH1	5:D:2075:HOH:O	2.42	0.51
1:C:60:ASP:O	1:C:64:ALA:CB	2.56	0.50
1:D:46:VAL:HG22	1:D:75:ILE:HD11	1.92	0.50
1:C:52:LEU:CB	1:C:53:GLU:CA	2.86	0.49
1:C:55:LEU:O	1:C:58:ALA:N	2.39	0.49
1:A:95:LEU:O	1:A:99:GLY:CA	2.59	0.49
1:B:63:GLN:NE2	1:B:73:ARG:HA	2.29	0.48
1:C:52:LEU:HD22	1:C:53:GLU:C	2.33	0.48
1:C:166:PRO:O	1:C:167:LEU:HB2	2.13	0.48
1:B:280:TRP:CD2	1:B:281:PRO:HD2	2.48	0.48
1:A:125:HIS:O	1:A:129:GLU:HB2	2.14	0.47
1:A:94:THR:O	1:A:98:PHE:HB2	2.15	0.47
1:A:95:LEU:O	1:A:99:GLY:HA2	2.15	0.47
1:C:124:TRP:CE3	1:C:170:HIS:HB3	2.49	0.47
1:C:55:LEU:O	1:C:57:SER:N	2.47	0.46
1:B:27:ALA:HB2	1:B:55:LEU:HD23	1.97	0.46
1:D:63:GLN:HG2	5:D:2089:HOH:O	2.16	0.46
1:A:286:ASP:CA	5:A:405:HOH:O	2.64	0.46
1:A:35:VAL:HG11	1:A:62:LEU:HD13	1.98	0.45
1:B:124:TRP:CE3	1:B:170:HIS:HB3	2.51	0.45
1:A:75:ILE:HD13	1:A:98:PHE:CZ	2.51	0.45
1:D:26:GLY:HA3	1:D:47:ILE:HG23	1.99	0.45
1:D:51:LYS:O	1:D:55:LEU:HB2	2.17	0.45
1:B:208:THR:O	1:B:209:ALA:HB2	2.17	0.44
1:D:50:ARG:O	1:D:78:GLN:NE2	2.51	0.44
1:D:84:GLU:OE2	1:D:140:LYS:NZ	2.50	0.44
1:D:117:GLU:OE2	5:D:2066:HOH:O	2.21	0.44
1:C:286:ASP:OD2	1:C:288:SER:OG	2.34	0.44
1:A:205:TYR:HB2	1:A:233:ILE:HD12	2.00	0.44
1:A:280:TRP:CD2	1:A:281:PRO:HD2	2.53	0.44
1:C:165:PHE:CD1	1:C:171:SER:OG	2.65	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:57:SER:O	1:D:61:GLU:HB2	2.18	0.43
1:A:158:ILE:O	1:A:201:PRO:HD2	2.18	0.43
1:B:270:HIS:HE1	1:D:272:TYR:O	2.02	0.43
1:D:280:TRP:CG	1:D:281:PRO:HD2	2.53	0.43
1:B:280:TRP:CG	1:B:281:PRO:HD2	2.54	0.42
1:D:24:THR:O	1:D:107:ASN:HB3	2.18	0.42
1:B:28:THR:HB	1:B:209:ALA:HA	2.00	0.42
1:C:158:ILE:HB	1:C:175:ARG:HG3	2.00	0.42
1:C:35:VAL:HG13	1:C:45:VAL:HG11	2.01	0.42
1:D:179:TYR:CZ	1:D:183:LYS:HE2	2.54	0.42
1:C:24:THR:O	1:C:107:ASN:HB3	2.20	0.42
1:C:127:VAL:HG21	1:C:170:HIS:CD2	2.54	0.42
1:A:24:THR:O	1:A:107:ASN:HB3	2.19	0.42
1:B:165:PHE:CD1	1:B:168:ALA:HB3	2.55	0.42
1:A:95:LEU:O	1:A:99:GLY:HA3	2.21	0.41
1:C:208:THR:CG2	5:C:373:HOH:O	2.68	0.41
1:B:159:VAL:O	1:B:161:THR:HG23	2.19	0.41
1:C:158:ILE:O	1:C:201:PRO:HD2	2.19	0.41
1:D:35:VAL:CG1	1:D:62:LEU:HD13	2.51	0.41
1:D:83:ASN:OD1	1:D:86:GLU:HG3	2.21	0.41
1:A:205:TYR:HB2	1:A:233:ILE:CD1	2.51	0.41
1:C:165:PHE:O	1:C:166:PRO:C	2.57	0.41
1:D:51:LYS:HA	1:D:52:LEU:CB	2.49	0.40
1:C:87:VAL:HG13	1:C:138:MET:SD	2.61	0.40
1:C:179:TYR:CE1	1:C:258:SER:HB2	2.57	0.40
1:D:52:LEU:H	1:D:53:GLU:CG	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	296/303 (98%)	284 (96%)	11 (4%)	1 (0%)	41 31
1	B	281/303 (93%)	271 (96%)	10 (4%)	0	100 100
1	C	254/303 (84%)	240 (94%)	11 (4%)	3 (1%)	13 4
1	D	288/303 (95%)	276 (96%)	10 (4%)	2 (1%)	22 12
All	All	1119/1212 (92%)	1071 (96%)	42 (4%)	6 (0%)	29 18

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	56	LYS
1	D	53	GLU
1	C	55	LEU
1	A	111	GLN
1	C	158	ILE
1	D	158	ILE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	215/245 (88%)	207 (96%)	8 (4%)	34 25
1	B	221/245 (90%)	217 (98%)	4 (2%)	59 55
1	C	191/245 (78%)	184 (96%)	7 (4%)	34 25
1	D	219/245 (89%)	210 (96%)	9 (4%)	30 21
All	All	846/980 (86%)	818 (97%)	28 (3%)	38 29

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	THR
1	A	74	VAL
1	A	121	SER
1	A	165	PHE
1	A	170	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	215	SER
1	A	253	PHE
1	A	288	SER
1	B	92	LYS
1	B	121	SER
1	B	165	PHE
1	B	170	HIS
1	C	15	LEU
1	C	75	ILE
1	C	93	SER
1	C	165	PHE
1	C	170	HIS
1	C	175	ARG
1	C	288	SER
1	D	50	ARG
1	D	52	LEU
1	D	53	GLU
1	D	55	LEU
1	D	57	SER
1	D	74	VAL
1	D	113	LEU
1	D	165	PHE
1	D	170	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	131	ASN
1	B	63	GLN
1	B	111	GLN
1	B	180	ASN
1	B	270	HIS
1	D	80	ASN
1	D	118	HIS
1	D	150	HIS
1	D	277	HIS

5.3.3 RNA ①

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	4001	-	4,4,4	0.33	0	6,6,6	0.44	0
2	SO4	A	304	-	4,4,4	0.27	0	6,6,6	0.39	0
2	SO4	D	2001	-	4,4,4	0.20	0	6,6,6	0.52	0
2	SO4	D	1001	-	4,4,4	0.24	0	6,6,6	0.58	0
4	ADE	B	4002	-	9,11,11	1.74	2 (22%)	7,15,15	2.71	3 (42%)
3	PO4	B	3001	-	4,4,4	0.68	0	6,6,6	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADE	B	4002	-	-	-	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	4002	ADE	C2-N3	4.19	1.38	1.32
4	B	4002	ADE	C4-N9	2.12	1.38	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	4002	ADE	N3-C2-N1	-5.80	119.62	128.68
4	B	4002	ADE	C2-N3-C4	3.24	121.03	113.45
4	B	4002	ADE	C4-C5-N7	-2.11	107.20	109.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	111:GLN	C	112:PHE	N	1.14
1	D	194:ILE	C	195:ARG	N	0.99

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/303 (98%)	0.49	16 (5%) 25 29	22, 34, 53, 59	1 (0%)
1	B	283/303 (93%)	0.37	9 (3%) 47 50	20, 33, 48, 53	0
1	C	264/303 (87%)	0.86	38 (14%) 2 2	24, 38, 56, 60	0
1	D	291/303 (96%)	0.63	32 (10%) 5 6	25, 37, 55, 60	1 (0%)
All	All	1135/1212 (93%)	0.58	95 (8%) 11 12	20, 35, 53, 60	2 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	52	LEU	6.8
1	C	296	THR	6.8
1	A	52	LEU	6.1
1	D	57	SER	6.0
1	D	50	ARG	5.6
1	C	72	ALA	5.3
1	A	68	PRO	5.3
1	C	210	VAL	5.2
1	A	303	LEU	4.9
1	D	112	PHE	4.7
1	C	59	ALA	4.6
1	C	52	LEU	4.5
1	C	70	LYS	4.3
1	C	297	PHE	4.3
1	A	66	LEU	4.2
1	C	60	ASP	4.1
1	C	221	PHE	4.1
1	C	208	THR	3.9
1	D	66	LEU	3.7
1	D	77	ILE	3.6
1	D	60	ASP	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	55	LEU	3.6
1	C	165	PHE	3.5
1	A	59	ALA	3.5
1	D	108	GLY	3.4
1	A	65	ASN	3.4
1	D	267	LEU	3.4
1	D	83	ASN	3.3
1	A	7	GLY	3.2
1	D	64	ALA	3.2
1	C	225	PHE	3.2
1	D	54	ARG	3.1
1	B	259	VAL	3.1
1	B	179	TYR	3.1
1	A	179	TYR	3.0
1	C	77	ILE	3.0
1	D	111	GLN	2.9
1	C	293	MET	2.9
1	B	96	ASP	2.9
1	A	284	ALA	2.9
1	D	59	ALA	2.9
1	D	75	ILE	2.8
1	D	196	ILE	2.8
1	D	303	LEU	2.8
1	C	125	HIS	2.8
1	C	159	VAL	2.7
1	D	91	VAL	2.7
1	D	259	VAL	2.7
1	C	53	GLU	2.7
1	B	176	ALA	2.7
1	A	210	VAL	2.6
1	D	58	ALA	2.6
1	C	26	GLY	2.6
1	C	71	GLN	2.6
1	D	98	PHE	2.6
1	C	86	GLU	2.6
1	C	96	ASP	2.6
1	D	72	ALA	2.5
1	C	51	LYS	2.5
1	C	56	LYS	2.5
1	D	73	ARG	2.5
1	C	259	VAL	2.5
1	C	166	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	111	GLN	2.4
1	C	66	LEU	2.4
1	D	163	ALA	2.4
1	C	55	LEU	2.4
1	B	125[A]	HIS	2.4
1	B	297	PHE	2.4
1	B	208	THR	2.3
1	C	28	THR	2.3
1	D	79	CYS	2.3
1	D	78	GLN	2.3
1	A	259	VAL	2.3
1	C	62	LEU	2.3
1	C	81	ILE	2.2
1	A	63	GLN	2.2
1	C	88	ASN	2.2
1	D	65	ASN	2.2
1	C	205	TYR	2.2
1	A	164	GLY	2.2
1	A	199	VAL	2.2
1	C	126	ALA	2.2
1	C	75	ILE	2.2
1	C	124	TRP	2.2
1	C	97	THR	2.1
1	D	146	TRP	2.1
1	A	176	ALA	2.1
1	D	90	LEU	2.1
1	B	160	PRO	2.1
1	D	99	GLY	2.1
1	D	136	PHE	2.1
1	C	6	LYS	2.0
1	C	164	GLY	2.0
1	A	254	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	3001	5/5	0.85	0.28	56,56,57,57	0
4	ADE	B	4002	10/10	0.85	0.18	47,48,49,49	0
2	SO4	A	304	5/5	0.95	0.15	53,56,56,56	0
2	SO4	D	1001	5/5	0.95	0.09	50,53,57,59	0
2	SO4	D	2001	5/5	0.96	0.12	60,60,64,64	0
2	SO4	B	4001	5/5	0.98	0.08	33,35,38,39	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.