



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 25, 2024 – 04:14 PM JST

PDB ID : 8Z1R  
Title : isocitrate lyase MoMcl1  
Authors : Liu, X.H.; Zhao, W.H.  
Deposited on : 2024-04-11  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	
1	B	558	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	508	3981	2494	711	758	18	0	1	0
1	B	516	4036	2528	723	766	19	0	1	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	HIS	-	expression tag	UNP G5EH03
A	51	HIS	-	expression tag	UNP G5EH03
A	52	HIS	-	expression tag	UNP G5EH03
A	53	HIS	-	expression tag	UNP G5EH03
A	54	HIS	-	expression tag	UNP G5EH03
A	55	HIS	-	expression tag	UNP G5EH03
A	56	SER	-	expression tag	UNP G5EH03
A	57	SER	-	expression tag	UNP G5EH03
A	58	GLY	-	expression tag	UNP G5EH03
A	59	LEU	-	expression tag	UNP G5EH03
A	60	VAL	-	expression tag	UNP G5EH03
A	61	PRO	-	expression tag	UNP G5EH03
A	62	ARG	-	expression tag	UNP G5EH03
A	63	GLY	-	expression tag	UNP G5EH03
A	64	SER	-	expression tag	UNP G5EH03
B	50	HIS	-	expression tag	UNP G5EH03
B	51	HIS	-	expression tag	UNP G5EH03
B	52	HIS	-	expression tag	UNP G5EH03
B	53	HIS	-	expression tag	UNP G5EH03
B	54	HIS	-	expression tag	UNP G5EH03
B	55	HIS	-	expression tag	UNP G5EH03
B	56	SER	-	expression tag	UNP G5EH03
B	57	SER	-	expression tag	UNP G5EH03
B	58	GLY	-	expression tag	UNP G5EH03
B	59	LEU	-	expression tag	UNP G5EH03

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	60	VAL	-	expression tag	UNP G5EH03
B	61	PRO	-	expression tag	UNP G5EH03
B	62	ARG	-	expression tag	UNP G5EH03
B	63	GLY	-	expression tag	UNP G5EH03
B	64	SER	-	expression tag	UNP G5EH03

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	321	Total 321	O 321	0	0
4	B	414	Total 414	O 414	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isocitrate lyase



- Molecule 1: Isocitrate lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.43Å 150.50Å 128.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.81 – 1.82 32.81 – 1.82	Depositor EDS
% Data completeness (in resolution range)	100.0 (32.81-1.82) 100.0 (32.81-1.82)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 1.82Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.184 , 0.215 0.183 , 0.215	Depositor DCC
$R_{free}$ test set	5021 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.5	Xtrriage
Anisotropy	0.233	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8766	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/4073	0.55	0/5530
1	B	0.37	0/4130	0.58	0/5606
All	All	0.35	0/8203	0.56	0/11136

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3981	0	3850	36	0
1	B	4036	0	3911	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	4	0	6	0	0
3	B	8	0	12	0	0
4	A	321	0	0	6	0
4	B	414	0	0	3	0
All	All	8766	0	7779	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:THR:HG22	1:B:176:THR:H	1.50	0.77
1:B:193[B]:ASN:ND2	4:B:805:HOH:O	2.29	0.64
1:A:174:THR:HG22	1:A:176:THR:H	1.63	0.63
1:A:185:ASP:OD1	4:A:801:HOH:O	2.17	0.60
1:A:204:MET:HE2	1:A:207:ARG:HH22	1.68	0.59
1:A:351:GLU:OE1	1:A:434:ARG:NH2	2.29	0.59
1:A:382:SER:HA	1:A:385:LYS:HE2	1.86	0.57
1:B:94:ARG:HB2	1:B:125:ARG:HH12	1.70	0.57
1:A:410:ARG:NH2	4:A:808:HOH:O	2.40	0.55
1:A:564:ILE:O	1:B:543:ARG:NH1	2.37	0.53
1:A:211:ASP:OD2	1:A:582:ARG:NH1	2.41	0.53
1:A:543:ARG:NH1	1:B:564:ILE:O	2.41	0.51
1:A:417:LEU:O	1:A:421:LYS:HD2	2.09	0.51
1:B:347:LEU:HD22	1:B:360:ILE:HG23	1.92	0.51
1:A:215:LYS:HD3	1:A:582:ARG:NE	2.26	0.51
1:B:76:ASP:O	1:B:80:GLU:HG2	2.11	0.51
1:B:287:GLU:OE1	4:B:801:HOH:O	2.19	0.50
1:B:265:GLU:HB2	4:B:804:HOH:O	2.12	0.50
1:A:343:LEU:O	1:A:347:LEU:HG	2.13	0.49
1:A:244:LEU:O	1:A:248:THR:HG23	2.13	0.49
1:A:265:GLU:HB2	4:A:803:HOH:O	2.13	0.48
1:A:333:LEU:HB3	1:A:373:LEU:HB3	1.96	0.47
1:B:277:LEU:HD13	1:B:439:HIS:HE1	1.79	0.47
1:B:209:HIS:CE1	1:B:213:ARG:HD2	2.51	0.45
1:A:204:MET:HE2	1:A:207:ARG:NH2	2.29	0.45
1:A:215:LYS:HG2	1:A:582:ARG:HH21	1.81	0.45
1:B:81:GLN:HG2	1:B:84:ARG:HH12	1.83	0.44
1:A:402:ARG:HE	1:A:445:ARG:NH1	2.14	0.44
1:A:498:ASN:HB3	1:A:502:HIS:CE1	2.53	0.44
1:A:313:ASP:OD1	4:A:803:HOH:O	2.21	0.43
1:A:555:ASN:OD1	1:A:559:ARG:NH1	2.52	0.43
1:A:330:GLU:OE2	1:A:402:ARG:HG3	2.19	0.43
1:A:128:TRP:O	1:A:132:GLN:HG2	2.19	0.43
1:B:353:ARG:HA	1:B:353:ARG:HD3	1.78	0.42
1:A:318:ARG:NH1	4:A:821:HOH:O	2.51	0.42
1:B:311:ARG:HA	1:B:462:TRP:O	2.18	0.42
1:A:167:TRP:HB3	1:A:528:LEU:HD21	2.01	0.41
1:A:213:ARG:HA	1:A:216:MET:HG3	2.03	0.41
1:A:176:THR:HG23	1:A:187:PRO:HG2	2.03	0.41
1:A:393:GLU:H	1:A:393:GLU:HG2	1.49	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:MET:HE3	1:A:204:MET:HB2	1.95	0.41
1:A:311:ARG:HA	1:A:462:TRP:O	2.21	0.41
1:A:543:ARG:HG2	1:B:158:HIS:CE1	2.56	0.41
1:A:309:ILE:HG12	1:A:460:LEU:HB2	2.02	0.41
1:A:365:ALA:O	1:A:369:ARG:HG3	2.20	0.41
1:A:576:GLY:C	1:B:181:PRO:HG3	2.41	0.40
1:A:548:GLU:HB2	1:A:552:ALA:HB2	2.04	0.40
1:B:215:LYS:HD2	1:B:582:ARG:NH1	2.36	0.40
1:A:174:THR:HG21	4:A:869:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	505/558 (90%)	496 (98%)	9 (2%)	0	100	100
1	B	515/558 (92%)	507 (98%)	8 (2%)	0	100	100
All	All	1020/1116 (91%)	1003 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/445 (92%)	405 (99%)	4 (1%)	76	70
1	B	414/445 (93%)	408 (99%)	6 (1%)	67	58
All	All	823/890 (92%)	813 (99%)	10 (1%)	71	64

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	LYS
1	A	223	SER
1	A	428	TRP
1	A	482	GLU
1	B	81	GLN
1	B	115	GLN
1	B	165	SER
1	B	400	LYS
1	B	428	TRP
1	B	445	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	388	ASN
1	B	558	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	B	701	-	3,3,3	0.46	0	2,2,2	0.36	0
3	EDO	B	702	-	3,3,3	0.47	0	2,2,2	0.50	0
3	EDO	A	702	-	3,3,3	0.54	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	701	-	-	0/1/1/1	-
3	EDO	B	702	-	-	0/1/1/1	-
3	EDO	A	702	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	508/558 (91%)	0.51	61 (12%) <b>4</b> <b>3</b>	15, 30, 80, 105	0
1	B	516/558 (92%)	0.34	40 (7%) <b>13</b> <b>10</b>	15, 25, 57, 83	0
All	All	1024/1116 (91%)	0.43	101 (9%) <b>7</b> <b>5</b>	15, 27, 69, 105	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	419	ALA	8.7
1	A	219	GLN	8.1
1	B	276	HIS	7.4
1	A	222	SER	6.2
1	B	73	ALA	6.2
1	A	218	ALA	6.0
1	B	419	ALA	5.7
1	B	74	ALA	5.3
1	A	360	ILE	5.2
1	B	360	ILE	5.1
1	A	385	LYS	5.1
1	B	275	GLY	5.0
1	A	502	HIS	4.9
1	B	77	ALA	4.9
1	A	421	LYS	4.8
1	B	357	ALA	4.7
1	B	219	GLN	4.6
1	A	420	ASP	4.6
1	A	388	ASN	4.6
1	A	365	ALA	4.5
1	A	418	GLY	4.4
1	B	356	SER	4.4
1	A	416	ILE	4.3
1	A	369	ARG	4.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	420	ASP	4.2
1	A	422	PRO	4.2
1	B	358	SER	4.2
1	B	76	ASP	4.1
1	A	358	SER	4.0
1	A	225	PRO	4.0
1	A	500	MET	3.9
1	B	223	SER	3.9
1	A	73	ALA	3.8
1	A	501	GLY	3.7
1	A	357	ALA	3.7
1	B	277	LEU	3.6
1	B	570	THR	3.3
1	A	394	TYR	3.3
1	A	381	VAL	3.2
1	A	74	ALA	3.2
1	A	361	ASP	3.2
1	B	217	SER	3.2
1	A	370	ASP	3.2
1	A	175	SER	3.1
1	A	341	ALA	3.1
1	B	585	GLY	3.1
1	B	582	ARG	3.0
1	B	362	ALA	3.0
1	B	369	ARG	3.0
1	A	363	TYR	3.0
1	B	222	SER	3.0
1	A	362	ALA	2.9
1	B	224	THR	2.9
1	B	271	GLY	2.9
1	A	423	ASP	2.9
1	B	234	ILE	2.9
1	A	414	LYS	2.8
1	A	392	ALA	2.8
1	B	563	GLU	2.8
1	A	368	THR	2.8
1	B	75	GLU	2.8
1	A	223	SER	2.8
1	B	353	ARG	2.8
1	A	393	GLU	2.8
1	A	397	GLY	2.7
1	A	387	HIS	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	361	ASP	2.7
1	A	359	GLU	2.7
1	A	226	TYR	2.6
1	B	272	LYS	2.6
1	B	274	CYS	2.6
1	A	415	ASP	2.6
1	B	80	GLU	2.6
1	B	359	GLU	2.5
1	B	573	LYS	2.5
1	A	364	GLU	2.5
1	B	587	ILE	2.5
1	A	562	LYS	2.4
1	B	78	LEU	2.4
1	B	577	ALA	2.4
1	A	406	ILE	2.4
1	A	372	PRO	2.3
1	A	411	ALA	2.3
1	A	413	ALA	2.3
1	A	174	THR	2.3
1	A	224	THR	2.3
1	A	356	SER	2.3
1	A	340	ALA	2.3
1	B	218	ALA	2.3
1	A	383	HIS	2.2
1	B	581	ASP	2.2
1	A	217	SER	2.2
1	A	215	LYS	2.2
1	A	569	LEU	2.1
1	A	570	THR	2.1
1	A	396	ALA	2.1
1	B	363	TYR	2.1
1	A	389	VAL	2.1
1	A	367	PHE	2.1
1	A	407	TRP	2.1
1	A	400	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	A	702	4/4	0.81	0.16	37,38,40,48	0
2	MG	A	701	1/1	0.89	0.33	39,39,39,39	0
3	EDO	B	701	4/4	0.89	0.32	37,40,42,55	0
3	EDO	B	702	4/4	0.89	0.12	30,34,35,44	0
2	MG	B	703	1/1	0.91	0.06	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around EDO A 702:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around MG A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.