



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 14, 2023 – 11:41 AM JST

PDB ID : 5Z5B  
Title : Crystal structure of Tk-PTP in the G95A mutant form  
Authors : Ku, B.; Yun, H.Y.; Kim, S.J.  
Deposited on : 2018-01-17  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 2% 88% 10% .
1	B	160	 2% 86% 9% . .
1	C	160	 6% 78% 14% . . 6%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein-tyrosine phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	156	1249	795	221	226	7	0	0	0
1	B	156	1255	797	221	230	7	0	0	0
1	C	150	1203	768	210	218	7	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	ALA	GLY	engineered mutation	UNP Q8X270
A	148	LYS	-	expression tag	UNP Q8X270
A	149	LEU	-	expression tag	UNP Q8X270
A	150	ALA	-	expression tag	UNP Q8X270
A	151	ALA	-	expression tag	UNP Q8X270
A	152	ALA	-	expression tag	UNP Q8X270
A	153	LEU	-	expression tag	UNP Q8X270
A	154	GLU	-	expression tag	UNP Q8X270
A	155	HIS	-	expression tag	UNP Q8X270
A	156	HIS	-	expression tag	UNP Q8X270
A	157	HIS	-	expression tag	UNP Q8X270
A	158	HIS	-	expression tag	UNP Q8X270
A	159	HIS	-	expression tag	UNP Q8X270
A	160	HIS	-	expression tag	UNP Q8X270
B	95	ALA	GLY	engineered mutation	UNP Q8X270
B	148	LYS	-	expression tag	UNP Q8X270
B	149	LEU	-	expression tag	UNP Q8X270
B	150	ALA	-	expression tag	UNP Q8X270
B	151	ALA	-	expression tag	UNP Q8X270
B	152	ALA	-	expression tag	UNP Q8X270
B	153	LEU	-	expression tag	UNP Q8X270
B	154	GLU	-	expression tag	UNP Q8X270
B	155	HIS	-	expression tag	UNP Q8X270

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	156	HIS	-	expression tag	UNP Q8X270
B	157	HIS	-	expression tag	UNP Q8X270
B	158	HIS	-	expression tag	UNP Q8X270
B	159	HIS	-	expression tag	UNP Q8X270
B	160	HIS	-	expression tag	UNP Q8X270
C	95	ALA	GLY	engineered mutation	UNP Q8X270
C	148	LYS	-	expression tag	UNP Q8X270
C	149	LEU	-	expression tag	UNP Q8X270
C	150	ALA	-	expression tag	UNP Q8X270
C	151	ALA	-	expression tag	UNP Q8X270
C	152	ALA	-	expression tag	UNP Q8X270
C	153	LEU	-	expression tag	UNP Q8X270
C	154	GLU	-	expression tag	UNP Q8X270
C	155	HIS	-	expression tag	UNP Q8X270
C	156	HIS	-	expression tag	UNP Q8X270
C	157	HIS	-	expression tag	UNP Q8X270
C	158	HIS	-	expression tag	UNP Q8X270
C	159	HIS	-	expression tag	UNP Q8X270
C	160	HIS	-	expression tag	UNP Q8X270

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	C	1	Total	C	O	0	0
			3	1	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Protein-tyrosine phosphatase



- Molecule 1: Protein-tyrosine phosphatase



- Molecule 1: Protein-tyrosine phosphatase



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	154.80Å 154.80Å 75.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	77.40 – 2.30 26.44 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.6 (77.40-2.30) 90.7 (26.44-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.51 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.219 , 0.265 0.226 , 0.264	Depositor DCC
$R_{free}$ test set	1374 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.5	Xtrriage
Anisotropy	0.074	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 42.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.88	0/1273	0.95	0/1718
1	B	0.78	0/1277	0.96	2/1721 (0.1%)
1	C	0.76	0/1226	0.93	1/1654 (0.1%)
All	All	0.81	0/3776	0.94	3/5093 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	139	VAL	CB-CA-C	-7.22	97.68	111.40
1	C	73	LEU	CA-CB-CG	5.48	127.91	115.30
1	B	125	ARG	NE-CZ-NH1	5.27	122.94	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1249	0	1261	6	1
1	B	1255	0	1265	10	1
1	C	1203	0	1218	15	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3	0	1	0	0
3	B	3	0	1	0	0
3	C	3	0	1	1	0
All	All	3718	0	3747	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:152:ALA:O	1:B:153:LEU:N	1.75	1.18
1:B:152:ALA:O	1:B:152:ALA:CA	2.08	1.01
1:B:117:ARG:HD2	1:C:63:ASP:OD2	1.95	0.65
1:A:8:ILE:HB	1:A:108:LEU:HD21	1.85	0.58
1:A:108:LEU:N	1:A:108:LEU:HD23	2.20	0.57

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:GLU:OE1	1:B:156:HIS:NE2[5_445]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	154/160 (96%)	147 (96%)	6 (4%)	1 (1%)	25 31
1	B	152/160 (95%)	144 (95%)	6 (4%)	2 (1%)	12 12
1	C	148/160 (92%)	134 (90%)	13 (9%)	1 (1%)	22 26

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	454/480 (95%)	425 (94%)	25 (6%)	4 (1%)	17 20

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	21	ARG
1	B	131	VAL
1	C	131	VAL
1	A	131	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	130/137 (95%)	124 (95%)	6 (5%)	27 38
1	B	132/137 (96%)	126 (96%)	6 (4%)	27 39
1	C	126/137 (92%)	119 (94%)	7 (6%)	21 29
All	All	388/411 (94%)	369 (95%)	19 (5%)	25 35

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	65	THR
1	C	137	MET
1	C	141	LYS
1	C	133	THR
1	B	65	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	71	GLN
1	B	134	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FMT	C	201	-	2,2,2	0.81	0	1,1,1	0.36	0
3	FMT	B	202	-	2,2,2	0.88	0	1,1,1	0.60	0
3	FMT	A	202	-	2,2,2	0.48	0	1,1,1	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	201	FMT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	156/160 (97%)	-0.24	3 (1%) 66 73	15, 28, 45, 76	0
1	B	156/160 (97%)	-0.11	3 (1%) 66 73	16, 32, 54, 75	0
1	C	150/160 (93%)	0.52	9 (6%) 21 28	33, 49, 70, 78	0
All	All	462/480 (96%)	0.05	15 (3%) 47 54	15, 35, 61, 78	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	149	LEU	5.1
1	A	155	HIS	4.6
1	C	84	ARG	3.5
1	C	73	LEU	2.9
1	B	91	ILE	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	201	1/1	0.94	0.06	42,42,42,42	0
3	FMT	C	201	3/3	0.94	0.13	31,31,35,37	0
2	CL	B	201	1/1	0.95	0.04	54,54,54,54	0
3	FMT	B	202	3/3	0.99	0.08	22,22,22,23	0
3	FMT	A	202	3/3	0.99	0.09	19,19,22,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.