



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 16, 2023 – 01:34 AM EDT

PDB ID : 1Z7D
Title : Ornithine aminotransferase PY00104 from Plasmodium Yoelii
Authors : Walker, J.R.; Alam, Z.; Amani, M.; Lew, J.; Wasney, G.; Boulanger, K.; Weigelt, J.; Sundstrom, M.; Arrowsmith, C.; Edwards, A.; Bochkarev, A.; Hui, R.; Vedadi, M.; Structural Genomics Consortium (SGC)
Deposited on : 2005-03-24
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	
1	B	433	
1	C	433	
1	D	433	
1	E	433	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	433	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into three segments: a green segment on the left labeled '68%', a yellow segment in the middle labeled '16%', and a grey segment on the right labeled '13%'. A small red square is at the beginning of the bar, and two small black dots are at the end of the bar. A '%' symbol is positioned above the start of the bar.</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 18172 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ornithine aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	378	2969	1898	497	555	19	0	0	0
1	B	377	2960	1893	495	553	19	0	0	0
1	C	369	2895	1851	483	542	19	0	0	0
1	D	375	2944	1885	491	549	19	0	0	0
1	E	374	2935	1879	489	548	19	0	0	0
1	F	375	2944	1885	491	549	19	0	0	0

There are 114 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	cloning artifact	UNP Q7RT90
A	-18	GLY	-	cloning artifact	UNP Q7RT90
A	-17	SER	-	cloning artifact	UNP Q7RT90
A	-16	SER	-	cloning artifact	UNP Q7RT90
A	-15	HIS	-	cloning artifact	UNP Q7RT90
A	-14	HIS	-	cloning artifact	UNP Q7RT90
A	-13	HIS	-	cloning artifact	UNP Q7RT90
A	-12	HIS	-	cloning artifact	UNP Q7RT90
A	-11	HIS	-	cloning artifact	UNP Q7RT90
A	-10	HIS	-	cloning artifact	UNP Q7RT90
A	-9	SER	-	cloning artifact	UNP Q7RT90
A	-8	SER	-	cloning artifact	UNP Q7RT90
A	-7	GLY	-	cloning artifact	UNP Q7RT90
A	-6	LEU	-	cloning artifact	UNP Q7RT90
A	-5	VAL	-	cloning artifact	UNP Q7RT90
A	-4	PRO	-	cloning artifact	UNP Q7RT90
A	-3	ARG	-	cloning artifact	UNP Q7RT90

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	UNP Q7RT90
A	-1	SER	-	cloning artifact	UNP Q7RT90
B	-19	MET	-	cloning artifact	UNP Q7RT90
B	-18	GLY	-	cloning artifact	UNP Q7RT90
B	-17	SER	-	cloning artifact	UNP Q7RT90
B	-16	SER	-	cloning artifact	UNP Q7RT90
B	-15	HIS	-	cloning artifact	UNP Q7RT90
B	-14	HIS	-	cloning artifact	UNP Q7RT90
B	-13	HIS	-	cloning artifact	UNP Q7RT90
B	-12	HIS	-	cloning artifact	UNP Q7RT90
B	-11	HIS	-	cloning artifact	UNP Q7RT90
B	-10	HIS	-	cloning artifact	UNP Q7RT90
B	-9	SER	-	cloning artifact	UNP Q7RT90
B	-8	SER	-	cloning artifact	UNP Q7RT90
B	-7	GLY	-	cloning artifact	UNP Q7RT90
B	-6	LEU	-	cloning artifact	UNP Q7RT90
B	-5	VAL	-	cloning artifact	UNP Q7RT90
B	-4	PRO	-	cloning artifact	UNP Q7RT90
B	-3	ARG	-	cloning artifact	UNP Q7RT90
B	-2	GLY	-	cloning artifact	UNP Q7RT90
B	-1	SER	-	cloning artifact	UNP Q7RT90
C	-19	MET	-	cloning artifact	UNP Q7RT90
C	-18	GLY	-	cloning artifact	UNP Q7RT90
C	-17	SER	-	cloning artifact	UNP Q7RT90
C	-16	SER	-	cloning artifact	UNP Q7RT90
C	-15	HIS	-	cloning artifact	UNP Q7RT90
C	-14	HIS	-	cloning artifact	UNP Q7RT90
C	-13	HIS	-	cloning artifact	UNP Q7RT90
C	-12	HIS	-	cloning artifact	UNP Q7RT90
C	-11	HIS	-	cloning artifact	UNP Q7RT90
C	-10	HIS	-	cloning artifact	UNP Q7RT90
C	-9	SER	-	cloning artifact	UNP Q7RT90
C	-8	SER	-	cloning artifact	UNP Q7RT90
C	-7	GLY	-	cloning artifact	UNP Q7RT90
C	-6	LEU	-	cloning artifact	UNP Q7RT90
C	-5	VAL	-	cloning artifact	UNP Q7RT90
C	-4	PRO	-	cloning artifact	UNP Q7RT90
C	-3	ARG	-	cloning artifact	UNP Q7RT90
C	-2	GLY	-	cloning artifact	UNP Q7RT90
C	-1	SER	-	cloning artifact	UNP Q7RT90
D	-19	MET	-	cloning artifact	UNP Q7RT90
D	-18	GLY	-	cloning artifact	UNP Q7RT90

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-17	SER	-	cloning artifact	UNP Q7RT90
D	-16	SER	-	cloning artifact	UNP Q7RT90
D	-15	HIS	-	cloning artifact	UNP Q7RT90
D	-14	HIS	-	cloning artifact	UNP Q7RT90
D	-13	HIS	-	cloning artifact	UNP Q7RT90
D	-12	HIS	-	cloning artifact	UNP Q7RT90
D	-11	HIS	-	cloning artifact	UNP Q7RT90
D	-10	HIS	-	cloning artifact	UNP Q7RT90
D	-9	SER	-	cloning artifact	UNP Q7RT90
D	-8	SER	-	cloning artifact	UNP Q7RT90
D	-7	GLY	-	cloning artifact	UNP Q7RT90
D	-6	LEU	-	cloning artifact	UNP Q7RT90
D	-5	VAL	-	cloning artifact	UNP Q7RT90
D	-4	PRO	-	cloning artifact	UNP Q7RT90
D	-3	ARG	-	cloning artifact	UNP Q7RT90
D	-2	GLY	-	cloning artifact	UNP Q7RT90
D	-1	SER	-	cloning artifact	UNP Q7RT90
E	-19	MET	-	cloning artifact	UNP Q7RT90
E	-18	GLY	-	cloning artifact	UNP Q7RT90
E	-17	SER	-	cloning artifact	UNP Q7RT90
E	-16	SER	-	cloning artifact	UNP Q7RT90
E	-15	HIS	-	cloning artifact	UNP Q7RT90
E	-14	HIS	-	cloning artifact	UNP Q7RT90
E	-13	HIS	-	cloning artifact	UNP Q7RT90
E	-12	HIS	-	cloning artifact	UNP Q7RT90
E	-11	HIS	-	cloning artifact	UNP Q7RT90
E	-10	HIS	-	cloning artifact	UNP Q7RT90
E	-9	SER	-	cloning artifact	UNP Q7RT90
E	-8	SER	-	cloning artifact	UNP Q7RT90
E	-7	GLY	-	cloning artifact	UNP Q7RT90
E	-6	LEU	-	cloning artifact	UNP Q7RT90
E	-5	VAL	-	cloning artifact	UNP Q7RT90
E	-4	PRO	-	cloning artifact	UNP Q7RT90
E	-3	ARG	-	cloning artifact	UNP Q7RT90
E	-2	GLY	-	cloning artifact	UNP Q7RT90
E	-1	SER	-	cloning artifact	UNP Q7RT90
F	-19	MET	-	cloning artifact	UNP Q7RT90
F	-18	GLY	-	cloning artifact	UNP Q7RT90
F	-17	SER	-	cloning artifact	UNP Q7RT90
F	-16	SER	-	cloning artifact	UNP Q7RT90
F	-15	HIS	-	cloning artifact	UNP Q7RT90
F	-14	HIS	-	cloning artifact	UNP Q7RT90

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	-13	HIS	-	cloning artifact	UNP Q7RT90
F	-12	HIS	-	cloning artifact	UNP Q7RT90
F	-11	HIS	-	cloning artifact	UNP Q7RT90
F	-10	HIS	-	cloning artifact	UNP Q7RT90
F	-9	SER	-	cloning artifact	UNP Q7RT90
F	-8	SER	-	cloning artifact	UNP Q7RT90
F	-7	GLY	-	cloning artifact	UNP Q7RT90
F	-6	LEU	-	cloning artifact	UNP Q7RT90
F	-5	VAL	-	cloning artifact	UNP Q7RT90
F	-4	PRO	-	cloning artifact	UNP Q7RT90
F	-3	ARG	-	cloning artifact	UNP Q7RT90
F	-2	GLY	-	cloning artifact	UNP Q7RT90
F	-1	SER	-	cloning artifact	UNP Q7RT90

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	127	Total O 127 127	0	0
2	B	87	Total O 87 87	0	0
2	C	56	Total O 56 56	0	0
2	D	82	Total O 82 82	0	0
2	E	67	Total O 67 67	0	0
2	F	106	Total O 106 106	0	0



• Molecule 1: ornithine aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.49Å 221.83Å 114.48Å 90.00° 100.89° 90.00°	Depositor
Resolution (Å)	49.74 – 2.10 49.74 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.1 (49.74-2.10) 99.1 (49.74-2.09)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.259 0.214 , 0.258	Depositor DCC
R_{free} test set	7970 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtrriage
Anisotropy	0.180	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18172	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.61	0/3023	0.69	0/4092
1	B	0.55	0/3014	0.68	2/4080 (0.0%)
1	C	0.50	0/2947	0.63	1/3990 (0.0%)
1	D	0.55	1/2998 (0.0%)	0.67	1/4058 (0.0%)
1	E	0.52	0/2989	0.65	0/4047
1	F	0.58	0/2998	0.70	1/4058 (0.0%)
All	All	0.55	1/17969 (0.0%)	0.67	5/24325 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	62	CYS	CB-SG	-5.17	1.73	1.81

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	77	LEU	CA-CB-CG	6.11	129.35	115.30
1	F	367	LEU	CA-CB-CG	5.86	128.77	115.30
1	B	309	LEU	CA-CB-CG	5.67	128.33	115.30
1	D	77	LEU	CA-CB-CG	5.21	127.27	115.30
1	C	263	LEU	CA-CB-CG	5.01	126.81	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2969	0	3015	56	0
1	B	2960	0	3007	57	0
1	C	2895	0	2939	64	0
1	D	2944	0	2995	61	0
1	E	2935	0	2982	40	0
1	F	2944	0	2995	58	0
2	A	127	0	0	3	0
2	B	87	0	0	2	0
2	C	56	0	0	4	0
2	D	82	0	0	1	0
2	E	67	0	0	2	0
2	F	106	0	0	4	0
All	All	18172	0	17933	305	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 305 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:366:LYS:HD2	2:F:514:HOH:O	1.49	1.10
1:C:374:THR:HG21	1:C:383:ARG:O	1.52	1.09
1:A:21:ALA:HB3	1:B:106:LEU:HD23	1.34	1.07
1:E:98:ASN:HB3	2:E:459:HOH:O	1.58	1.03
1:A:374:THR:HG21	1:A:383:ARG:O	1.60	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	374/433 (86%)	356 (95%)	14 (4%)	4 (1%)	14 9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	373/433 (86%)	353 (95%)	18 (5%)	2 (0%)	29	26
1	C	365/433 (84%)	350 (96%)	13 (4%)	2 (0%)	29	26
1	D	371/433 (86%)	352 (95%)	16 (4%)	3 (1%)	19	15
1	E	370/433 (86%)	350 (95%)	19 (5%)	1 (0%)	41	41
1	F	371/433 (86%)	348 (94%)	18 (5%)	5 (1%)	12	7
All	All	2224/2598 (86%)	2109 (95%)	98 (4%)	17 (1%)	19	15

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	290	SER
1	C	288	HIS
1	C	379	ASP
1	D	288	HIS
1	D	379	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	331/378 (88%)	306 (92%)	25 (8%)	13	10
1	B	330/378 (87%)	309 (94%)	21 (6%)	17	14
1	C	322/378 (85%)	302 (94%)	20 (6%)	18	15
1	D	328/378 (87%)	307 (94%)	21 (6%)	17	14
1	E	327/378 (86%)	304 (93%)	23 (7%)	15	12
1	F	328/378 (87%)	304 (93%)	24 (7%)	14	11
All	All	1966/2268 (87%)	1832 (93%)	134 (7%)	16	13

5 of 134 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	106	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	204	GLU
1	F	367	LEU
1	C	99	LEU
1	C	77	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	355	ASN
1	F	22	HIS
1	D	65	ASN
1	F	267	HIS
1	E	267	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/433 (87%)	0.13	4 (1%) 80 84	16, 34, 57, 71	0
1	B	377/433 (87%)	0.34	15 (3%) 38 44	20, 38, 63, 71	0
1	C	369/433 (85%)	0.36	24 (6%) 18 23	27, 43, 72, 81	0
1	D	375/433 (86%)	0.21	10 (2%) 54 60	23, 40, 57, 83	0
1	E	374/433 (86%)	0.50	25 (6%) 17 22	23, 43, 68, 81	0
1	F	375/433 (86%)	0.24	5 (1%) 77 80	20, 36, 58, 71	0
All	All	2248/2598 (86%)	0.30	83 (3%) 41 48	16, 39, 64, 83	0

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	288	HIS	10.3
1	D	287	GLU	5.7
1	B	289	GLY	4.8
1	D	289	GLY	4.8
1	C	135	ASN	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.