

Full wwPDB X-ray Structure Validation Report (i)

Apr 22, 2025 – 10:14 AM JST

PDB ID : 8Z8L / pdb 00008z8l

Title: structure of IL-4Ra in complex with an nanobody 4E9

Authors : Song, G.; Qiu, W.

Deposited on : 2024-04-22

Resolution : 3.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0rc1

EDS: 3.0

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.006 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.42

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	1115 (4.20-3.72)
Clashscore	180529	1009 (4.16-3.76)
Ramachandran outliers	177936	1006 (4.18-3.74)
Sidechain outliers	177891	1117 (4.20-3.72)
RSRZ outliers	164620	1116 (4.20-3.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	В	214	46%	40%	• 12%			
2	A	124	50%	42%	• 7%			
3	С	2	1	.00%				



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Interleukin-4 receptor subunit alpha.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	В	188	Total 1506	C 962	N 252	O 282	S 10	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	182	ALA	CYS	engineered mutation	UNP P24394
В	208	HIS	-	expression tag	UNP P24394
В	209	HIS	-	expression tag	UNP P24394
В	210	HIS	-	expression tag	UNP P24394
В	211	HIS	-	expression tag	UNP P24394
В	212	HIS	-	expression tag	UNP P24394
В	213	HIS	-	expression tag	UNP P24394
В	214	ALA	-	expression tag	UNP P24394

• Molecule 2 is a protein called nanobody dAb1.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	A	115	Total 832	C 516	N 141	O 169	S 6	0	0	0


• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
3	С	2	Total 28	C 16	N 2	O 10	0	0	0



 \bullet Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6).$




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total C N O 14 8 1 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Interleukin-4 receptor subunit alpha



Chain C: 100%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants	142.94Å 142.94Å 149.09Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.82 - 3.96	Depositor
Resolution (A)	19.82 - 3.96	EDS
% Data completeness	86.2 (19.82-3.96)	Depositor
(in resolution range)	85.3 (19.82-3.96)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.21 (at 3.94Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D	0.301 , 0.334	Depositor
R, R_{free}	0.300 , 0.335	DCC
R_{free} test set	263 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	191.4	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.30 , 147.4	EDS
L-test for twinning ²	$ < L > = 0.44, < L^2> = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2380	wwPDB-VP
Average B, all atoms (Å ²)	208.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	В	0.25	0/1553	0.53	0/2130	
2	A	0.28	0/847	0.52	0/1152	
All	All	0.26	0/2400	0.53	0/3282	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1506	0	1413	75	0
2	A	832	0	760	46	0
3	С	28	0	25	0	0
4	В	14	0	13	0	0
All	All	2380	0	2211	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
1:B:134:TYR:HB2	1:B:152:VAL:HB	<u>distance (Å)</u> 1.59	overlap (Å) 0.84
1:B:43:LEU:HB2	2:A:56:TYR:HB3		0.83
1:B:140:SER:HA	1:B:172:TYR:HA	1.60	0.83
		1.58	
1:B:41:PHE:CE1	2:A:49:ARG:CZ	2.64	0.81
2:A:29:TYR:HB3	2:A:97:VAL:HA	1.68	0.76
1:B:41:PHE:CD1	2:A:49:ARG:NH2	2.56	0.74
2:A:90:VAL:HG22	2:A:110:LEU:HB2	1.68	0.73
1:B:98:PRO:HG2	1:B:188:SER:HB3	1.70	0.73
1:B:41:PHE:CE1	2:A:49:ARG:NH1	2.58	0.71
2:A:6:GLU:HG3	2:A:22:CYS:HB2	1.72	0.70
2:A:11:LEU:HD11	2:A:114:SER:HB3	1.74	0.70
2:A:29:TYR:CD1	2:A:50:SER:HB2	2.28	0.69
2:A:90:VAL:HA	2:A:110:LEU:HA	1.76	0.68
1:B:35:LEU:HD12	1:B:49:CYS:HB3	1.77	0.67
1:B:54:ASN:HD22	1:B:60:VAL:HG23	1.59	0.66
1:B:141:GLU:HB2	1:B:173:ARG:HG3	1.78	0.66
1:B:99:ARG:HB3	1:B:121:PRO:HB2	1.79	0.65
1:B:22:LYS:HA	1:B:58:GLY:HA2	1.77	0.65
1:B:41:PHE:HE1	2:A:29:TYR:HH	1.45	0.64
1:B:107:HIS:HB3	1:B:115:LEU:HB3	1.79	0.64
1:B:41:PHE:CD1	2:A:49:ARG:CZ	2.80	0.64
1:B:41:PHE:HE1	2:A:29:TYR:OH	1.83	0.61
1:B:13:TYR:HD2	1:B:128:LEU:HD21	1.67	0.60
1:B:66:ASP:OD1	1:B:67:ASP:N	2.35	0.59
2:A:88:THR:HG23	2:A:112:THR:HA	1.84	0.59
2:A:96:GLY:HA3	2:A:101:CYS:HA	1.83	0.59
1:B:3:VAL:HA	1:B:23:MET:HA	1.86	0.58
1:B:80:ALA:HB2	1:B:85:LEU:HD11	1.85	0.58
1:B:34:ARG:HB2	1:B:79:TRP:HB2	1.85	0.57
1:B:26:PRO:HA	1:B:56:GLY:O	2.05	0.57
1:B:40:VAL:HG21	1:B:73:ASN:HB3	1.87	0.56
2:A:26:GLY:O	2:A:74:ASN:ND2	2.40	0.55
1:B:54:ASN:OD1	1:B:55:GLY:N	2.40	0.54
2:A:79:GLN:HE22	2:A:81:ASN:ND2	2.06	0.54
1:B:23:MET:SD	1:B:33:LEU:HD11	2.47	0.54
1:B:15:SER:HA	1:B:65:MET:O	2.08	0.53
1:B:131:HIS:CD2	1:B:180:ALA:HB1	2.44	0.53
2:A:33:TRP:NE1	2:A:76:LEU:HG	2.23	0.52
2:A:51:GLY:O	2:A:53:ARG:NH1	2.42	0.52
1:B:39:LEU:HD13	1:B:74:TYR:CZ	2.45	0.52
2:A:24:ALA:HB3	2:A:74:ASN:ND2	2.25	0.52
1:B:125:ASP:OD1	1:B:125:ASP:N	2.38	0.51
1.D.120.ASP.ODI	1:D:120:A5F:N	2.38	0.01

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from pred		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\rm \mathring{A})$	overlap (Å)
2:A:57:TYR:HB2	2:A:62:LYS:HG2	1.94	0.49
1:B:136:VAL:HG22	1:B:176:VAL:HG23	1.93	0.49
1:B:54:ASN:HB3	1:B:60:VAL:HG23	1.95	0.49
1:B:19:CYS:HB2	1:B:61:CYS:HB3	1.94	0.48
1:B:113:THR:HA	1:B:162:ALA:HB2	1.93	0.48
1:B:175:ARG:HD2	1:B:190:TRP:HB3	1.96	0.48
1:B:129:TYR:CE2	1:B:155:LEU:HD12	2.49	0.48
2:A:29:TYR:HD1	2:A:50:SER:HB2	1.77	0.48
2:A:36:GLN:OE1	2:A:42:ARG:N	2.47	0.48
1:B:39:LEU:HD13	1:B:74:TYR:CE1	2.48	0.48
2:A:45:VAL:HG13	2:A:61:VAL:HG11	1.96	0.48
1:B:98:PRO:HD2	1:B:186:THR:O	2.14	0.47
1:B:98:PRO:HG3	1:B:179:TRP:HA	1.96	0.47
1:B:134:TYR:CD1	1:B:178:ALA:HB2	2.49	0.47
2:A:4:LEU:HD11	2:A:95:VAL:HG22	1.96	0.46
1:B:139:TRP:CZ3	1:B:175:ARG:HG2	2.51	0.46
1:B:136:VAL:HG22	1:B:176:VAL:CG2	2.45	0.46
1:B:12:ASP:HA	1:B:97:LYS:O	2.15	0.46
1:B:139:TRP:HA	1:B:147:PHE:HA	1.98	0.46
1:B:39:LEU:HD11	1:B:72:ASP:HB3	1.97	0.46
2:A:13:GLN:HG3	2:A:14:PRO:HD2	1.98	0.46
1:B:39:LEU:HG	2:A:49:ARG:HH22	1.81	0.46
1:B:99:ARG:HB2	1:B:122:TYR:CZ	2.51	0.46
1:B:36:LEU:HD13	1:B:79:TRP:CD1	2.51	0.45
1:B:170:ILE:HG22	1:B:171:SER:H	1.81	0.45
1:B:104:LEU:HA	1:B:117:THR:O	2.16	0.45
2:A:36:GLN:O	2:A:89:ALA:HB1	2.16	0.45
2:A:90:VAL:HG22	2:A:110:LEU:CB	2.43	0.45
1:B:41:PHE:HE1	2:A:49:ARG:NH1	2.13	0.45
2:A:33:TRP:CE2	2:A:78:LEU:HB2	2.52	0.45
2:A:92:TYR:HD1	2:A:108:GLY:HA3	1.81	0.45
1:B:124:PRO:HA	1:B:129:TYR:CD2	2.52	0.44
1:B:134:TYR:HD1	1:B:178:ALA:HB2	1.82	0.44
2:A:47:ALA:HB3	2:A:56:TYR:HB2	1.98	0.44
1:B:39:LEU:HG	2:A:49:ARG:NH2	2.33	0.44
1:B:177:ARG:HB3	1:B:190:TRP:HA	1.99	0.44
2:A:58:ALA:O	2:A:62:LYS:HG3	2.18	0.44
1:B:21:TRP:CE2	1:B:59:CYS:HB2	2.53	0.44
2:A:100:ASN:O	2:A:102:ARG:N	2.51	0.44
1:B:36:LEU:HD13	1:B:79:TRP:HD1	1.83	0.44
1:B:52:GLU:OE2	1:B:62:HIS:NE2	2.45	0.44

Continued on next page...



Continued from previous page...

A + 1		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:50:ILE:HD12	1:B:51:PRO:HD2	1.99	0.43
1:B:17:SER:HB2	1:B:65:MET:HE2	2.00	0.43
2:A:35:ARG:HG3	2:A:89:ALA:HB3	2.00	0.43
2:A:35:ARG:HD3	2:A:91:TYR:CZ	2.54	0.43
1:B:29:CYS:HA	1:B:33:LEU:HG	2.01	0.43
2:A:65:PHE:CG	2:A:80:MET:HG2	2.53	0.43
2:A:96:GLY:HA2	2:A:103:ASN:CG	2.40	0.43
1:B:84:LEU:HD23	1:B:84:LEU:HA	1.86	0.42
1:B:175:ARG:HG3	1:B:190:TRP:CE3	2.55	0.42
1:B:41:PHE:HE1	2:A:29:TYR:CZ	2.37	0.42
2:A:4:LEU:HB2	2:A:106:GLY:HA3	2.00	0.42
1:B:4:LEU:N	1:B:22:LYS:O	2.53	0.41
1:B:141:GLU:HG3	1:B:195:LYS:NZ	2.35	0.41
2:A:90:VAL:HG13	2:A:109:THR:C	2.40	0.41
1:B:40:VAL:CG2	1:B:73:ASN:HB3	2.50	0.41
1:B:139:TRP:HZ3	1:B:175:ARG:HG2	1.85	0.41
2:A:1:GLN:HG3	2:A:104:TYR:HE1	1.86	0.41
1:B:37:TYR:HA	1:B:75:THR:O	2.21	0.41
2:A:30:CYS:N	2:A:96:GLY:O	2.41	0.41
1:B:76:LEU:HB3	1:B:86:TRP:HZ3	1.85	0.41
1:B:175:ARG:HD2	1:B:190:TRP:CB	2.51	0.41
1:B:80:ALA:O	1:B:82:GLN:N	2.54	0.41
1:B:3:VAL:HG13	1:B:21:TRP:HD1	1.86	0.40
1:B:13:TYR:HB2	1:B:96:VAL:HG13	2.02	0.40
1:B:132:LEU:HD22	1:B:178:ALA:HB1	2.04	0.40
1:B:171:SER:HA	1:B:197:HIS:HA	2.03	0.40
2:A:24:ALA:HB3	2:A:74:ASN:HD21	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percei	ntiles
1	В	182/214~(85%)	165 (91%)	16 (9%)	1 (0%)	25	60
2	A	113/124 (91%)	100 (88%)	13 (12%)	0	100	100
All	All	295/338~(87%)	265 (90%)	29 (10%)	1 (0%)	37	70

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	81	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	d Rotameric Outliers		Percentiles		
1	В	165/192~(86%)	162 (98%)	3 (2%)	54 71		
2	A	83/98 (85%)	81 (98%)	2 (2%)	44 62		
All	All	248/290 (86%)	243 (98%)	5 (2%)	50 68		

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	13	TYR
1	В	29	CYS
1	В	175	ARG
2	A	93	CYS
2	A	101	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	79	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res Link		Bo	ond leng	$ ag{ths}$	В	ond ang	les
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	С	1	1,3	14,14,15	0.25	0	17,19,21	0.46	0
3	NAG	С	2	3	14,14,15	0.26	0	17,19,21	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	С	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	С	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	С	1	NAG	C4-C5-C6-O6
3	С	1	NAG	O5-C5-C6-O6



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
IVIOI	туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	В	301	1	14,14,15	0.34	0	17,19,21	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	В	301	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	В	301	NAG	C8-C7-N2-C2
4	В	301	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	В	188/214 (87%)	-0.57	2 (1%) 77 62	178, 218, 264, 281	0
2	A	115/124~(92%)	-0.57	0 100 100	156, 183, 212, 253	0
All	All	303/338 (89%)	-0.57	2 (0%) 84 70	156, 203, 261, 281	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	128	LEU	2.6
1	В	143	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	NAG	С	2	14/15	0.72	0.08	214,214,214,214	0
3	NAG	С	1	14/15	0.90	0.07	213,213,213,213	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	NAG	В	301	14/15	0.66	0.09	206,206,206,206	0

6.5 Other polymers (i)

There are no such residues in this entry.

