



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2026 – 12:04 PM EDT

PDB ID : 9ZCL / pdb_00009zcl
Title : N17.3.2 NRASQ61K HLA A1 complex
Authors : Sharma, V.K.; Gallagher, D.T.; Mariuzza, R.A.
Deposited on : 2025-11-23
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	10	<div> <div>12%</div> <div>90% 10%</div> </div>
2	C	276	<div> <div>12%</div> <div>80% 7% 12%</div> </div>
3	D	100	<div> <div>93% 7%</div> </div>
4	A	212	<div> <div>6%</div> <div>85% 11%</div> </div>
5	B	241	<div> <div>%</div> <div>87% 11%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6386 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ILE-LEU-ASP-THR-ALA-GLY-LYS-GLU-GLU-TYR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	P	10	Total	C	N	O	0	0	0
			80	50	11	19			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	7	LYS	GLN	conflict	UNP P01116

- Molecule 2 is a protein called HLA class I histocompatibility antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	243	Total	C	N	O	S	0	0	0
			1837	1140	345	343	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	MET	-	initiating methionine	UNP Q59GJ2

- Molecule 3 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	100	Total	C	N	O	S	0	0	0
			808	514	135	155	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP P61769


- Molecule 4 is a protein called N17.3.2 TCR alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	204	Total	C	N	O	S	0	0	0
			1506	951	242	306	7			

- Molecule 5 is a protein called N17.3.2 TCR beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	B	236	Total	C	N	O	S	1	0	0
			1813	1145	309	350	9			

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	O	S	0	0
			5	4	1		
6	C	1	Total	O	S	0	0
			5	4	1		
6	D	1	Total	O	S	0	0
			5	4	1		
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	P	8	Total 8	O 8	0	0
7	C	109	Total 109	O 109	0	0
7	D	60	Total 60	O 60	0	0
7	A	54	Total 54	O 54	0	0
7	B	86	Total 86	O 86	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ILE-LEU-ASP-THR-ALA-GLY-LYS-GLU-GLU-TYR

Chain P: 



- Molecule 2: HLA class I histocompatibility antigen

Chain C: 





- Molecule 3: Beta-2-microglobulin

Chain D: 





- Molecule 4: N17.3.2 TCR alpha

Chain A: 



- Molecule 5: N17.3.2 TCR beta

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	160.39Å 160.39Å 89.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.83 – 2.30 29.83 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.83-2.30) 99.9 (29.83-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.31Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.206 , 0.256 0.208 , 0.256	Depositor DCC
R_{free} test set	2638 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6386	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	P	0.39	0/80	0.52	0/105
2	C	0.48	0/1884	0.70	1/2543 (0.0%)
3	D	0.36	0/831	0.55	0/1132
4	A	0.38	0/1539	0.61	0/2089
5	B	0.33	0/1862	0.51	0/2539
All	All	0.40	0/6196	0.60	1/8408 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
5	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	208	PHE	CA-CB-CG	6.81	120.61	113.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
5	B	106	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	80	0	77	1	0
2	C	1837	0	1613	12	0
3	D	808	0	738	3	0
4	A	1506	0	1330	17	0
5	B	1813	0	1661	16	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
6	C	10	0	0	0	0
6	D	5	0	0	0	0
7	A	54	0	0	1	0
7	B	86	0	0	3	0
7	C	109	0	0	1	0
7	D	60	0	0	1	0
7	P	8	0	0	0	0
All	All	6386	0	5419	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:183:ASP:HB2	2:C:209:TYR:HB2	1.68	0.74
2:C:209:TYR:HB3	2:C:210:PRO:HD3	1.75	0.68
2:C:259:CYS:N	2:C:272:LEU:HD12	2.12	0.64
2:C:210:PRO:HD2	2:C:263:HIS:CE1	2.33	0.62
1:P:2:LEU:O	2:C:163:ARG:NH2	2.36	0.59
4:A:120:GLN:N	4:A:120:GLN:OE1	2.36	0.59
5:B:196:PHE:O	5:B:202:ASN:ND2	2.35	0.56
4:A:36:TRP:HB2	4:A:49:ILE:HG22	1.90	0.54
3:D:91:LYS:NZ	7:D:202:HOH:O	2.40	0.53
2:C:192:HIS:O	2:C:200:THR:N	2.41	0.53
5:B:205:ARG:NH1	5:B:207:GLN:HB2	2.26	0.51
5:B:192:VAL:HG11	5:B:196:PHE:HD1	1.75	0.51
2:C:210:PRO:HD2	2:C:263:HIS:HE1	1.74	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:116:LYS:NZ	7:A:404:HOH:O	2.44	0.49
4:A:44:GLU:HB2	7:B:402:HOH:O	2.12	0.49
4:A:203:THR:HG23	4:A:205:PHE:CE1	2.48	0.49
3:D:36:GLU:O	3:D:82:VAL:HA	2.12	0.49
5:B:207:GLN:HG3	5:B:230:ILE:HG23	1.96	0.48
5:B:59:GLU:HG2	5:B:60:VAL:HG13	1.95	0.48
4:A:172:MET:HE3	4:A:172:MET:HA	1.96	0.47
5:B:131:ILE:HG23	5:B:194:ALA:HB1	1.96	0.47
4:A:203:THR:HG23	4:A:205:PHE:HE1	1.79	0.47
5:B:126:PRO:HD2	5:B:197:TRP:CZ2	2.50	0.47
5:B:12:VAL:HG22	5:B:148:PRO:HG3	1.95	0.46
5:B:70:ASN:HB2	7:B:445:HOH:O	2.15	0.46
4:A:51:ARG:NH2	4:A:59:GLU:HB2	2.30	0.45
5:B:31:SER:HA	5:B:49:SER:O	2.17	0.45
2:C:174:ASN:HD22	2:C:174:ASN:HA	1.64	0.45
3:D:23:LEU:O	3:D:67:TYR:HA	2.18	0.44
2:C:170:ARG:HG3	7:C:506:HOH:O	2.17	0.44
4:A:120:GLN:CD	4:A:121:ASN:H	2.25	0.44
4:A:160:TYR:O	4:A:181:ALA:HA	2.19	0.43
2:C:36:PHE:CG	2:C:67:MET:HG2	2.54	0.42
5:B:188:SER:C	5:B:189:ARG:HD3	2.44	0.42
4:A:42:SER:OG	4:A:44:GLU:OE1	2.32	0.42
5:B:79:GLU:N	7:B:408:HOH:O	2.53	0.42
5:B:147:TYR:CD1	5:B:148:PRO:HA	2.54	0.42
4:A:63:ARG:CZ	4:A:83:GLN:HG3	2.49	0.42
4:A:152:SER:H	4:A:197:SER:HB3	1.85	0.42
5:B:126:PRO:HG3	5:B:139:LEU:HD12	2.01	0.41
2:C:32:GLN:NE2	2:C:35:ARG:HB3	2.36	0.41
4:A:51:ARG:HD3	4:A:55:ASP:HB2	2.03	0.41
4:A:165:CYS:HB3	5:B:189:ARG:NH2	2.35	0.41
4:A:64:TYR:HB3	4:A:66:TRP:CH2	2.56	0.40
2:C:21:ARG:HD2	2:C:39:ASP:CG	2.47	0.40
4:A:160:TYR:CE1	5:B:175:GLU:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
2	C	235/276 (85%)	223 (95%)	12 (5%)	0	100	100
3	D	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
4	A	202/212 (95%)	191 (95%)	11 (5%)	0	100	100
5	B	234/241 (97%)	227 (97%)	7 (3%)	0	100	100
All	All	777/839 (93%)	745 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	8/8 (100%)	8 (100%)	0	100	100
2	C	171/233 (73%)	165 (96%)	6 (4%)	32	48
3	D	88/95 (93%)	86 (98%)	2 (2%)	44	63
4	A	157/192 (82%)	155 (99%)	2 (1%)	61	77
5	B	188/208 (90%)	185 (98%)	3 (2%)	55	73
All	All	612/736 (83%)	599 (98%)	13 (2%)	47	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	178	THR
2	C	182	THR
2	C	208	PHE
2	C	243	LYS
2	C	260	HIS
2	C	272	LEU
3	D	31	HIS
3	D	70	PHE
4	A	60	ILE
4	A	73	SER
5	B	4	VAL
5	B	43	LEU
5	B	127	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	32	GLN
2	C	141	GLN
2	C	174	ASN
3	D	8	GLN
3	D	13	HIS
4	A	184	ASN
5	B	28	ASN
5	B	171	GLN
5	B	229	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	C	301	-	4,4,4	0.34	0	6,6,6	0.10	0
6	SO4	C	302	-	4,4,4	0.14	0	6,6,6	0.27	0
6	SO4	A	301	-	4,4,4	0.28	0	6,6,6	0.21	0
6	SO4	D	101	-	4,4,4	0.23	0	6,6,6	0.20	0
6	SO4	B	301	-	4,4,4	0.25	0	6,6,6	0.28	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	10/10 (100%)	-0.71	0 100 100	22, 23, 25, 32	0
2	C	243/276 (88%)	0.40	34 (13%) 6 7	19, 34, 77, 93	0
3	D	100/100 (100%)	-0.32	0 100 100	20, 30, 45, 54	0
4	A	204/212 (96%)	0.45	12 (5%) 28 30	25, 47, 67, 85	0
5	B	236/241 (97%)	0.04	3 (1%) 75 76	22, 36, 57, 69	1 (0%)
All	All	793/839 (94%)	0.20	49 (6%) 26 28	19, 37, 70, 93	1 (0%)

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	272	LEU	6.3
2	C	259	CYS	5.3
2	C	209	TYR	4.9
2	C	215	LEU	4.7
4	A	136	ASP	4.1
4	A	135	SER	3.9
2	C	247	VAL	3.8
2	C	270	LEU	3.6
2	C	203	CYS	3.6
2	C	266	LEU	3.6
2	C	204	TRP	3.6
2	C	180	GLN	3.5
2	C	269	PRO	3.5
2	C	185	PRO	3.4
2	C	182	THR	3.1
2	C	261	VAL	2.9
2	C	214	THR	2.9
2	C	210	PRO	2.9
2	C	264	GLU	2.8
4	A	137	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	206	LEU	2.8
2	C	267	PRO	2.7
2	C	207	GLY	2.7
2	C	230	LEU	2.7
2	C	260	HIS	2.6
2	C	246	ALA	2.6
5	B	239	ALA	2.6
4	A	205	PHE	2.6
2	C	187	THR	2.5
4	A	191	ALA	2.4
2	C	208	PHE	2.4
2	C	179	LEU	2.4
4	A	185	LYS	2.4
2	C	265	GLY	2.4
2	C	244	TRP	2.4
4	A	192	ASN	2.3
5	B	27	MET	2.3
4	A	199	ILE	2.3
4	A	133	LYS	2.3
2	C	186	LYS	2.3
4	A	196	ASN	2.2
2	C	90	ASP	2.2
2	C	184	PRO	2.2
2	C	176	LYS	2.1
4	A	193	ALA	2.1
4	A	22	ASP	2.1
2	C	178	THR	2.0
2	C	228	THR	2.0
5	B	194	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	SO4	B	301	5/5	0.91	0.14	54,58,64,67	0
6	SO4	D	101	5/5	0.96	0.10	46,51,53,55	0
6	SO4	A	301	5/5	0.97	0.09	36,43,50,57	0
6	SO4	C	301	5/5	0.97	0.12	32,36,45,49	0
6	SO4	C	302	5/5	0.98	0.07	35,37,40,42	0

6.5 Other polymers

There are no such residues in this entry.