



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 06:44 am BST

PDB ID : 4ZDI  
Title : Crystal structure of the *M. tuberculosis* CTP synthase PyrG (apo form)  
Authors : Bellinzoni, M.; Barilone, N.; Alzari, P.M.  
Deposited on : 2015-04-17  
Resolution : 3.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1161 (3.60-3.44)
Clashscore	141614	1244 (3.60-3.44)
Ramachandran outliers	138981	1206 (3.60-3.44)
Sidechain outliers	138945	1207 (3.60-3.44)
RSRZ outliers	127900	1080 (3.60-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	
1	B	587	
1	C	587	
1	D	587	
1	E	587	
1	F	587	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	587	 <p>% 86% 10%</p>
1	H	587	 <p>% 85% 11%</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 30937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CTP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	531	3877	2451	676	739	11	0	0	0
1	B	534	3888	2458	671	747	12	0	0	0
1	C	533	3856	2442	662	740	12	0	0	0
1	D	531	3932	2490	686	745	11	0	0	0
1	E	530	3911	2479	684	738	10	0	0	0
1	F	526	3862	2447	669	735	11	0	0	0
1	G	527	3893	2466	678	738	11	0	0	0
1	H	525	3710	2338	641	720	11	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P9WHK7
B	0	GLY	-	expression tag	UNP P9WHK7
C	0	GLY	-	expression tag	UNP P9WHK7
D	0	GLY	-	expression tag	UNP P9WHK7
E	0	GLY	-	expression tag	UNP P9WHK7
F	0	GLY	-	expression tag	UNP P9WHK7
G	0	GLY	-	expression tag	UNP P9WHK7
H	0	GLY	-	expression tag	UNP P9WHK7

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	H	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CTP synthase

Chain A: 





- Molecule 1: CTP synthase

Chain B: 



- Molecule 1: CTP synthase

Chain C: 






- Molecule 1: CTP synthase





Chain D: 





• Molecule 1: CTP synthase




• Molecule 1: CTP synthase




• Molecule 1: CTP synthase



• Molecule 1: CTP synthase







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	196.83Å 196.83Å 184.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 3.52 49.21 – 3.52	Depositor EDS
% Data completeness (in resolution range)	98.6 (49.21-3.52) 98.3 (49.21-3.52)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 3.48Å)	Xtrriage
Refinement program	BUSTER 2.10.1	Depositor
R, $R_{free}$	0.213 , 0.221 0.220 , 0.231	Depositor DCC
$R_{free}$ test set	4292 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	101.7	Xtrriage
Anisotropy	0.721	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 80.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.089 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	30937	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	110.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/3965	0.54	0/5425
1	B	0.40	0/3975	0.54	0/5440
1	C	0.40	0/3941	0.54	0/5400
1	D	0.40	0/4019	0.54	0/5494
1	E	0.40	0/3999	0.53	0/5470
1	F	0.40	0/3948	0.53	0/5401
1	G	0.40	0/3977	0.53	0/5433
1	H	0.40	0/3791	0.53	0/5204
All	All	0.40	0/31615	0.53	0/43267

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3877	0	3650	9	0
1	B	3888	0	3655	7	0
1	C	3856	0	3615	11	0
1	D	3932	0	3774	9	0
1	E	3911	0	3743	13	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3862	0	3668	9	0
1	G	3893	0	3749	8	0
1	H	3710	0	3377	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
All	All	30937	0	29231	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:54:ASP:HB2	1:H:110:ARG:HH22	1.60	0.67
1:B:428:THR:HB	1:B:444:GLY:HA3	1.82	0.61
1:C:305:ALA:HB2	1:C:358:VAL:HG11	1.83	0.60
1:G:305:ALA:HB2	1:G:358:VAL:HG11	1.83	0.60
1:E:305:ALA:HB2	1:E:358:VAL:HG11	1.83	0.59
1:D:305:ALA:HB2	1:D:358:VAL:HG11	1.83	0.59
1:F:305:ALA:HB2	1:F:358:VAL:HG11	1.83	0.59
1:H:305:ALA:HB2	1:H:358:VAL:HG11	1.83	0.58
1:B:305:ALA:HB2	1:B:358:VAL:HG11	1.87	0.57
1:G:313:LEU:HD23	1:G:316:ALA:HB2	1.87	0.55
1:D:313:LEU:HD23	1:D:316:ALA:HB2	1.89	0.55
1:B:22:LEU:HD22	1:B:188:LEU:HD13	1.89	0.54
1:E:54:ASP:HB2	1:H:110:ARG:NH2	2.24	0.53
1:E:313:LEU:HD23	1:E:316:ALA:HB2	1.91	0.53
1:C:207:VAL:HG11	1:C:240:CYS:HB3	1.91	0.51
1:H:312:GLU:H	1:H:339:TRP:HB3	1.75	0.51
1:C:228:VAL:HG13	1:C:232:LEU:HD23	1.93	0.51
1:E:347:THR:HG23	1:E:350:GLY:H	1.77	0.49
1:G:347:THR:HG23	1:G:350:GLY:H	1.77	0.49
1:D:347:THR:HG23	1:D:350:GLY:H	1.78	0.49
1:A:428:THR:HB	1:A:444:GLY:HA3	1.95	0.49
1:C:347:THR:HG23	1:C:350:GLY:H	1.78	0.49

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:347:THR:HG23	1:F:350:GLY:H	1.77	0.49
1:B:347:THR:HG23	1:B:350:GLY:H	1.77	0.49
1:D:207:VAL:HG11	1:D:240:CYS:HB3	1.95	0.48
1:H:313:LEU:HD23	1:H:316:ALA:HB2	1.95	0.48
1:H:347:THR:HG23	1:H:350:GLY:H	1.77	0.48
1:A:347:THR:HG23	1:A:350:GLY:H	1.77	0.47
1:F:207:VAL:HG11	1:F:240:CYS:HB3	1.95	0.47
1:F:48:ASP:HB2	1:F:97:THR:HG22	1.97	0.47
1:A:121:VAL:HA	1:A:125:ILE:HB	1.96	0.47
1:G:207:VAL:HG11	1:G:240:CYS:HB3	1.95	0.47
1:H:207:VAL:HG11	1:H:240:CYS:HB3	1.98	0.46
1:D:48:ASP:HB2	1:D:97:THR:HG22	1.99	0.45
1:B:207:VAL:HG11	1:B:240:CYS:HB3	1.99	0.45
1:H:48:ASP:HB2	1:H:97:THR:HG22	1.99	0.45
1:C:497:GLY:HA3	1:C:507:PHE:HB2	1.99	0.45
1:C:389:VAL:HB	1:C:518:VAL:HG13	1.98	0.45
1:C:312:GLU:H	1:C:339:TRP:HB3	1.81	0.44
1:A:451:TYR:HB2	1:A:474:HIS:CD2	2.54	0.43
1:D:451:TYR:HB2	1:D:474:HIS:CD2	2.54	0.43
1:G:482:ASN:HA	1:G:485:ARG:HG3	2.01	0.43
1:A:482:ASN:HA	1:A:485:ARG:HG3	2.01	0.43
1:H:451:TYR:HB2	1:H:474:HIS:CD2	2.54	0.43
1:H:482:ASN:HA	1:H:485:ARG:HG3	2.01	0.43
1:G:392:LEU:HD22	1:G:525:PRO:HD2	2.01	0.42
1:D:392:LEU:HD22	1:D:525:PRO:HD2	2.01	0.42
1:E:451:TYR:HB2	1:E:474:HIS:CD2	2.53	0.42
1:F:482:ASN:HA	1:F:485:ARG:HG3	2.00	0.42
1:D:482:ASN:HA	1:D:485:ARG:HG3	2.01	0.42
1:E:392:LEU:HD22	1:E:525:PRO:HD2	2.01	0.42
1:G:451:TYR:HB2	1:G:474:HIS:CD2	2.53	0.42
1:A:207:VAL:HG11	1:A:240:CYS:HB3	2.01	0.42
1:E:482:ASN:HA	1:E:485:ARG:HG3	2.01	0.42
1:F:392:LEU:HD22	1:F:525:PRO:HD2	2.02	0.42
1:A:392:LEU:HD22	1:A:525:PRO:HD2	2.01	0.42
1:C:328:GLY:HA2	1:C:542:VAL:HG11	2.01	0.42
1:B:482:ASN:HA	1:B:485:ARG:HG3	2.01	0.42
1:F:451:TYR:HB2	1:F:474:HIS:CD2	2.54	0.42
1:E:248:ILE:HG13	1:E:268:LEU:HD13	2.02	0.41
1:F:248:ILE:HG13	1:F:268:LEU:HD13	2.02	0.41
1:G:248:ILE:HG13	1:G:268:LEU:HD13	2.02	0.41
1:A:328:GLY:HA2	1:A:542:VAL:HG11	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:248:ILE:HG13	1:B:268:LEU:HD13	2.02	0.41
1:D:248:ILE:HG13	1:D:268:LEU:HD13	2.02	0.41
1:E:47:LEU:HD23	1:E:96:VAL:HB	2.01	0.41
1:C:248:ILE:HG13	1:C:268:LEU:HD13	2.03	0.41
1:C:392:LEU:HD22	1:C:525:PRO:HD2	2.02	0.41
1:A:248:ILE:HG13	1:A:268:LEU:HD13	2.03	0.41
1:F:121:VAL:O	1:F:125:ILE:HB	2.21	0.41
1:H:392:LEU:HD22	1:H:525:PRO:HD2	2.02	0.41
1:E:207:VAL:HG11	1:E:240:CYS:HB3	2.02	0.41
1:E:328:GLY:HA2	1:E:542:VAL:HG11	2.04	0.40
1:E:58:MET:HA	1:H:116:GLY:HA2	2.02	0.40
1:C:482:ASN:HA	1:C:485:ARG:HG3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/587 (90%)	510 (97%)	15 (3%)	2 (0%)	34	71
1	B	530/587 (90%)	511 (96%)	16 (3%)	3 (1%)	25	64
1	C	529/587 (90%)	507 (96%)	18 (3%)	4 (1%)	19	59
1	D	527/587 (90%)	511 (97%)	13 (2%)	3 (1%)	25	64
1	E	526/587 (90%)	508 (97%)	15 (3%)	3 (1%)	25	64
1	F	520/587 (89%)	504 (97%)	14 (3%)	2 (0%)	34	71
1	G	521/587 (89%)	507 (97%)	12 (2%)	2 (0%)	34	71
1	H	519/587 (88%)	504 (97%)	13 (2%)	2 (0%)	34	71
All	All	4199/4696 (89%)	4062 (97%)	116 (3%)	21 (0%)	29	67

All (21) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	393	CYS
1	B	18	VAL
1	B	393	CYS
1	C	18	VAL
1	C	393	CYS
1	D	393	CYS
1	E	393	CYS
1	F	393	CYS
1	G	393	CYS
1	H	393	CYS
1	A	18	VAL
1	D	18	VAL
1	E	18	VAL
1	C	408	GLY
1	F	115	LEU
1	G	115	LEU
1	B	115	LEU
1	C	115	LEU
1	H	115	LEU
1	D	115	LEU
1	E	115	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	382/480 (80%)	377 (99%)	5 (1%)	69	86
1	B	384/480 (80%)	379 (99%)	5 (1%)	69	86
1	C	378/480 (79%)	372 (98%)	6 (2%)	62	83
1	D	397/480 (83%)	391 (98%)	6 (2%)	65	84
1	E	392/480 (82%)	389 (99%)	3 (1%)	81	91
1	F	386/480 (80%)	379 (98%)	7 (2%)	59	81
1	G	394/480 (82%)	389 (99%)	5 (1%)	69	86
1	H	350/480 (73%)	346 (99%)	4 (1%)	73	87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	3063/3840 (80%)	3022 (99%)	41 (1%)	69 86

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LYS
1	A	77	LEU
1	A	111	ARG
1	A	223	ARG
1	A	315	ASP
1	B	24	LYS
1	B	77	LEU
1	B	115	LEU
1	B	223	ARG
1	B	315	ASP
1	C	24	LYS
1	C	77	LEU
1	C	115	LEU
1	C	117	ASP
1	C	223	ARG
1	C	471	SER
1	D	24	LYS
1	D	54	ASP
1	D	77	LEU
1	D	115	LEU
1	D	197	GLU
1	D	223	ARG
1	E	24	LYS
1	E	77	LEU
1	E	223	ARG
1	F	24	LYS
1	F	77	LEU
1	F	122	ILE
1	F	199	LYS
1	F	223	ARG
1	F	313	LEU
1	F	315	ASP
1	G	24	LYS
1	G	77	LEU
1	G	117	ASP
1	G	199	LYS
1	G	223	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	H	77	LEU
1	H	144	ASN
1	H	199	LYS
1	H	223	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	171	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	531/587 (90%)	0.03	1 (0%) 95 91	68, 97, 122, 166	0
1	B	534/587 (90%)	0.01	3 (0%) 89 81	69, 97, 125, 159	0
1	C	533/587 (90%)	0.03	2 (0%) 92 87	68, 98, 133, 163	0
1	D	531/587 (90%)	0.08	1 (0%) 95 91	80, 107, 132, 174	0
1	E	530/587 (90%)	0.09	5 (0%) 84 73	87, 116, 139, 173	0
1	F	526/587 (89%)	0.10	7 (1%) 77 65	92, 118, 141, 168	0
1	G	527/587 (89%)	0.13	4 (0%) 86 75	96, 121, 142, 164	0
1	H	525/587 (89%)	0.00	3 (0%) 89 81	96, 120, 149, 188	0
All	All	4237/4696 (90%)	0.06	26 (0%) 89 81	68, 111, 140, 188	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	370	ARG	3.5
1	E	366	GLY	3.4
1	E	194	PRO	3.1
1	F	194	PRO	2.7
1	B	194	PRO	2.7
1	G	193	ALA	2.7
1	F	531	PRO	2.5
1	E	200	THR	2.5
1	G	324	LEU	2.4
1	B	366	GLY	2.3
1	B	195	SER	2.3
1	G	306	LEU	2.3
1	H	505	VAL	2.3
1	C	369	ILE	2.2
1	H	193	ALA	2.2
1	F	171	GLN	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	250	THR	2.2
1	F	372	ILE	2.2
1	C	248	ILE	2.1
1	E	493	LEU	2.1
1	F	370	ARG	2.1
1	A	194	PRO	2.0
1	F	392	LEU	2.0
1	E	542	VAL	2.0
1	G	31	LEU	2.0
1	H	250	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	G	601	1/1	0.16	0.23	112,112,112,112	0
2	CA	D	601	1/1	0.55	0.18	119,119,119,119	0
2	CA	F	601	1/1	0.60	0.25	124,124,124,124	0
2	CA	H	601	1/1	0.64	0.21	117,117,117,117	0
2	CA	B	601	1/1	0.68	0.27	97,97,97,97	0
2	CA	C	601	1/1	0.69	0.31	113,113,113,113	0
2	CA	E	601	1/1	0.78	0.27	114,114,114,114	0
2	CA	A	601	1/1	0.81	0.27	94,94,94,94	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.