

wwPDB X-ray Structure Validation Summary Report (i)

Apr 11, 2023 – 02:54 pm BST

PDB ID : 7ZD0

Title: Crystal structure of Pseudomonas aeruginosa S-adenosyl-L-homocysteine

hydrolase inhibited by Cd2+ ions

Authors : Malecki, P.H.; Gawel, M.; Brzezinski, K.

Deposited on : 2022-03-29

Resolution : 1.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.32.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.32.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	472	93%	
1	В	472	93%	
1	С	472	93%	
1	D	472	92%	5% •



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 16326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Adenosylhomocysteinase.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Λ	461	Total	С	N	О	S	0	4	0
1	A	401	3579	2256	618	682	23	0	4	0
1	В	460	Total	С	N	О	S	0	3	0
1	Б	400	3567	2249	617	679	22	0		
1	С	461	Total	С	N	О	S	0	6	0
1		401	3592	2265	622	681	24	U	0	
1	D	461	Total	С	N	О	S	0	6	0
1	ש	401	3591	2262	621	685	23	U	0	U

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9I685
A	-1	ASN	-	expression tag	UNP Q9I685
A	0	ALA	-	expression tag	UNP Q9I685
В	-2	SER	-	expression tag	UNP Q9I685
В	-1	ASN	ı	expression tag	UNP Q9I685
В	0	ALA	-	expression tag	UNP Q9I685
С	-2	SER	ı	expression tag	UNP Q9I685
С	-1	ASN	-	expression tag	UNP Q9I685
С	0	ALA	-	expression tag	UNP Q9I685
D	-2	SER	ı	expression tag	UNP Q9I685
D	-1	ASN	-	expression tag	UNP Q9I685
D	0	ALA	-	expression tag	UNP Q9I685


• Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0	
2	A	1	44	21	7	14	2	U		
9	В	1	Total	С	N	О	Р	0	0	
2	Б	1	44	21	7	14	2			
2	С	1	Total	С	N	О	Р	0	0	
2		1	44	21	7	14	2		U	
2	D	1	Total	С	N	О	Р	0	0	
2	ט	$D \mid I \mid$	44	21	7	14	2	U		


 \bullet Molecule 3 is ADENOSINE (three-letter code: ADN) (formula: $C_{10}H_{13}N_5O_4)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	Λ	1	Total C N	О	0	0
3	А	1	19 10 5	4	0	0
3	B	1	Total C N	О	0	0
3	Ъ	1	19 10 5	4		
3	С	1	Total C N	О	0	0
3	C	1	19 10 5	4	0	0
3	D	1	Total C N	О	0	0
3	D	1	19 10 5	4		U


 \bullet Molecule 4 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: $\mathrm{C_4H_{10}O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 4 2	0	0
4	A	1	Total C O 6 4 2	0	0
4	С	1	Total C O 6 4 2	0	0
4	С	1	Total C O 6 4 2	0	0
4	С	1	Total C O 6 4 2	0	0
4	D	1	Total C O 6 4 2	0	0

• Molecule 5 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: $C_3H_8O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 5 3 2	0	0
5	A	1	Total C O 5 3 2	0	0

• Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf			
6	A	1	Total K	0	0			
	11	_	1 1	Ŭ	Ŭ			
6	B	1	Total K	0	0			
	Б	В	D	Ъ	1	1 1		
6	С	1	Total K	0	0			
0		C	1	1 1	0	0		
6	D	1	Total K	0	0			
U	D	$D \mid I \mid$	1 1	U	U			

• Molecule 7 is CADMIUM ION (three-letter code: CD) (formula: Cd) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	5	Total Cd 5 5	0	0
7	В	5	Total Cd 6 6	0	1


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	С	4	Total Cd 5 5	0	1
7	D	5	Total Cd 5 5	0	0

• Molecule 8 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	В	1	Total 8	C 6	O 2	0	0

• Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	D	1	Total Cl 1 1	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	488	Total O 497 497	0	9
10	В	365	Total O 374 374	0	11
10	С	407	Total O 414 414	0	10

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	D	376	Total O 380 380	0	5




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenosylhomocysteinase









4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	176.97Å 134.16Å 108.88Å	Depositor
a, b, c, α , β , γ	90.00° 106.06° 90.00°	Depositor
Resolution (Å)	44.10 - 1.87	Depositor
recondition (11)	44.10 - 1.87	EDS
% Data completeness	92.5 (44.10-1.87)	Depositor
(in resolution range)	93.3 (44.10-1.87)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.81 (at 1.87Å)	Xtriage
Refinement program	PHENIX 1.19.2	Depositor
D.D.	0.147 , 0.162	Depositor
R, R_{free}	0.147 , 0.160	DCC
R_{free} test set	1029 reflections (0.55%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 45.3	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
	0.000 for -1/2 *h + 1/2 *k + 1,1/2 *h - 1/2 *k + 1,1	
Estimated twinning fraction	/2*h+1/2*k 0.000 for -1/2*h-1/2*k+l,-1/2*h-1/2*k-l,1/2	Xtriage
G		
E. E. convolution	*h-1/2*k	EDC
F_o, F_c correlation	0.98	EDS
Total number of atoms	16326	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NAD, BU1, K, PDO, HEZ, ADN, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.40	0/3648	0.59	1/4931~(0.0%)
1	В	0.36	0/3636	0.58	1/4916~(0.0%)
1	С	0.37	0/3667	0.59	0/4955
1	D	0.36	0/3660	0.57	1/4947~(0.0%)
All	All	0.37	0/14611	0.58	3/19749 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	373	LEU	CA-CB-CG	5.71	128.43	115.30
1	D	373	LEU	CA-CB-CG	5.12	127.06	115.30
1	A	373	LEU	CA-CB-CG	5.03	126.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3579	0	3584	12	0
1	В	3567	0	3575	14	0
1	С	3592	0	3608	11	0
1	D	3591	0	3592	16	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
2	A	44	0	26	1	0
2	В	44	0	26	1	0
2	С	44	0	26	1	0
2	D	44	0	26	1	0
3	A	19	0	13	2	0
3	В	19	0	13	2	0
3	С	19	0	13	2	0
3	D	19	0	13	2	0
4	A	12	0	20	2	0
4	С	18	0	30	1	0
4	D	6	0	10	1	0
5	A	10	0	16	1	0
6	A	1	0	0	0	0
6	В	1	0	0	0	0
6	С	1	0	0	0	0
6	D	1	0	0	0	0
7	A	5	0	0	0	0
7	В	6	0	0	0	0
7	С	5	0	0	0	0
7	D	5	0	0	0	0
8	В	8	0	14	1	0
9	D	1	0	0	0	0
10	A	497	0	0	2	0
10	В	374	0	0	3	0
10	С	414	0	0	0	0
10	D	380	0	0	2	0
All	All	16326	0	14605	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:14:LYS:HE2	4:C:503:BU1:H21	1.70	0.72
1:D:14:LYS:HE3	4:D:503:BU1:H32	1.76	0.67
1:B:287:LEU:HD12	10:B:781:HOH:O	1.96	0.65
1:A:148:HIS:HB3	4:A:504:BU1:H41	1.84	0.58
2:A:501:NAD:C4N	3:A:502:ADN:H3'	2.35	0.56

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/472 (98%)	452 (98%)	10 (2%)	1 (0%)	47	37
1	В	461/472 (98%)	453 (98%)	8 (2%)	0	100	100
1	С	465/472~(98%)	455 (98%)	9 (2%)	1 (0%)	47	37
1	D	465/472 (98%)	455 (98%)	9 (2%)	1 (0%)	47	37
All	All	1854/1888 (98%)	1815 (98%)	36 (2%)	3 (0%)	47	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	С	61	HIS
1	D	61	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	381/385 (99%)	378 (99%)	3 (1%)	81	80
1	В	380/385 (99%)	379 (100%)	1 (0%)	92	92
1	С	383/385 (100%)	381 (100%)	2 (0%)	88	88
1	D	382/385~(99%)	379 (99%)	3 (1%)	81	80
All	All	1526/1540 (99%)	1517 (99%)	9 (1%)	86	86

5 of 9 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	D	373	LEU
1	D	409	LEU
1	В	373	LEU
1	С	42	ARG
1	С	335	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	129	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 43 ligands modelled in this entry, 26 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type C	Chain	Chain	n Ros	Pog	Pag	Dog	Dec	Dog	Pos	Ros	Dog	Dog	Dog	Dog	Dog	Dog	Dag	Des	Dec	Dog	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	gles
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2																		
4	BU1	D	503	-	5,5,5	0.39	0	4,4,4	0.40	0																		
3	ADN	С	502	-	18,21,21	0.66	0	18,31,31	0.84	1 (5%)																		
4	BU1	A	503	-	5,5,5	0.34	0	4,4,4	0.52	0																		



Mol	Type	Chain	Res	Link	Вс	ond leng	$ ag{ths}$	Bond angles			
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	PDO	A	505	-	4,4,4	0.36	0	3,3,3	0.36	0	
3	ADN	A	502	_	18,21,21	0.64	0	18,31,31	0.93	1 (5%)	
3	ADN	D	502	-	18,21,21	0.67	0	18,31,31	0.87	1 (5%)	
5	PDO	A	506	-	4,4,4	0.33	0	3,3,3	0.38	0	
2	NAD	С	501	-	42,48,48	0.57	0	50,73,73	0.71	1 (2%)	
4	BU1	A	504	-	5,5,5	0.33	0	4,4,4	0.60	0	
4	BU1	С	504	_	5,5,5	0.38	0	4,4,4	0.47	0	
2	NAD	A	501	-	42,48,48	0.55	0	50,73,73	0.78	1 (2%)	
4	BU1	С	503	-	5,5,5	0.32	0	4,4,4	0.44	0	
4	BU1	С	505	-	5,5,5	0.33	0	4,4,4	0.38	0	
8	HEZ	В	503	_	7,7,7	0.12	0	6,6,6	0.42	0	
2	NAD	D	501	-	42,48,48	0.55	0	50,73,73	0.72	1 (2%)	
3	ADN	В	502	-	18,21,21	0.64	0	18,31,31	0.89	1 (5%)	
2	NAD	В	501	-	42,48,48	0.54	0	50,73,73	0.74	1 (2%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BU1	D	503	-	-	2/3/3/3	-
3	ADN	С	502	-	-	0/2/22/22	0/3/3/3
4	BU1	A	503	-	-	1/3/3/3	-
5	PDO	A	505	_	-	2/2/2/2	-
3	ADN	A	502	-	-	0/2/22/22	0/3/3/3
3	ADN	D	502	-	-	0/2/22/22	0/3/3/3
5	PDO	A	506	_	-	1/2/2/2	-
2	NAD	С	501	_	-	5/26/62/62	0/5/5/5
4	BU1	A	504	-	-	2/3/3/3	-
4	BU1	С	504	-	-	2/3/3/3	-
2	NAD	A	501	-	-	5/26/62/62	0/5/5/5
4	BU1	С	503	-	-	2/3/3/3	-
4	BU1	С	505	-	-	0/3/3/3	-
8	HEZ	В	503	-	-	3/5/5/5	-
2	NAD	D	501	-	-	5/26/62/62	0/5/5/5
3	ADN	В	502	-		0/2/22/22	0/3/3/3
2	NAD	В	501	-	-	5/26/62/62	0/5/5/5



There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	502	ADN	C5-C6-N6	2.33	123.90	120.35
2	С	501	NAD	C5A-C6A-N6A	2.29	123.83	120.35
3	В	502	ADN	C5-C6-N6	2.27	123.81	120.35
2	В	501	NAD	C5A-C6A-N6A	2.27	123.80	120.35
2	A	501	NAD	C5A-C6A-N6A	2.26	123.79	120.35

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NAD	O4D-C1D-N1N-C2N
2	A	501	NAD	O4D-C1D-N1N-C6N
2	A	501	NAD	C2D-C1D-N1N-C2N
2	A	501	NAD	C2D-C1D-N1N-C6N
2	В	501	NAD	O4D-C1D-N1N-C2N

There are no ring outliers.


13 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	503	BU1	1	0
3	С	502	ADN	2	0
3	A	502	ADN	2	0
3	D	502	ADN	2	0
5	A	506	PDO	1	0
2	С	501	NAD	1	0
4	A	504	BU1	2	0
2	A	501	NAD	1	0
4	С	503	BU1	1	0
8	В	503	HEZ	1	0
2	D	501	NAD	1	0
3	В	502	ADN	2	0
2	В	501	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.




Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	461/472 (97%)	-0.06	8 (1%) 70 72	22, 32, 50, 73	0
1	В	460/472 (97%)	0.15	27 (5%) 22 24	23, 38, 64, 79	0
1	С	461/472 (97%)	-0.01	15 (3%) 46 47	25, 35, 54, 72	0
1	D	461/472 (97%)	0.27	38 (8%) 11 12	23, 37, 62, 76	0
All	All	1843/1888 (97%)	0.09	88 (4%) 30 32	22, 35, 59, 79	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	10	PHE	7.5
1	В	409	LEU	5.8
1	D	412	ALA	5.6
1	A	355	ASP	5.4
1	D	411	ALA	5.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathrm{\AA}^2)$	Q < 0.9
4	BU1	С	505	6/6	0.59	0.19	54,62,66,71	0
4	BU1	С	504	6/6	0.70	0.21	54,60,70,76	0
4	BU1	D	503	6/6	0.75	0.17	55,61,68,81	0
4	BU1	A	503	6/6	0.76	0.09	49,57,60,64	0
5	PDO	A	506	5/5	0.84	0.22	52,55,68,69	0
8	HEZ	В	503	8/8	0.85	0.11	52,58,69,71	0
7	CD	С	510	1/1	0.88	0.11	98,98,98,98	1
5	PDO	A	505	5/5	0.89	0.08	49,52,61,61	0
4	BU1	A	504	6/6	0.90	0.12	50,58,69,73	0
7	CD	A	511	1/1	0.90	0.09	100,100,100,100	1
4	BU1	С	503	6/6	0.91	0.09	51,54,64,71	0
7	CD	D	509	1/1	0.92	0.06	69,69,69,69	1
7	CD	В	506	1/1	0.92	0.05	92,92,92,92	1
7	CD	С	508	1/1	0.93	0.09	99,99,99,99	1
3	ADN	D	502	19/19	0.95	0.09	28,31,37,38	0
2	NAD	С	501	44/44	0.95	0.09	25,29,33,38	0
3	ADN	С	502	19/19	0.95	0.11	25,28,36,39	0
2	NAD	В	501	44/44	0.96	0.08	26,29,31,33	0
7	CD	A	512	1/1	0.96	0.04	91,91,91,91	1
2	NAD	A	501	44/44	0.96	0.10	24,27,29,30	0
2	NAD	D	501	44/44	0.96	0.08	24,28,31,37	0
3	ADN	A	502	19/19	0.96	0.10	21,25,30,31	0
3	ADN	В	502	19/19	0.96	0.11	27,30,34,39	0
7	CD	A	510	1/1	0.96	0.07	73,73,73,73	1
6	K	D	504	1/1	0.97	0.09	38,38,38,38	1
7	CD	D	507	1/1	0.97	0.12	81,81,81,81	1
7	CD	С	509[A]	1/1	0.98	0.05	55,55,55,55	1
7	CD	С	509[B]	1/1	0.98	0.05	81,81,81,81	1
7	CD	В	507[A]	1/1	0.98	0.07	58,58,58,58	1
7	CD	В	507[B]	1/1	0.98	0.07	74,74,74,74	1
7	CD	D	508	1/1	0.98	0.07	66,66,66,66	1
7	CD	В	509	1/1	0.98	0.04	54,54,54,54	1
6	K	В	504	1/1	0.98	0.10	41,41,41,41	1
7	CD	В	508	1/1	0.99	0.03	96,96,96,96	1
6	K	С	506	1/1	0.99	0.10	37,37,37,37	1
7	CD	A	509	1/1	0.99	0.05	49,49,49,49	1
7	CD	D	505	1/1	0.99	0.04	44,44,44,44	1
7	CD	D	506	1/1	1.00	0.05	36,36,36,36	1
6	K	A	507	1/1	1.00	0.07	39,39,39,39	1
7	CD	A	508	1/1	1.00	0.05	31,31,31,31	0
7	CD	С	507	1/1	1.00	0.07	37,37,37,37	1
7	CD	В	505	1/1	1.00	0.06	39,39,39,39	1
9	CL	D	510	1/1	1.00	0.10	32,32,32,32	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around CD A 511:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CD D 509: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around CD B 506: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around CD C 508:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ADN D 502:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD C 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around ADN C 502: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around NAD B 501: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)





Electron density around CD A 512: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around NAD A 501:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAD D 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ADN A 502:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ADN B 502:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around CD A 510:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around K D 504: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around CD D 507: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around CD B 507 (A):

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CD B 507 (B): $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around CD D 508:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



























Electron density around CD A 509: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around CD D 505: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)
























6.5 Other polymers (i)

There are no such residues in this entry.

